

Fields and Galois Theory

Lectures by Uzi Vishne
Summary by Ari Feiglin (ari.feiglin@gmail.com)

Contents

1 Field Extensions

1

1 Field Extensions

Suppose $F \subseteq K$ are fields, then K is certainly also an F -vector space and therefore has a dimension and we denote it $[K : F] := \dim_F K$.

1.0.1 Theorem

Suppose $F \subseteq K$ and V is a K -vector space, then V is also a vector space over F as well, and $\dim_F V = [K : F] \dim_K V$.

Proof: Let $B_1 \subseteq V$ be a basis for V over K and $B_2 \subseteq K$ be a basis for K over F , then define $B = \{\alpha v \mid \alpha \in B_2, v \in B_1\}$. This is a basis for V in F , it is linearly independent since if $\alpha_1 v_1, \dots, \alpha_n v_n \in B$ and $\beta_1, \dots, \beta_n \in F$ then $\sum_{i=1}^n \beta_i \alpha_i v_i = 0$ implies $\beta_i \alpha_i = 0$ for all i since B_1 is a basis, and this means that β_i or α_i is zero, but $\alpha_i v_i \in B$ so $\beta_i = 0$ as required. B spans V since for $v \in B$ there exist $v_1, \dots, v_n \in B_1$ and $\alpha_1, \dots, \alpha_n \in K$ such that $v = \sum_{i=1}^n \alpha_i v_i$ and α_i can be written as the linear combination of elements in B_2 by elements of F which gives a linear combination of elements in B of F . So B is indeed a basis for V over F . Finally $B \cong B_2 \times B_1$ since $(\alpha, v) \mapsto \alpha v$ is a bijection: it is obviously surjective and $\alpha_1 v_1 = \alpha_2 v_2$ implies $\alpha_1 = \alpha_2, v_1 = v_2$ since v_1, v_2 are independent. Thus we have

$$\dim_F V = |B| = |B_2 \times B_1| = [K : F] \dim_K V$$

In particular if $F \subseteq K \subseteq E$ are fields then $[E : F] = [E : K] \cdot [K : F]$.

The following are methods of constructing fields:

- (1) If R is a commutative ring and $M \triangleleft R$ is a maximal ideal then R/M is a field. Specifically if $R = F[x]$ and p is an irreducible polynomial, $\langle p \rangle$ is maximal and $F[x]/\langle p \rangle$ is a field.
- (2) If F is a field, then the set of rational functions is also a field:

$$F \subseteq F(x) := \left\{ \frac{f(x)}{g(x)} \mid f, g \in F[x], g(x) \neq 0 \right\}$$

In general if R is an integral domain then its field of fractions/quotients $q(R) := \left\{ \frac{a}{b} \mid a, b \in R, b \neq 0 \right\}$ is a field. And $F(x)$ is the quotient field of $F[x]$.

- (3) If $F_0 \subseteq F_1 \subseteq F_2 \subseteq \dots$ is a chain of fields then so is $\bigcup F_n$ (the theory of fields is inductive, this holds for arbitrary chains, not just inductive ones). So for example $F(\lambda_1, \lambda_2, \dots)$ is a field since we can define $F_n = F(\lambda_1, \dots, \lambda_n)$ (the quotient field of $F[\lambda_1, \dots, \lambda_n]$) and the union of this chain is $F(\lambda_1, \lambda_2, \dots)$.

Let F be a field and $F \subseteq K$ a ring with $a \in K$, we define a homomorphism $F[\lambda] \xrightarrow{\psi_a} K$ defined by $\alpha \mapsto \alpha$ for $\alpha \in F$ and $\lambda \mapsto a$, meaning

$$\psi_a \left(\sum \alpha_i \lambda^i \right) = \sum \alpha_i a^i \quad (\psi_a(f) = f(a))$$

In particular ψ_a is a linear transformation from F to K , and is called the *evaluation homomorphism* at a . The kernel of the homomorphism is

$$\ker \psi_a = \{f \in F[\lambda] \mid f(a) = 0\} \triangleleft F[\lambda]$$

1.0.2 Definition

$a \in K$ is **algebraic** if $\ker \psi_a \neq 0$ and **transcendental** if the kernel is trivial.

If a is transcendental then $\ker \psi_a = 0$ and so $\text{Im } \psi_a = \{f(a) \mid f \in F[\lambda]\} = F[a] \cong F[\lambda]$. In fact we get

$$\begin{array}{ccc} F & \subseteq & F[a] \subseteq F(x) \\ & \cong & \cong \\ & & F[\lambda] \end{array}$$

Now if a is algebraic, since $F[x]$ is a euclidean domain and therefore a PID, the kernel has a generator $\ker \psi_a = \langle h \rangle = h \cdot F[\lambda]$. So $h(a) = 0$ and $f(a) = 0 \implies h \mid f$, and h is called the *minimal polynomial* of a . And so

$$F[\lambda] / \langle h \rangle = F[\lambda] / \ker \psi_a \cong \text{Im } \psi_a = \{f(a) \mid f \in F[\lambda]\} = F[a] = \text{span}\{1, a, \dots, a^{n-1}\} \subseteq K$$

2 Field Extensions

where $n = \deg h$, since $f(x) = q(x)h(x) + r(x)$ where $\deg r < \deg h = n$ and so $f(a) = r(a)$. $\{1, \dots, a^{n-1}\}$ is a basis due to h being minimal, a zeroing linear combination would give a zeroing polynomial of a of degree less than h . This means that the dimension of $F[a]$ as an F -vector space is n , ie. $[F[a] : F] = n$.

Since K is an integral domain and therefore so too is $F[a]$ and this means that $\langle h \rangle$ is a prime ideal (since R/I is an integral domain if and only if I is prime), this means that h is a prime (irreducible) polynomial. And since $F[a]$ is a PID, prime and maximal ideals are one and the same, so $\langle h \rangle$ is maximal and therefore $F[a]/\langle h \rangle \cong F[a]$ is a field. Let us summarize this:

1.0.3 Proposition

Let $F \subseteq K$ where K is an integral domain and $a \in K$ is algebraic in F , let h_a be its minimal polynomial. Then (1) h_a is irreducible, (2) $F[a]$ is a field, (3) $[F[a] : F] = \deg h_a$.

So for example let $a \in K \setminus F$ be algebraic then $F \subseteq F[a] \subseteq K$ and suppose $[K : F] = p$ is prime. Then $p = [K : F] = [K : F[a]] \cdot [F[a] : F]$, and since $a \in F[a] \setminus F$ this means $[F[a] : F] > 1$ so $[F[a] : F] = p$ and $[K : F[a]] = 1$ since p is prime so $F[a] = K$.

1.0.4 Corollary

Suppose F is a field and $F \subseteq K$ is an integral domain with finite dimension. Then every element of K is algebraic and K is a field.

Proof: Let $a \in K$ then $[K : F] = [K : F[a]] \cdot [F[a] : F]$ so $[F[a] : F]$ is finite. If a were transcendental then $F[a] \cong F[x]$ and $F[x]$ has infinite dimension over F . K is a field since every $a \in K$ must have a multiplicative inverse, since $F[a]$ is a field. ■

Notice that $[F[a, b] : F[a]] \leq [F[b] : F]$ since if h_b is b 's minimal polynomial in F then it is also a zeroing polynomial in $F[a]$. This means that

$$[F[a, b] : F] = [F[a, b] : F[a]] \cdot [F[a] : F] \leq [F[b] : F] \cdot [F[a] : F]$$

1.0.5 Corollary

Let F be a field and K a field extension, define

$$\text{Alg}_F(K) := \{a \in K \mid a \text{ is algebraic over } F\}.$$

This is a field. Furthermore $F \subseteq \text{Alg}_F(K)$ is an algebraic extension (all elements of $\text{Alg}_F(K)$ are algebraic in F), and $\text{Alg}_F(K) \subseteq K$ is a purely transcendental extension (all elements in $K \setminus \text{Alg}_F(K)$ are transcendental in $\text{Alg}_F(K)$).

Proof: Notice that $F[a \cdot b], F[a + b] \subseteq F[a, b]$ and so $[F[a, b] : F] \leq [F[b] : F] \cdot [F[a] : F] < \infty$, so $\text{Alg}_F(K)$ is closed under addition and multiplication (and obviously additive inverses). For a algebraic, $F[a]$ is a field so $a^{-1} \in F[a]$ and so $F[a^{-1}] \subseteq F[a]$ and therefore $[F[a^{-1}] : F] < \infty$ so a^{-1} is algebraic as well (and so by symmetry $F[a] = F[a^{-1}]$). So $\text{Alg}_F(K)$ is indeed a field.

To show that $\text{Alg}_F(K) \subseteq K$ is a pure transcendental extension, notice that if $F_1 \subseteq F_2 \subseteq F_3$ where $F_1 \subseteq F_2$ is algebraic, if $a \in F_3$ is algebraic in F_2 it is also algebraic in F_1 . Indeed if $f \in F_2[x]$ such that $f(a) = 0$, let its coefficients be b_i then a is algebraic in $F_1[b_0, \dots, b_n]$ and so

$$[F_1[b_0, \dots, b_n, a] : F_1[b_0, \dots, b_n]] = [F_1[b_0, \dots, b_n, a] : F_1[b_0, \dots, b_n]] \cdot [F_1[b_0, \dots, b_n] : F_1]$$

and this is finite since b_0, \dots, b_n are algebraic in F_1 as they are in F_2 , so both terms are finite. So if K had any algebraic numbers not in $\text{Alg}_F(K)$, they would be algebraic in F and thus in $\text{Alg}_F(K)$ in contradiction. ■

1.0.6 Proposition

Let F be a field and $f \in F[\lambda]$ be irreducible, then there exists a field extension $F \subseteq K$ such that f has a root in K , and $[K : F] = \deg f$.

Proof: since f is irreducible, $\langle f \rangle$ is prime and $F[\lambda]$ is a PID so it is maximal. So $K := F[\lambda]/\langle f \rangle$ is a field, and its dimension is $\deg f$, since it can be generated by $\{1, x, \dots, x^{\deg f-1}\}$. Now recall that by the second isomorphism theorem, $F/F \cap \langle f \rangle \cong F + \langle f \rangle / \langle f \rangle \subseteq F[\lambda]/\langle f \rangle = K$. But since elements of $\langle f \rangle$ are multiples of f , which is disjoint from F , so $F \cap \langle f \rangle = (0)$ so $F/F \cap \langle f \rangle \cong F$, and so F can be embedded into K and is thus for all intents and purposes, a subfield of K . Now define $\alpha := \lambda + \langle f \rangle$, and suppose $f(\lambda) = \sum_{i=0}^n a_i \lambda^i$ where $a_i \in F$ (viewing f as a polynomial over K , a_i is actually $a_i + \langle f \rangle$). Then

$$f(\alpha) = \sum_{i=0}^n a_i (\lambda + \langle f \rangle)^i = \sum_{i=0}^n a_i (\lambda^i + \langle f \rangle) = \sum_{i=0}^n a_i \lambda^i + \langle f \rangle = f + \langle f \rangle = \langle f \rangle = 0_K$$

so α is indeed a root of $f(\lambda)$, as required. \blacksquare

1.0.7 Corollary

Let F be a field and $f \in F[\lambda]$ any polynomial. Then there exists a field extension $F \subseteq K$ such that f has a root in K and $[K : F] \leq \deg f$.

Proof: find f 's irreducible factorization $f = f_1 \cdots f_t$, then extend F to a field K such that f_1 has a root in K , and by above $[K : F] = \deg f_1 \leq \deg f$. \blacksquare

1.0.8 Definition

Let F be a field, and f a polynomial over F . A field $F \subseteq K$ **splits** f if there exist $\alpha_1, \dots, \alpha_n \in K$ such that $f(\lambda) = (\lambda - \alpha_1) \cdots (\lambda - \alpha_n)$.

1.0.9 Theorem

Every polynomial f over a field F has a field K which splits it, such that $[K : F] \leq (\deg f)!$.

Proof: by induction on $n = \deg f$. For $n = 1$ then f already has a root, and so take $F = K$ and $[K : F] = 1 = (\deg f)!$. Now suppose $\deg f = n + 1$, then by above there exists a field extension $F \subseteq K_0$ such that there exists an $\alpha_1 \in K_0$ such that $f(\alpha_1) = 0$ and $[K_0 : F] \leq \deg f = n + 1$. And so $(\lambda - \alpha_1) \mid f(\lambda)$, so $f(\lambda) = (\lambda - \alpha_1)g(\lambda)$. Then $\deg g = n$, and g is a polynomial over K_0 , so there exists a field extension $F \subseteq K_0 \subseteq K$ such that $g(\lambda) = (\lambda - \alpha_2) \cdots (\lambda - \alpha_{n+1})$ for $\alpha_i \in K$ and $[K : K_0] \leq n!$. Then $f(\lambda) = (\lambda - \alpha_1) \cdots (\lambda - \alpha_{n+1})$ for $\alpha_i \in K$ and $[K : F] = [K : K_0][K_0 : F] \leq (n + 1)n! = (n + 1)!$. \blacksquare

Notice the following

- (1) the split of a polynomial over any field into its roots is unique,
- (2) the number of roots is $\leq \deg f$.

Recall that a field F is *algebraically closed* if it splits every polynomial in $F[\lambda]$.

1.0.10 Definition

Let F be a field, then $F \subseteq \overline{F}$ is an **algebraic closure** of F if \overline{F} is algebraically closed.

Note

Every field has a unique (up to isomorphism) algebraic closure.

So let $f(\lambda) \in F[\lambda]$, then $f(\lambda) \in \overline{F}[\lambda]$ and so $f = (\lambda - \alpha_1) \cdots (\lambda - \alpha_n)$ for $\alpha_i \in \overline{F}$. Then take $F \subseteq K = F[\alpha_1, \dots, \alpha_n] \subseteq \overline{F}$, it can be shown that $[K : F] \leq (\deg f)!$.

Now suppose $F \subseteq K$ are fields, and E is a field which F is embeddable into, suppose $\varphi: F \hookrightarrow E$ is an embedding. An embedding $\varphi': K \hookrightarrow E$ is an *extension* of φ if $\varphi'|_F = \varphi$. Denote

$$\eta_{F \subseteq K}^E := \#\{\varphi' \text{ is an extension of } \varphi\}$$

where φ is held constant and understood. Then

1.0.11 Proposition

Suppose $K = F[\alpha]$, then $\eta_{F \subseteq K}^E$ is equal to the number of roots the minimal polynomial of α in F has in E .

Proof: since α generates K over F , every extension of φ is defined by its image on α . Let h be the minimal polynomial of α over F . Denote $\hat{b} := \varphi(b)$ for all $b \in F$, and this definition extends to polynomials, $\sum_{i=0}^n b_i x^i = \sum_{i=0}^n \hat{b}_i x^i$. Then if φ' is an extension of φ ,

$$\hat{h}(\varphi'(\alpha)) = \varphi'(h(\alpha)) = \varphi'(0) = 0$$

this is since if $h(\lambda) = \sum_{i=0}^n a_i \lambda^i$, then $\hat{h}(\lambda) = \sum_{i=0}^n \hat{a}_i \lambda^i$, so

$$\hat{h}(\varphi'(\alpha)) = \sum_{i=0}^n \hat{a}_i \varphi'(\alpha)^i = \sum_{i=0}^n \varphi(a_i) \varphi'(\alpha)^i = \sum_{i=0}^n \varphi'(a_i) \varphi'(\alpha)^i = \varphi' \left(\sum_{i=0}^n a_i \alpha^i \right) = \varphi'(h(\alpha))$$

so $\varphi'(\alpha)$ must be one of \hat{h} 's roots, precisely as stated. ■

1.0.12 Definition

A polynomial f which splits over E is called **separable** in E if its linear factors are distinct (ie. all of its roots in E are distinct).

1.0.13 Theorem

Let $F \subseteq K$ be a finite extension (meaning $[K : F] < \infty$), and let $\varphi: F \hookrightarrow E$ be a given embedding. Then

- (1) $\eta_{F \subseteq K}^E \leq [K : F]$,
- (2) if K is generated by the roots of f , assuming that E splits f , then $1 \leq \eta_{F \subseteq K}^E$,
- (3) if f is separable over E , then $\eta_{F \subseteq K}^E = [K : F]$.

Proof: suppose $K = F[\alpha_1, \dots, \alpha_n]$ (the generators of K can be taken to be the basis of K as an F -vector space). We prove this by induction on n , for $n = 1$ this is given by the previous proposition, since $\eta_{F \subseteq K}^E$ is the number of roots h has in E , and $[K : F] = \deg h$ which is at least this. Define $F_1 := F[\alpha_1]$, then

$$\begin{aligned} \eta_{F \subseteq K}^E &= \#\{\varphi'': K \longrightarrow E \text{ is an extension of } \varphi\} \\ &= \#\bigcup \{\varphi'': F_1 \longrightarrow E \text{ is an extension of } \varphi' \mid \varphi': F_1 \longrightarrow E \text{ is an extension of } \varphi\} \\ &= \sum_{\varphi'} \eta_{F_1 \subseteq K}^E = \eta_{F \subseteq F_1}^E \cdot \eta_{F_1 \subseteq K}^E \subseteq [F_1 : F] \cdot [K : F_1] = [K : F] \end{aligned}$$

For (2), by the assumption there is an extension of $F \hookrightarrow E$ to $F_1 \hookrightarrow E$, and continue inductively. For (3), since f is separable, makes the bound an equality. ■

1.0.14 Definition

Let f be a polynomial over F , a field $F \subseteq K$ is a **splitting field** if it is the smallest field in which the polynomial splits.