

Mathcord Mathematical Logic

Problem Set 1

Submission October 28

Problem 1

- (1) Show that $\{\neg, \rightarrow\}$ is functional complete.
- (2) Show that \vee can be represented in terms of \rightarrow alone.
- (3) Show that any formula φ over the signature \rightarrow has $w\varphi = 1$ for the valuation w which evaluates every variable as 1.
- (4) Show that \wedge cannot be represented in terms of \rightarrow alone.
- (5) Show that \leftrightarrow cannot be represented in terms of \rightarrow alone.

Problem 2

We define the boolean *nand* connective, \uparrow , as follows:

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

and *nor*, \downarrow :

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

- (1) Verify that $\alpha \uparrow \beta \equiv \neg(\alpha \wedge \beta)$ and $\alpha \downarrow \beta \equiv \neg(\alpha \vee \beta)$.
- (2) Show that both $\{\uparrow\}$ and $\{\downarrow\}$ are functional complete.
- (3) Show that if \circ is a boolean connective and $\{\circ\}$ is functional complete, $\circ \in \{\uparrow, \downarrow\}$.

Problem 3

Let $\varphi \rightarrow \psi$ be a tautology, then show that there exists a formula θ which uses only variables common to both φ and ψ (i.e. $\text{var}\theta \subseteq \text{var}\varphi \cap \text{var}\psi$), such that both $\varphi \rightarrow \theta$ and $\theta \rightarrow \psi$ are tautologies.

Problem 4

Let φ be a formula, we define its *dual*, φ^δ , recursively as follows:

$$\pi^\delta = \pi \text{ for prime } \pi, \quad (\neg\alpha)^\delta = \neg\alpha^\delta, \quad (\alpha \wedge \beta)^\delta = (\alpha^\delta \vee \beta^\delta), \quad (\alpha \vee \beta)^\delta = (\alpha^\delta \wedge \beta^\delta)$$

Equivalently, it just substitutes all \wedge in φ with \vee and all \vee with \wedge .

Let $f: \{0, 1\}^n \rightarrow \{0, 1\}$ be a boolean function, we define its *dual* to be

$$f^\delta \bar{x} = \neg f(\neg \bar{x})$$

- (1) Show that if α represents f , then α^δ represents f^δ .
- (2) Show that $f \mapsto f^\delta$ is a bijection on boolean functions.
- (3) Using the fact that every boolean function can be represented by a DNF, show that every boolean function can be represented by a CNF.

Problem 5

Let $\bar{x} = (x_1, \dots, x_n)$ and $\bar{y} = (y_1, \dots, y_n)$ be boolean vectors. We write $\bar{x} \leq \bar{y}$ to mean $x_i \leq y_i$ for all $i = 1, \dots, n$. A boolean function f is said to be *increasing* if $\bar{x} \leq \bar{y}$ implies $f\bar{x} \leq f\bar{y}$.

Show that every increasing boolean function can be represented in the logical signature $\{\wedge, \vee\}$ and vice versa: every formula in this logical signature is increasing.