









METHOD	FORMULA	CONDITIONS	QUANTITY REPRESENTED BY INTEGRAND	WHEN TO USE
CROSS-SECTION	$\int_a^b A(t) dt$	A IS CONTINUOUS ON $a \leq t \leq b$	AREA OF A CROSS-SECTION PERPENDICULAR TO $t$ -AXIS 	CROSS-SEC. IS TRIANGULAR, SQUARE, ETC. (NOT A DISC OR CYLINDER)
DISC	$\int_a^b \pi f(t)^2 dt$	f CONTINUOUS ON $a \leq t \leq b$	AREA OF A CROSS-SECTION PERPENDICULAR TO $t$ -AXIS 	CROSS-SEC. IS A DISC OF RADIUS $f(t)$ REGION BOUNDED BY $f$ & $t$ -AXIS IS REVOLVED @ $t$ -AXIS 
WASHER (SUBTRACTIVE DISC)	$\int_a^b \pi (f(t)^2 - g(t)^2) dt$	f, g CONT'S ON $a \leq t \leq b$ , $f(t) \geq g(t)$	AREA OF A CROSS-SECTION PERPENDICULAR TO $t$ -AXIS 	CROSS-SEC. IS A DISC OF RADIUS $f(t)$ WITH A HOLE OF RADIUS $g(t)$ REGION BDD. BY $f$ & $g$ IS REVOLVED @ $t$ -AXIS 
SHELL	$\int_a^b 2\pi t f(t) dt$	f CONT'S ON $a \leq t \leq b$	SURFACE AREA OF AN OPEN CYLINDER OF RADIUS $t$ AND HEIGHT $f(t)$ PERPENDICULAR TO $t$ -AXIS 	REGION BDD BY $f$ AND $t$ -AXIS IS REVOLVED @ AXIS PERPENDICULAR TO $t$ -AXIS! 
SUBTRACTIVE SHELL	$\int_a^b 2\pi t (f(t) - g(t)) dt$	f, g CONT'S ON $a \leq t \leq b$ , $f(t) \geq g(t)$	SURFACE AREA OF AN OPEN CYLINDER OF RADIUS $t$ AND HEIGHT $f(t) - g(t)$ PERP. TO $t$ -AXIS 	REGION BDD BY $f$ AND $g$ IS REVOLVED @ AXIS PERPENDICULAR TO $t$ -AXIS! 