

AT LEAST ONE

~~TWO~~ OPEN PROBLEMS~~X~~
IN CATEGORICAL
QUANTUM THEORY

ARI ROSENFIELD
DEPT. OF MATHEMATICS

WHAT IS THIS?

ORIGINAL PLAN: TALK ABOUT TWO PROJECTS
I STARTED, BUT DIDN'T FINISH, WHICH I
THOUGHT COULD BE ACCESSIBLE TO PEOPLE
WHO DON'T KNOW TONS OF CATEGORY
THEORY.

ONE PROBLEM IS NO LONGER OPEN;
IT'S A VERY PRETTY FAMILY OF RESULTS,
SO I'LL TELL YOU ABOUT IT ANYWAY.

LED TO A NEW OPEN PROBLEM, BUT
IT MIGHT BE HARD.

WHAT IS THIS?

MAIN GOAL:

IF ANY OF THIS SOUNDS FUN AND/OR
YOU'RE TRYING TO LEARN SOME CATEGORY
THEORY AND MAYBE PUBLISH A MATH
PAPER, LET'S TALK AFTER!

WHY CQT?

- CT :



"STUDY THE TRANSFORMATIONS A CLASS OF OBJECTS CAN UNDERGO, AS LONG AS YOU CAN COMPOSE THEM."

- CQT : "REFORMULATE QUANTUM THEORY IN A GENERAL DAGGER COMPACT CATEGORY." Hilb is ONE OF THESE.

WHY CQT?

- FOR COMPUTING: WANT TO VIEW QUANTUM INFORMATION THEORY AS ONE AMONGST MANY COMPUTATIONAL THEORIES, SEE WHAT DISTINGUISHES IT.
- PRACTICAL APPLICATIONS: ??? ??




1

MODELING CONCURRENCY IN SYMMETRIC MONOIDAL CATEGORIES

EXAMPLE
OF MORPHISM
IN $X \amalg Y$

(j/w
CARMEN CONSTANTIN
CHRIS HEUNEN
NESTA v.d. SCHAAF)



BACKGROUND

TRADITIONALLY, QUANTUM COMPUTATION
HAPPENS IN THE CATEGORY

Hilb {

- OBJECTS: FINITE DIMENSIONAL HILBERT SPACES
(QUDIT REGISTERS)
- MORPHISMS: BOUNDED \mathbb{C} -LINEAR MAPS
(QUANTUM CIRCUITS)

BACKGROUND

Hilb HAPPENS TO BE A
SYMMETRIC MONOIDAL CATEGORY:


MONOIDAL:

CAN PUT TWO
REGISTERS SIDE-BY-SIDE
TO FORM A NEW ONE:

$$\begin{matrix} \mathbb{C}^2 \\ \mathbb{C}^2 \end{matrix} \xrightarrow{\quad} \mathbb{C}^2 \otimes \mathbb{C}^2$$

AND THERE'S A
"TRIVIAL REGISTER" w/r/t \otimes

SAME WITH
CIRCUITS:



SYMMETRIC:

SWITCHING LEFT WITH RIGHT IS
UNPROBLEMATIC. (i.e., \otimes IS COMMUTATIVE.)

BACKGROUND

SYMMETRIC MONOIDAL
CATEGORY

OBJECT

MORPHISM

RESOURCE THEORY


TYPE OF RESOURCE

PROCESS

BACKGROUND

ANY SYMMETRIC MONOIDAL CATEGORY $(\mathcal{X}, \otimes, I)$
COMES WITH:

a STRING-DIAGRAMMATIC CALCULUS
FOR ITS MORPHISMS.



MORPHISM
= STRING DIAGRAM
= CIRCUIT DIAGRAM

BUT NOW WE
IMPOSE RULES
FOR SENSIBLY
TRANSFORMING
ONE TO ANOTHER.

BACKGROUND

ANY SYMMETRIC MONOIDAL CATEGORY $(\mathcal{X}, \otimes, I)$
COMES WITH:

- LATTICE OF OPENS OF A
TOPLOGICAL SPACE
b) AN INTERNAL \vee TOPOLOGICAL SPACE
CALLED $I_{Sub}(\mathcal{X})$.

IT'S A LATTICE OF DISTINGUISHED
MONOMORPHISMS INTO THE
TENSOR UNIT I , CALLED SUBUNITS.

(CATEGORIFICATION OF CENTRAL IDEMPOTENTS
OF A RING.)

BACKGROUND

FACT: EACH OF THESE CORRESPONDS TO A SYMMETRIC MONOIDAL SUBCATEGORY OF \mathcal{X} , AND VICE VERSA. ("SUBUNIT" BECAUSE EACH IS THE TENSOR UNIT FOR A SYMM-MONOIDAL SUBCATEGORY.)

FOR NARRATIVE PURPOSES, THINK OF THEM AS ADMISSIBLE LOCATIONS FOR PROCESSES TO OCCUR AT.

BACKGROUND

ANY SYMMETRIC MONOIDAL CATEGORY $(\mathcal{X}, \otimes, I)$
COMES WITH:

c) NOTIONS OF RESTRICTION OF A
MORPHISM TO A SUBUNIT AND
SUPPORT OF A MORPHISM ON A SUBUNIT.

"THERE'S A PLACE IN \mathcal{X} WHERE A
GIVEN PROCESS OCCURS, AND IT
ONLY OCCURS THERE."



Contents lists available at ScienceDirect

Journal of Pure and Applied Algebra

www.elsevier.com/locate/jpaa



Tensor topology

Pau Enrique Moliner, Chris Heunen ^{*,1}, Sean Tull

University of Edinburgh, United Kingdom of Great Britain and Northern Ireland



BACKGROUND

SAY THAT TWO PROCESSES CAN
EXCHANGE RESOURCES IF THEY
HAVE COMMON SUPPORT.

THE PROBLEM


WE'D EXPECT SYMMETRIC MONOIDAL
CAT's TO BE GOOD FOR MODELING
COMPUTATIONAL CONCURRENCY:

- SEQUENTIAL PROCESSES ✓
(COMPOSING MORPHISMS)
- PARALLEL PROCESSES ✓
(TENSORING MORPHISMS)

THE PROBLEM

WE'D LIKE PROCESSES IN A GIVEN \mathcal{X}
TO BE ABLE TO EXCHANGE RESOURCES.

e.g., TELEPORTATION:




(PRESENCE OF
DUAL OBJECTS)

NICE PROPERTIES OF
 $Hilb_{fd}$ LET US
ASSUME ALL PROCESSES
INVOLVED HAVE COMMON
SUPPORT.

"LIVE IN THE SAME
UNIVERSE."

THE PROBLEM



THOSE NICE PROPERTIES DON'T HOLD FOR A GENERAL $(\mathcal{X}, \otimes, I)$.

MIGHT NOT BE ABLE TO FORM JOINS IN $\text{ISub}(\mathcal{X})$.

CAN REMEDY THIS BY FINDING A NEW CATEGORY ?? WITH A NICER SUBUNIT LATTICE.

THE PROBLEM

WHY WOULD WE WANT TO FIX THIS PROBLEM?

BECAUSE WE WANT TO DO
SHEAF THEORY.

Advances in Mathematics 416 (2023) 108900



Contents lists available at ScienceDirect

Advances in Mathematics

journal homepage: www.elsevier.com/locate/aim



BUT THAT WOULD BE
A MUCH LONGER TALK.

SEE
FOR MORE ON
THIS

Sheaf representation of monoidal categories [☆]



Rui Soares Barbosa ^a, Chris Heunen ^{b,*}

^a INL – International Iberian Nanotechnology Laboratory, Braga, Portugal

^b University of Edinburgh, United Kingdom

SO FAR

CONSTRUCTED A CANDIDATE FOR ?? :

GIVEN $x, y \in \text{SymMonCat}$, FORM

$x \amalg y \in \text{SymMonCat}$ WHOSE


MORPHISMS ARE 2-COLORED STRING

DIAGRAMS.

(HARD PART: ALGEBRAIC FORMALISM. MORPHISMS
ARE ACTUALLY MATRICES WITH ENTRIES IN
A JUDICIOUSLY CHOSEN SEMIRING.)

TO - DO

- VERIFY THAT $X \amalg Y$ IS A PUSHOUT
IN SymMonCat. (REQUIRES SOME LIGHT
HIGHER CT.)
- FIND A NORMAL FORM FOR $\text{Mor}(X \amalg Y)$.
- GENERALIZE TO $n > 2$. (i.e., WRITE
DOWN WHAT AN n -COLORED STRING-
DIAGRAM CALCULUS SHOULD BE.)



(2)

FUNCTIONAL ENTROPY

(j/w MAICOL OCHOA)

I LIED. THIS IS MERELY
CQT-ADJACENT. NEEDED
CATCHY TITLE

BACKGROUND

OVERARCHING PROJECT: FIND SETS OF AXIOMS WHICH CHARACTERIZE ENTROPY (RELATIVE ENTROPY, ENTROPY DIFFERENCE) AS A FUNCTION FROM SOME CLASS OF (GENERALIZED) PROBABILITY SPACES INTO $[0, \infty)$ OR $[0, \infty]$.

FROM 1950S:

FADDEEV,



KHINCHIN.

BACKGROUND

REMINDER: SAY $(X, \mu) \xrightarrow{f} (Y, \nu)$
IS A MEASURE-PRESERVING MAP
OF FINITE PROBABILITY SPACES.

THE SHANNON ENTROPY DIFFERENCE
OF f IS

$$H(\mu) - H(\nu) = \sum_y \nu_i \ln \nu_i - \sum_x \mu_j \ln \mu_j$$

(ALSO RECALL: ALWAYS NON NEGATIVE.)

BACKGROUND

CT PERSPECTIVE: WE LIKE WHEN
"MAPS" BETWEEN CATEGORIES ARE
FUNCTORIAL, RESPECTING COMPOSITIONAL
STRUCTURE.

ANALOGY:

- FOR GROUP THEORY, WE LIKE OUR MAPS TO BE HOMOMORPHISMS
- FOR MEASURE THEORY, WE LIKE OUR MAPS TO BE MEASURE-PRESERVING
... ETC.

BACKGROUND

FROM THAT PERSPECTIVE, NATURAL
TO ASK WHAT HAPPENS IF WE
REQUIRE ENTROPY TO BEHAVE
FUNCTORIALLY.

TURNS OUT, WE NECESSARILY GET
SHANNON ENTROPY DIFFERENCE:

BACKGROUND

(BUILDING ON FADDEEV'S CHARACTERIZATION)

THEOREM. (BAEZ - FRITZ - LEINSTER, 2011)

IF $F: \text{FinProb} \rightarrow [0, \infty)$ IS ANY
CONTINUOUS FUNCTION WHICH IS

- CONVEX LINEAR

$$F(\lambda f \oplus (1-\lambda)g) = \lambda Ff + (1-\lambda)Fg$$

- FUNCTORIAL

$$F(fg) = Ff + Fg$$

THEN F IS A REAL MULTIPLE OF
THE SHANNON ENTROPY DIFFERENCE.



BACKGROUND

- The charm of this result is that the first two hypotheses look like linear conditions, and none of the hypotheses hint at any special role for the function $-p \ln p$, but it emerges in the conclusion.

THEN, ANALOGUES FOR :

- FINITE MEASURE SPACES
- TSALLIS ENTROPY (MATH. ECOLOGY?)

PARZYNAT, 2022 :

- VON NEUMANN ENTROPY OF FINITE-DIMENSIONAL NONCOMMUTATIVE PROBABILITY SPACES

"PROBLEM"

CAN WE EXTEND ANY OF THESE
RESULTS TO THE COUNTABLY INFINITE
/INFINITE-DIMENSIONAL CASE? YES:

"SOLUTION"

THEOREM. (GAGNÉ-PANANGADEN, 2023)

IF A FUNCTOR $F: \text{StdBorel} \longrightarrow [0, \infty]$
IS LOWER-SEMICONINUOUS, CONVEX LINEAR,
+ 1 OTHER THING, THEN ON MORPHISMS,
 F IS A REAL MULTIPLE OF
KULLBACK-LEIBLER DIVERGENCE.

GENERALIZES B-F-L; ANSWERS OUR
QUESTION SINCE A COUNTABLE SET /W
DISCRETE σ -ALGEBRA GIVES A StdBorel.

OBVIOUS NEW QUESTION: WHAT'S THE
ANALOGUE FOR VON NEUMANN ENTROPY?


AT THE MOMENT, I DON'T HAVE A
CONJECTURE (DON'T UNDERSTAND THE
PAPER WELL ENOUGH YET!)

LIKELY OPEN, MAY BE HARD,
CERTAINLY WORTHWHILE.



THANKS FOR LISTENING!

"THE FUNDAMENTALS OF
CATEGORICAL QUANTUM MECHANICS



SLIDES:

ari-rosenfield.github.io