

Logic

Aria Ruler(Forouzandeh)

March 11, 2026

This page intentionally left blank.

Contents

1	Atomes	1
1.1	Basic tools	1
1.2	constants and variables	3
2	propositional logic (zeroth-order logic)	5
2.1	What is proposition?	5
2.2	Rules of propositions.	6
3	first-order logic	13
3.1	what is predicate?	13
3.2	Rules of predicates.	14
4	The beginning of set theory	19
4.1	Axioms of set theory	19
4.2	definitions of set theory	21
5	Relations and functions	29
5.1	cartesian product	29
5.2	Relations	30
5.3	Functions	32
5.4	Family of sets	34

This page intentionally left blank.

Chapter 1

Atomes

1.1 Basic tools

we start the book by building logic step by step.block by block in first it might seem a little annoying but I advice you to be patient.book will show its power soon. like euclidian geometry this book has undefined concepts. I tried to explain them for you to understand.but they don't have any definition.

Undefined 1.1.1. *Urelements* are the most fundamental entities in a logical. They cannot be constructed, defined, or described in terms of other things. Their existence is primitive—accepted without internal structure or relational properties. Urelements simply are.

In this book, we will denote individual urelements using lowercase letters $u, v, w, \dots$

Undefined 1.1.2. *Type* parallel lines used to show that an object exists (you will undrestnd what is object soon.) (we will show you how to use them).

1.	u			
2.		v		
3.			w	

Rule 1.1.1. *fusion* it's the Rule that explain how sets made. (this rule is schema.it means it works everywhere for every object.)

$m.$	u_0	
	$\vdots$	
$m + n.$	u_n	
$o.$		$\{u_0, u_1 \dots u_n\}$

Undefined 1.1.3. Set we can't say what exactly sets are but we know sets can be made by fusion (if "a" is a set we write "set(a)" from now we denote sets with uppercase letters like A, B, C, ...).

definition 1.1.1. Object urelements and sets.(we write obj instead of object). (from now we denote objects with lowercase letters like a, b, c, ...).

Rule 1.1.2. fission it show us how to break an obj. (this rule is scheme it means it works everywhere for every obj.)

$o.$		$\{a_0, a_1 \dots a_n\}$
$m.$	a_0	
	$\vdots$	
$m + n.$	a_n	

Example 1.1.1. $\{a, b, c\} / \therefore \{a, b\}$ (prove if $\{a, b, c\}$ exists then $\{a, c\}$ exists)

1.		$\{a, b, c\}$
2.	a	$fi1$
3.	b	$fi1$
4.	c	$fi1$
5.		$\{a, c\} \quad fu2,3,4$

Example 1.1.2. $\{a, b\} / \therefore \{b, a\}$

1.		$\{a, b\}$
2.	a	$fi1$
3.	b	$fi1$
4.		$\{b, a\} \quad fu2,3$

1.2 constants and variables

obj are two types. we know them and we know what exactly are we talking about and unknowns are mystery. all the obj we have used until now were known ones we call them **"constants"**.

Undefined 1.2.1. *variables* A variable is a symbol (usually a letter like x, y , or z) that represents an unknown obj.

definition 1.2.1. *predicate* " $set()$ " and identity " $=$ " and membership " $\in$ " symbols are predicates. " $set()$ " is unary and identity and membership are binary.

This page intentionally left blank.

Chapter 2

propositional logic (zeroth-order logic)

2.1 What is proposition?

before answering to this question let me show you what is well formed formula(WWF).

definition 2.1.1. Atomic formula every single obj or or two obj(conatants and variable) and a binary predicate ($=, \in$) between them like $x = b, y \in C, \dots$
we denote atomic formulas with A_i .

Undefined 2.1.1. logical operators ($\neg, \rightarrow, \wedge, \vee$) tool that used between atoms to make more complex formulas (all of the opratores are binary except " $\neg$ " that is unary).

definition 2.1.2. well formed formula(WWF) these are formulas.

1. atomic formulas (A_i)
2. if A is a formula then $\neg A$ is a formula too.
3. if A, B are two formulas then $A \bigcirc B$ is a formula too. ($\bigcirc$ can be $\rightarrow, \wedge, \vee$).

for example ($A_0 \rightarrow A_1, \neg A_3$)

we easily call WWF "formula" and also we denote WWF with $A, B, C, \dots$

you can easily see that formula are made up by atomic formulas.in general we say formula is an atomic formula or combination of them.

definition 2.1.3. open formula formulas that variables used in one of its atomic formulas.

the formulas that are not open call "**closed formulas**" or "**sentence**". Now question is "are very closed formulas proposition?" (propositions in classic logic are sentences that are true or false?) the answer is unfortunately **No**.

definition 2.1.4. Atomic proposition every atomic formula that can be proven by tools introduced in chapter 1($I =, E =, I \in, E \in, I \perp, \perp E$).
we denote atomic formulas with p_i .

definition 2.1.5. *proposition* every WWF formula that its atomes are all atomic propositions. we denote propositions with $p, q, r, \dots$

definition 2.1.6. *open proposition* assume formula of a proposition but Substitute some obj and replace them with variables. we call it open proposition and denote it with capital letter (P, Q, R) shows other types of **predicates** and variables like $(P(x), Q(x, y), \dots)$

definition 2.1.7. *Substitution for obj* we define Substitution an obj with variable like this.

1. for variables if $x = y$ then $y[a/x] := a$ and if $x \neq y$ then $y[a/x] := y$.
2. for constants $c[a/x] := c$.

definition 2.1.8. *Substitution for formulas* we define Substitution for a formula like this.

1. for atomic formulas we define it like this. if it's a single object we do it as we explain in "**Substitution for obj**". if it's like $x = y$ or $x \in y$ (x, y can be both constants and variables). we define it like this.

$$(x = y)[a/z] := (x[a/z] = y[a/z])$$

$$(x \in y)[a/z] := (x[a/z] \in y[a/z])$$

2. if formula A is complex we define it like this.

$$(B \circ C)[a/x] := (B[a/x] \circ C[a/x])$$

($\circ$ can be $\rightarrow, \wedge, \vee$).


$$(\neg B)[a/x] := (\neg B[a/x])$$

simply means in a formula put an obj like b instead of another like a . we denote it by $A[b/a]$ or $A[\frac{b}{a}]$

2.2 Rules of propositions.

definition 2.2.1. *proposition schema(variable proposition)* a proposition that is unknown for us and it works as variable. (we show them with $\varphi, \psi, \theta, \dots$)

Undefined 2.2.1. *Assumption for conditional proof(acp)* is a tool for assuming a proposition. (any proposition can't get out but it can get in. right line must be shorter than left one. (it can't fall out). and every hypothesis that opens must be closed.)



Rule 2.2.1. rules of logical operations ($\rightarrow I, \rightarrow E, \wedge I, \wedge E, \vee I, \vee E, \neg I, \neg E, \perp I, \perp E$) *this rule is scheme it means it works everywhere for every propositions. (thats why we use φ, ψ instead of p, q)*

$\begin{array}{ l} \varphi \\ \vdots \\ \psi \end{array}$ $\therefore \varphi \rightarrow \psi \quad \rightarrow I$	$\begin{array}{l} \varphi \rightarrow \psi \\ \hline \varphi \end{array}$ $\therefore \psi \quad \rightarrow E$
$\begin{array}{l} \varphi \\ \psi \\ \hline \end{array}$ $\therefore \varphi \wedge \psi \quad \wedge I$	$\begin{array}{l} \varphi \wedge \psi \\ \hline \end{array}$ $\therefore \varphi \quad \wedge E$ $\begin{array}{l} \varphi \wedge \psi \\ \hline \end{array}$ $\therefore \psi \quad \wedge E$
$\begin{array}{l} \varphi \\ \hline \end{array}$ $\therefore \varphi \vee \psi \quad \vee I$ $\begin{array}{l} \varphi \\ \hline \end{array}$ $\therefore \psi \vee \varphi \quad \vee I$	$\varphi \vee \psi$ $\begin{array}{ l} \varphi \\ \vdots \\ \theta \end{array}$ $\begin{array}{ l} \psi \\ \vdots \\ \theta \end{array}$ $\therefore \theta \quad \vee E$
$\begin{array}{ l} \varphi \\ \vdots \\ \perp \end{array}$ $\therefore \neg \varphi \quad \neg I$	$\begin{array}{ l} \neg \varphi \\ \vdots \\ \perp \end{array}$ $\therefore \varphi \quad \neg E$
$\begin{array}{l} \varphi \\ \neg \varphi \\ \hline \end{array}$ $\therefore \perp \quad \perp I$	$\begin{array}{l} \perp \\ \hline \end{array}$ $\therefore \varphi \quad \perp E$

definition 2.2.2. $\varphi \leftrightarrow \psi : (\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$

definition 2.2.3. schema of prove you've got knowed by now what is schema of a rule. if you want to prove a theorem that it's schema form. you can't prove it directly. here we use a technique calls **"schema of prove"**. it let us prove for all situations at the same time.

Theorem 2.2.1. (schema) ($\leftrightarrow I, \leftrightarrow E$)

$ \begin{array}{c c} \varphi \\ \vdots \\ \psi \\ \hline \psi \\ \vdots \\ \varphi \\ \hline \therefore \varphi \leftrightarrow \psi \quad \leftrightarrow I \end{array} $	$ \begin{array}{ccc} \varphi \leftrightarrow \psi & & \varphi \leftrightarrow \psi \\ \frac{\varphi}{\therefore \psi} & \leftrightarrow E & \frac{\psi}{\therefore \varphi} \quad \leftrightarrow E \end{array} $
--	--

1	φ	<i>acp</i>
	$\vdots$	
n	ψ	
$n+1$	ψ	<i>acp</i>
	$\vdots$	
m	φ	
$m+1$	$\varphi \rightarrow \psi$	$\rightarrow, I1, n$
$m+2$	$\psi \rightarrow \varphi$	$\rightarrow, In+1, m$
$m+3$	$(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$	$\wedge I, m+1, m+2$
$m+4$	$\varphi \leftrightarrow \psi$	<i>def.</i> , $m+3$

1	$\varphi \leftrightarrow \psi$	<i>ass</i>
2	φ	<i>ass</i>
3	$(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$	<i>def.1</i>
4	$\varphi \rightarrow \psi$	$\wedge E3$
5	ψ	$\rightarrow E1, 4$

Example 2.2.1. for conatants p, q, r we can prove these. Proof Is Left As An Exercise To The Reader. 😊

1. $p \rightarrow q, q \rightarrow r / \therefore p \rightarrow r$
2. $q / \therefore p \rightarrow q$
3. $(p \wedge q) \wedge r / \therefore p \wedge (q \wedge r)$
4. $p \wedge q / \therefore q \wedge p$
5. $p \rightarrow q / \therefore (p \wedge r) \rightarrow q$
6. $(p \wedge q) \rightarrow r / \therefore p \rightarrow (q \rightarrow r)$

$$7. p \rightarrow (q \rightarrow r) / \therefore (p \wedge r) \rightarrow q$$

$$8. p \vee q / \therefore q \vee p$$

$$9. (p \vee q) \vee r / \therefore p \vee (q \vee r)$$

$$10. p \wedge (q \vee r) / \therefore (p \wedge q) \vee (p \wedge r)$$

$$11. (p \rightarrow q) \wedge (r \rightarrow s) / \therefore (p \vee r) \rightarrow (q \vee s)$$

$$12. p \rightarrow q, \neg q / \therefore \neg p$$

$$13. p \rightarrow q / \therefore \neg p \vee q$$

proofs.

$ \begin{array}{lll} 1 & p \rightarrow q & ass \\ 2 & q \rightarrow r & ass \\ 3 & \left \begin{array}{l} p \\ q \\ r \end{array} \right. & \begin{array}{l} acp \\ \rightarrow E1, 3 \\ \rightarrow E2, 4 \end{array} \\ 4 & & \\ 5 & & \\ 6 & p \rightarrow r & \rightarrow I \end{array} $	$ \begin{array}{lll} 1 & q & ass \\ 2 & \left \begin{array}{l} p \\ q \end{array} \right. & \begin{array}{l} acp \\ ass \end{array} \\ 3 & & \\ 4 & p \rightarrow q & \rightarrow I \end{array} $	$ \begin{array}{lll} 1 & (p \wedge q) \wedge r & ass \\ 2 & p \wedge q & \wedge E, 1 \\ 3 & r & \wedge E, 1 \\ 4 & p & \wedge E, 2 \\ 5 & q & \wedge E, 2 \\ 6 & q \wedge r & \wedge I, 3, 5 \\ 7 & p \wedge (q \wedge r) & \wedge I, 4, 6 \end{array} $
$ \begin{array}{lll} 1 & p \wedge q & ass \\ 2 & p & \wedge E, 1 \\ 3 & q & \wedge E, 1 \\ 4 & q \wedge p & \wedge I, 2, 3 \end{array} $	$ \begin{array}{lll} 1 & p \rightarrow q & ass \\ 2 & \left \begin{array}{l} p \wedge r \\ p \\ q \end{array} \right. & \begin{array}{l} acp \\ \wedge E, 2 \\ \rightarrow E, 1, 3 \end{array} \\ 3 & & \\ 4 & & \\ 5 & (p \wedge r) \rightarrow q & \rightarrow I \end{array} $	$ \begin{array}{lll} 1 & (p \wedge q) \rightarrow r & ass \\ 2 & \left \begin{array}{l} p \\ \left \begin{array}{l} q \\ p \wedge q \\ r \end{array} \right. \end{array} \right. & \begin{array}{l} acp \\ acp \\ \wedge I2, 3 \\ \rightarrow E1, 4 \end{array} \\ 3 & & \\ 4 & & \\ 5 & & \\ 6 & \left \begin{array}{l} q \rightarrow r \end{array} \right. & \rightarrow I \\ 7 & p \rightarrow (q \rightarrow r) & \rightarrow I \end{array} $

1	$p \rightarrow (q \rightarrow r)$	<i>ass</i>	1	$p \vee q$	<i>ass</i>	1	$(p \vee q) \vee r$	<i>ass</i>
2	$p \wedge q$	<i>acp</i>	2	p	<i>acp</i>	2	$p \vee q$	<i>acp</i>
3	p	$\wedge E, 2$	3	$p \vee q$	$\vee I, 2$	3	p	<i>acp</i>
4	$q \rightarrow r$	$\rightarrow E, 1, 3$	4	q	<i>acp</i>	4	$p \vee (q \vee r)$	$\vee I, 3$
5	q	$\wedge E, 2$	5	$q \vee p$	$\vee I, 4$	5	q	<i>acp</i>
6	r	$\rightarrow E, 4, 5$	6	$q \vee p$	$\vee E, 1$	6	$q \vee r$	$\vee I, 5$
7	$(p \wedge q) \rightarrow r$	$\rightarrow I$				7	$p \vee (q \vee r)$	$\vee I, 6$
						8	$p \vee (q \vee r)$	$\vee E, 2$
						9	r	<i>acp</i>
						10	$q \vee r$	$\vee I, 9$
						11	$p \vee (q \vee r)$	$\vee I, 10$
						12	$p \vee (q \vee r)$	$\vee E, 1$

1	$p \wedge (q \vee r)$	<i>ass</i>	1	$(p \rightarrow q) \wedge (r \rightarrow s)$	<i>ass</i>
2	p	$\wedge E1$	2	$(p \rightarrow q)$	$\wedge E1$
3	$q \vee r$	$\wedge E1$	3	$(r \rightarrow s)$	$\wedge E1$
4	q	<i>acp</i>	4	$p \vee r$	<i>acp</i>
5	$p \wedge q$	$\wedge E2, 4$	5	p	<i>acp</i>
6	$(p \wedge q) \vee (p \wedge r)$	$\vee I5$	6	q	$\rightarrow E2, 5$
7	r	<i>acp</i>	7	$q \vee s$	$\vee I, 6$
8	$p \wedge r$	$I \wedge 2, 7$	8	r	<i>acp</i>
9	$(p \wedge q) \vee (p \wedge r)$	$\vee I8$	9	s	$\rightarrow E3, 8$
10	$(p \wedge q) \vee (p \wedge r)$	$\vee I8$	10	$q \vee s$	$\vee I9$
11	$(p \wedge q) \vee (p \wedge r)$	$\vee E3$	11	$q \vee s$	$\vee E, 4$
			12	$(p \vee r) \rightarrow (q \vee s)$	$\rightarrow I$

1	$p \rightarrow q$	<i>ass</i>
2	$\neg q$	<i>ass</i>
3	$\left \begin{array}{l} p \\ q \\ \perp \end{array} \right.$	<i>acp</i>
4		$\rightarrow E, 1, 3$
5		$\perp I, 2, 4$
6	$\neg p$	$\neg I$

1	$p \rightarrow q$	<i>ass</i>
2	$\left \neg(\neg p \vee q) \right.$	<i>acp</i>
3	$\left \left p \right. \right.$	<i>acp</i>
4	$\left \left q \right. \right.$	$\rightarrow E, 1, 3$
5	$\left \left \neg p \vee q \right. \right.$	$\vee I, 4$
6	$\left \left \perp \right. \right.$	$\perp I, 2, 5$
7	$\left \neg p \right.$	$\neg I$
8	$\left \neg p \vee q \right.$	$\vee I, 7$
9	$\left \perp \right.$	$\perp I, 2, 8$
10	$\neg p \vee q$	$\neg I$

Exercise 2.2.1. for conatants p, q, r prove.

1. $p / \therefore p$
2. $/ \therefore p \vee \neg p$
3. $\neg p, p \vee q / \therefore q$
4. $\neg p / \therefore p \rightarrow q$
5. $p \rightarrow q \therefore / \therefore \neg q \rightarrow \neg p$
6. $(p \vee q) \wedge (p \vee r) \therefore / \therefore (p \vee (q \wedge r))$
7. $\neg p \wedge \neg q \therefore / \therefore \neg(p \vee q)$
8. $\neg p \vee \neg q \therefore / \therefore \neg(p \wedge q)$
9. $(\neg p \rightarrow q) / \therefore ((p \rightarrow q) \rightarrow q)$
10. $/ \therefore (p \wedge q \rightarrow r) \wedge (p \wedge s \rightarrow r) \rightarrow (p \wedge (q \vee s) \rightarrow r)$
11. $/ \therefore (p \wedge q \rightarrow r) \vee (p \wedge s \rightarrow r) \rightarrow (p \wedge (q \wedge s) \rightarrow r)$
12. $/ \therefore (p \rightarrow (q \rightarrow r)) \rightarrow ((p \rightarrow q) \rightarrow (p \rightarrow r))$
13. $/ \therefore (p \rightarrow q) \rightarrow ((q \rightarrow r) \rightarrow (p \rightarrow r))$

Attention 2.2.1. we can rewrite **Example** and **Exercise** propositions and proofs with proposition schemas and schema of proves. so they are all rules.

This page intentionally left blank.

Chapter 3

first-order logic

3.1 what is predicate?

definition 3.1.1. *Big AND and OR* assume a set like “ $\{a_0, a_1, a_2, \dots, a_n\}$ ” we define **big AND** like this.

$$\bigwedge_{i=0}^n P(a_i) : P(a_0) \wedge P(a_1) \wedge P(a_2) \wedge \dots \wedge P(a_n)$$

and **big OR**

$$\bigvee_{i=0}^n P(a_i) : P(a_0) \vee P(a_1) \vee P(a_2) \vee \dots \vee P(a_n)$$

definition 3.1.2. *Universe of discourse* we specify a set like “ $\{a_0, a_1, a_2, \dots, a_n\}$ ” as universe of a Logic world.

Rule 3.1.1. *use of open propositions* assume universe of discourse is “ $\{a_0, a_1, a_2, \dots, a_n\}$ ”

$$c \quad P(x)$$

means :

$$c_0 \quad P[a_0/x]$$

$$c_1 \quad P[a_1/x]$$

$$c_2 \quad P[a_2/x]$$

$$\vdots$$

$$c_n \quad P[a_n/x]$$

if Px was been assemed then like this

$$c \quad \left| \begin{array}{l} P(x) \\ \vdots \end{array} \right.$$

means :

$$\begin{array}{c} c_0 \\ c_1 \\ c_2 \\ c_n \end{array} \quad \left| \begin{array}{l} P[a_0/x] \\ \vdots \\ P[a_1/x] \\ \vdots \\ P[a_2/x] \\ \vdots \\ P[a_n/x] \\ \vdots \end{array} \right.$$

definition 3.1.3. predicate schema(variable predicate) a predicate that is unknown for us and it works as variable.(we show them with $\phi, \psi, \theta, \dots$)

definition 3.1.4. Universal quantification assume universe of discourse is “ $\{a_0, a_1, a_2, \dots, a_n\}$ ” we define $\forall x P(x)$

$$\forall x \phi(x) : \bigwedge_{i=0}^n \phi[a_i/x]$$

definition 3.1.5. Existential quantification assume universe of discourse is “ $\{a_0, a_1, a_2, \dots, a_n\}$ ” we define $\exists x P(x)$

$$\exists x \phi(x) : \bigvee_{i=0}^n \phi[a_i/x]$$

as you can see $\forall x \phi(x)$ and $\exists x \phi(x)$ are sentences.

definition 3.1.6. Free variables variables that are not bound by a quantifier. (you can easily see that every formula with no free variables are sentences) (previously we sayed that every sencece is not necessary a proposition)

3.2 Rules of predicates.

Theorem 3.2.1. (schema) ($\forall I, \forall E, \exists I, \exists E$) assume universe of discourse is “ $\{a_0, a_1, a_2, \dots, a_n\}$ ”

<i>Rules</i>	<i>Terms and Conditions</i>
$\frac{\phi(x)}{\therefore \forall x\phi(x)} \quad \forall I$	1. x is a variable. 2. $\phi(x)$ must not be assumed.
$\frac{\forall x\phi(x)}{\therefore \phi[a/x]} \quad \forall E$	1. a is an obj in universe of discourse.(it can be constant or variable.)
$\frac{\phi[a/x]}{\therefore \exists x\phi(x)} \quad \exists I$	1. a is an obj in universe of discourse.(it can be conatant or variable.)
$\frac{\begin{array}{c} \exists x\phi(x) \\ \left \begin{array}{c} \phi(x) \\ \vdots \\ \theta \end{array} \right. \\ \therefore \theta \end{array}}{\quad \quad \quad \exists E}$	1. x is a variable. 2. $\phi(x)$ must not be assumed in previous lines.(except ones that are closed) 3. x is not free in θ

1	$\phi(x)$	<i>ass</i>
2	$\phi[a_0/x]$	<i>def.1</i>
	$\vdots$	
n	$\phi[a_n/x]$	<i>def.1</i>
$n+1$	$\bigwedge_{i=0}^n \phi[a_i/x]$	$\wedge I, 2, 3, \dots, n+2$
$n+2$	$\forall x \phi(x)$	<i>def.n+3</i>

1	$\phi[a/x]$	
2	$\bigvee_{i=0}^n \phi[a_i/x]$	$\vee I1$
3	$\exists x \phi(x)$	<i>def2</i>

1	$\forall x \phi(x)$	<i>ass</i>
2	$\bigwedge_{i=0}^n \phi[a_i/x]$	<i>def.1</i>
3	$\phi[a/x]$	$\wedge E2$

1	$\exists x \phi(x)$	
2	$\bigvee_{i=0}^n \phi[a_i/x]$	
3	$\phi[a_0/x]$	<i>acp</i>
	$\vdots$	
n	θ	
$n+1$	$\phi[a_1/x]$	<i>acp</i>
	$\vdots$	
m	θ	
	$\vdots$	
l	$\phi[a_2/x]$	<i>acp</i>
	$\vdots$	
r	θ	
$r+1$	θ	$\exists E, 3$

Example 3.2.1. for conatants P, Q, R we can prove these. Proof Is Left As An Exercise To The Reader. 😊

1. $\forall x(P(x) \wedge Q(x)) \therefore / \therefore \forall x P(x) \wedge \forall x Q(x)$
2. $\exists x(P(x) \vee Q(x)) \therefore / \therefore \exists x P(x) \vee \exists x Q(x)$
3. $\forall x P(x) \vee \forall x Q(x) / \therefore \forall x(P(x) \vee Q(x))$
4. $\exists x(P(x) \wedge Q(x)) / \therefore \exists x P(x) \wedge \exists x Q(x)$
5. $\forall x(P(x) \rightarrow Q(x)) / \therefore \forall x P(x) \rightarrow \forall x Q(x)$
6. $\exists x P(x) \rightarrow \exists x Q(x) / \therefore \exists x(P(x) \rightarrow Q(x))$
7. $\forall x P(x) \wedge Q \therefore / \therefore \forall x(P(x) \wedge Q)$
8. $\forall x P(x) \vee Q \therefore / \therefore \forall x(P(x) \vee Q)$
9. $\exists x P(x) \wedge Q \therefore / \therefore \exists x(P(x) \wedge Q)$

10. $\exists xP(x) \vee Q \therefore / \therefore \exists x(P(x) \vee Q)$
11. $Q \rightarrow \forall xP(x) \therefore / \therefore \forall x(Q \rightarrow P(x))$
12. $Q \rightarrow \exists xP(x) \therefore / \therefore \exists x(Q \rightarrow P(x))$
13. $\forall xP(x) \rightarrow Q \therefore / \therefore \exists x(P(x) \rightarrow Q)$
14. $\exists xP(x) \rightarrow Q \therefore / \therefore \forall x(P(x) \rightarrow Q)$
15. $\exists x\neg P(x) \therefore / \therefore \neg\forall xP(x)$
16. $\forall x\neg P(x) \therefore / \therefore \neg\exists xP(x)$

Exercise 3.2.1. for conatants P, Q, R prove.

1. $\exists x(P(x) \wedge \exists yQ(y) \rightarrow \forall zR(z)) / \therefore \forall y\forall z(\forall xP(x) \wedge Q(y) \rightarrow R(z))$
2. $\forall x\exists y(P(x) \wedge Q(y) \rightarrow \forall zR(z)) / \therefore \forall z(\exists xP(x) \wedge \forall yQ(y) \rightarrow R(z))$
3. $\forall xP(x) \rightarrow \exists yQ(y) / \therefore \exists x\exists y(P(x) \rightarrow Q(y))$
4. $\forall x\forall z\exists y\exists w(P(y, z) \vee Q(z) \rightarrow R(x, w)) / \therefore (\exists z\forall yP(y, z) \vee \exists zQ(z)) \rightarrow \forall x\exists wR(x, w)$
5. $\exists xP(x) \vee \exists x(Q(x) \wedge R(x)) / \therefore \exists x(P(x) \vee Q(x)) \wedge \exists x(P(x) \vee R(x))$
6. $\exists x\exists z\forall y(P(x, y) \rightarrow Q(z) \vee R(x, y)) / \therefore \forall x\exists yP(x, y) \rightarrow (\exists zQ(z) \vee \exists x\exists zR(x, y))$
7. $\forall x\exists y(\exists zP(x, y, z) \wedge Q(x, y)) \vee \forall x\exists y\exists z(P(x, y, z) \wedge R(x, y)) / \therefore \forall x\exists y\exists z(P(x, y, z) \wedge (Q(x, y) \vee R(x, y)))$

Attention 3.2.1. we can rewrite **Example** and **Exercise** propositions and proofs with proposition schemas and schema of proves.so they are all rules.

This page intentionally left blank.

Chapter 4

The beginning of set theory

4.1 Axioms of set theory

we emphasize before that set has no definition. but we can describe what set is with rules and axioms.

Axiom 4.1.1. Existence of universe of set theory *There exists a set V (we denote it as V . it simply means the set of all sets), known as the Von Neumann Universe, which contains every set. (this axiom can't be formalized in first order logic). (there must be another axiom let us use logic for set theory but we ignore that)*

definition 4.1.1. Subset *a subset is a set where every element is also an element of another, larger set called the superset*

$$A \subseteq B : \forall x(x \in A \rightarrow x \in B)$$

Theorem 4.1.1. (schema) $(\subseteq I, \subseteq E)$

$\begin{array}{c} x \in A \\ \vdots \\ x \in B \end{array}$ $\therefore A \subseteq B \quad \subseteq I$	$\begin{array}{c} A \subseteq B \\ a \in A \\ \hline \therefore a \in B \quad \subseteq E \end{array}$
--	--

1	$x \in A$	<i>acp</i>	1	$A \subseteq B$	<i>ass</i>
	$\vdots$		2	$a \in A$	<i>ass</i>
n	$x \in B$		3	$\forall x(x \in A \rightarrow x \in A)$	<i>def.1</i>
$n+1$	$x \in A \rightarrow x \in B$	$I \rightarrow, 1, n$	4	$a \in A \rightarrow a \in B$	$\forall E, 4$
$n+2$	$\forall x(x \in A \rightarrow x \in B)$	$\forall I, n+1$	5	$a \in B$	$E \rightarrow 3, 5$
$n+3$	$A \subseteq B$	<i>def. n+2</i>			

1	$A \subseteq B$	<i>ass</i>
2	$B \subseteq C$	<i>ass</i>
3	$\left \begin{array}{l} x \in A \\ x \in B \\ x \in C \end{array} \right.$	<i>acp</i>
4		$\subseteq E, 1, 3$
5		$\subseteq E, 2, 4$
6	$A \subseteq C$	$\subseteq I, 3, 5$

$$\forall X, Y (\forall x (x \in X \leftrightarrow x \in Y) \rightarrow X = Y)$$
$$\forall X, Y ((X \subseteq Y \wedge Y \subseteq X) \rightarrow X = Y)$$

$ \begin{array}{ l} x \in A \\ \vdots \\ x \in B \\ \hline x \in B \\ \vdots \\ x \in A \end{array} $	$\frac{A = B \quad a \in A}{\therefore a \in B}$ $E =$ $\frac{A = B \quad a \in B}{\therefore a \in A}$ $E =$
---	---

1	$x \in A$	acp
	$\vdots$	
n	$x \in B$	
$n + 1$	$A \subseteq B$	$\subseteq I, 1, n$
$n + 2$	$x \in B$	acp
	$\vdots$	
m	$x \in A$	
$m + 1$	$B \subseteq A$	$\subseteq I, n + 2, m$
$m + 2$	$\forall X, Y ((X \subseteq Y \wedge Y \subseteq X) \rightarrow X = Y)$	<i>Extensionality</i>
$m + 3$	$(A \subseteq B \wedge B \subseteq A) \rightarrow A = B$	$\forall E, m + 2$
$m + 4$	$A \subseteq B \wedge B \subseteq A$	$\wedge I, n + 1, m + 1$
$m + 5$	$A = B$	$E \rightarrow, m + 3, m + 4$

Axiom 4.1.3. Weak comprehension (schema) this axioms says for every open proposition with just one variable like $\phi(x, x_0, x_1, \dots, x_n)$ there is a set such that every x is a member of that set if and only if it holds $\phi(x, x_0, x_1, \dots, x_n)$.

$$\forall x_0, x_1, \dots, x_n \exists X \forall x (\phi(x, x_0, x_1, \dots, x_n) \leftrightarrow x \in X)$$

by axiom of extentionality we can show that X is unique. so we denote it as “ $\{x | \phi(x, x_0, x_1, \dots, x_n)\}$ ”.

this axiom show that why Russell’s paradox doesn’t happened here because if ou assume “ $R = \{x | x \notin x\}$ ” is a set, you can’t prove or disprove “ $R \in R$ ”. (we don’t need to prove that now. we will prove it in “**MetaLogic**”.)

4.2 definitions of set theory

Let’s talk about some definitions in set theoty. These definitions will come in handy later.

definition 4.2.1. Empty set also known as the null set, is the unique mathematical set containing no elements at all. (uniqueness can be proven.)

$$\emptyset := \{x \mid \perp\}$$

therefore.

$$\forall x (x \in \emptyset \leftrightarrow \perp)$$

therefore.

$$\forall x (x \notin \emptyset)$$

Example 4.2.1. $\therefore \forall X (\emptyset \subseteq X)$

1	$x \in \emptyset$	<i>acp</i>
2	$\forall x(x \notin \emptyset)$	<i>def</i>
3	$x \notin \emptyset$	$\forall E, 2$
4	$\perp$	$\perp I, 1, 3$
5	$x \in X$	$\perp E, 4$
6	$\emptyset \subseteq X$	$\subseteq I, 1, 5$
7	$\forall X(\emptyset \subseteq X)$	$\forall I, 6$

Lemma 4.2.1. $A = \emptyset \therefore / \therefore \forall y(y \notin A)$

1	$\forall y(y \notin A)$	<i>ass</i>
2	$y \in A$	<i>acp</i>
3	$y \notin A$	$\forall E, 1$
4	$\perp$	$\perp I, 2, 3$
5	$y \in \emptyset$	$\perp E, 4$
6	$y \in \emptyset$	<i>acp</i>
7	$\forall y(y \notin \emptyset)$	<i>def.</i>
8	$y \notin \emptyset$	$\forall E, 7$
9	$\perp$	$\perp I, 6, 8$
10	$y \in A$	$\perp E, 9$
11	$A = \emptyset$	$I =$

1	$A = \emptyset$	<i>ass</i>
2	$y \in A$	<i>acp</i>
3	$\forall y(y \notin \emptyset)$	<i>def.</i>
4	$y \in \emptyset$	$E =, 1, 2$
5	$y \notin \emptyset$	$\forall E, 3$
6	$\perp$	$\perp I, 4, 5$
7	$y \notin A$	$\neg I$
8	$\forall y(y \notin A)$	$\forall I, 7$

Theorem 4.2.1. (*schema*) $(I\emptyset \neq, E\emptyset \neq, \emptyset = I, \emptyset = E)$

$\frac{a \in A}{\therefore A \neq \emptyset} \quad I\emptyset \neq$	$\begin{array}{c} A \neq \emptyset \\ \left \begin{array}{c} x \in A \\ \vdots \\ \theta \end{array} \right. \\ \therefore \theta \quad E\emptyset \neq \end{array}$
$\begin{array}{c} \left \begin{array}{c} x \in A \\ \vdots \\ \perp \end{array} \right. \\ \therefore A = \emptyset \quad \emptyset = I \end{array}$	$\frac{A = \emptyset}{\therefore a \notin A} \quad \emptyset = E$

- 1 $a \in A$ *ass*
- 2 $\exists x(x \in A)$ $\exists I, 1$
- 3 $\neg(\forall x(x \notin A))$ *De Morgan 2*
- 4 $A \neq \emptyset$ *lemma 4.4.1, 3*

- 1 $\left| \begin{array}{c} x \in A \\ \vdots \\ \perp \end{array} \right.$ *acp*
- n
- $n+1$ $x \notin A$ $\neg I$
- $n+2$ $\forall x(x \notin A)$ $\forall I, n+1$
- $n+3$ $A = \emptyset$ *lemma 4.4.1, 1*

- 1 $A = \emptyset$ *ass*
- 2 $\forall x(x \notin A)$ *lemma 4.4.1, 1*
- 3 $a \notin A$ $\forall E2$

definition 4.2.2. pairing set

$$\{a_0, a_1, a_2, \dots, a_n\} := \{x \mid x = a_0 \vee x = a_1 \vee x = a_2 \vee \dots \vee x = a_n\}$$

therefore.

$$\forall x(x \in \{a_0, a_1, a_2, \dots, a_n\} \leftrightarrow (x = a_0 \vee x = a_1 \vee x = a_2 \vee \dots \vee x = a_n))$$

Theorem 4.2.2. (schema) $(I\{\}, E\{\})$

$\frac{a = a_i}{\therefore a \in \{a_0, a_1, a_2, \dots, a_n\}} \quad I\{\}$	$a \in \{a_0, a_1, a_2, \dots, a_n\}$ $\left \begin{array}{l} a = a_0 \\ \vdots \\ \theta \end{array} \right.$ $\vdots$ $\left \begin{array}{l} a = a_n \\ \vdots \\ \theta \end{array} \right.$ $\therefore \theta \quad E\{\}$
--	--

Proof Is Left As An Exercise To The Reader. 😊

definition 4.2.3. Union and Intersection

Intersection:

$$A \cap B := \{x \mid x \in A \wedge x \in B\}$$

therefore.

$$\forall x(x \in A \cap B \leftrightarrow (x \in A \wedge x \in B))$$

Union:

$$A \cup B := \{x \mid x \in A \vee x \in B\}$$

therefore.

$$\forall x(x \in A \cup B \leftrightarrow (x \in A \vee x \in B))$$

Theorem 4.2.3. (*schema*) $(\cap I, \cap E, \cup I, \cup E)$

$\frac{a \in A}{\therefore a \in A \cap B} \quad \cap I$	$\frac{a \in A \cap B}{\therefore a \in A} \quad \cap E \quad \frac{a \in A \cap B}{\therefore a \in B} \quad \cap E$
$\frac{a \in A}{\therefore a \in A \cup B} \quad \cup I$ $\frac{a \in A}{\therefore a \in B \cup A} \quad \cup I$	$\frac{a \in A \cup B}{\therefore \theta} \quad \cup E$

Proof Is Left As An Exercise To The Reader. 😊

definition 4.2.4. Big Union and Intersection

Intersection:

$$\bigcap A := \{x \mid \forall X (X \in A \rightarrow x \in X)\}$$

i.e.

$$\bigcap A := \{x \mid \forall X \in A \quad x \in X\}$$

therefore.

$$\forall x (x \in \bigcap A \leftrightarrow \forall X (X \in A \rightarrow x \in X))$$

Union:

$$\bigcup A := \{x \mid \exists X (X \in A \wedge x \in X)\}$$

i.e.

$$\bigcup A := \{x \mid \exists X \in A \quad x \in X\}$$

therefore.

$$\forall x (x \in \bigcup A \leftrightarrow (\exists X (X \in A \wedge x \in X)))$$

Theorem 4.2.4. (*schema*) $(\cap I, \cap E, \cup I, \cup E)$

$\begin{array}{c} \left \begin{array}{l} X \in A \\ \vdots \\ a \in X \end{array} \right. \\ \hline \therefore a \in \bigcap A \quad \bigcap I \end{array}$	$\begin{array}{c} a \in \bigcap A \\ B \in A \\ \hline \therefore a \in B \quad \bigcap E \end{array}$
$\begin{array}{c} a \in B \\ B \in A \\ \hline \therefore a \in \bigcup A \quad \bigcup I \end{array}$	$\begin{array}{c} a \in \bigcup A \\ \left \begin{array}{l} a \in X \\ X \in A \\ \vdots \\ \theta \end{array} \right. \\ \hline \therefore \theta \quad \bigcup E \end{array}$

Proof Is Left As An Exercise To The Reader. 😊

definition 4.2.5. Complement

$$A^c := \{x \mid x \notin A\}$$

therefore.

$$\forall x(x \in A^c \leftrightarrow x \notin A)$$

Theorem 4.2.5. (schema) $((\)^c I, (\)^c E)$

$\begin{array}{c} \left \begin{array}{l} a \in A \\ \vdots \\ \perp \end{array} \right. \\ \hline \therefore a \notin A \quad (\)^c I \end{array}$	$\begin{array}{c} \left \begin{array}{l} a \notin A \\ \vdots \\ \perp \end{array} \right. \\ \hline \therefore a \in A \quad (\)^c E \end{array}$
--	--

Proof Is Left As An Exercise To The Reader. 😊

Observation 4.2.1. *we can show Universal set with...*

$$V = \{x \mid \top\}$$

therefore.

$$\forall x(x \in V \leftrightarrow \top)$$

therefore.

$$\forall x(x \in V)$$

definition 4.2.6. Subtraction

$$A - B := \{x \mid x \in A \wedge x \notin B\}$$

therefore.

$$\forall x \left(x \in A - B \leftrightarrow (x \in A \wedge x \notin B) \right)$$

Theorem 4.2.6. (schema) $(-I, -E)$

$\frac{a \in A \quad a \notin B}{\therefore a \in A - B} \quad -I$	$\frac{a \in A - B}{\therefore a \in A} \quad -E \quad \frac{a \in A - B}{\therefore a \notin B} \quad -E$
--	--

Proof Is Left As An Exercise To The Reader. 😊

Lemma 4.2.2. $\therefore A - B = A \cap B^c$

Proof Is Left As An Exercise To The Reader. 😊

definition 4.2.7. Powerset

$$P(A) := \{X \mid X \subseteq A\}$$

therefore.

$$\forall X \left(X \in P(A) \leftrightarrow (X \subseteq A) \right)$$

Theorem 4.2.7. (schema) (PI, PE)

$\frac{A \subseteq B}{\therefore A \in P(B)} \quad PI$	$\frac{A \in P(B)}{\therefore A \subseteq B} \quad PE$
--	--

Proof Is Left As An Exercise To The Reader. 😊

This page intentionally left blank.

Chapter 5

Relations and functions

5.1 cartesian product

definition 5.1.1. *Ordered pair*

$$(x, y) := \{\{x\}, \{x, y\}\}.$$

definition 5.1.2. *Tuples* (n is a arbitrary number.) (this is a temporary definition, we will give a better one in the next chapter.)

$$(x_1, x_2, \dots, x_n) := ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Lemma 5.1.1. $\therefore (x_0, y_0) = (x_1, y_1) \leftrightarrow ((x_0 = x_1) \wedge (y_0 = y_1))$

Proof Is Left As An Exercise To The Reader. 😊

definition 5.1.3. *Cartesian product*


$$A \times B := \{u \mid \exists x, y ((x, y) = u \wedge x \in A \wedge y \in B)\}.$$

or i.e.

$$A \times B := \{(x, y) \mid x \in A \wedge y \in B\}.$$

Example 5.1.1. if $A = \{a, b, c\}$ and $B = \{0, 1, 2, 3\}$ then

$$A \times B = \{(a, 0), (a, 1), (a, 2), (a, 3), (b, 0), (b, 1), (b, 2), (b, 3), (c, 0), (c, 1), (c, 2), (c, 3)\}$$



Exercise 5.1.1. for sets A, B, C , prove.

1. $\therefore A \times (B \cap C) = (A \times B) \cap (A \times C)$
2. $\therefore A \times (B \cup C) = (A \times B) \cup (A \times C)$
3. $\therefore A \times (B - C) = (A \times B) - (A \times C)$

5.2 Relations

definition 5.2.1. Relation we call a set like R a **Relation** when there are two sets like X and Y s.t. $R \subseteq X \times Y$.

$$Rel(R) : \exists X, Y (R \subseteq X \times Y).$$


definition 5.2.2. Domain and Range(also called Image)

Domain of a set R is all x that there exists a y s.t. " $(x, y) \in R$ "

$$Dom(R) := \{x \mid \exists y \quad (x, y) \in R\}$$

Range(Image) of a set R is all y that there exists a x s.t. $(x, y) \in R$

$$Range(R) := \{y \mid \exists x \quad (x, y) \in R\}.$$



definition 5.2.3. Inverse of a Relation

$$R^{-1} = \{u \mid \exists x, y (u = (x, y) \wedge (y, x) \in R)\}$$

or i.e.

$$R^{-1} = \{(x, y) \mid (y, x) \in R\}$$

therefore.

$$\forall x, y ((x, y) \in R^{-1} \leftrightarrow (y, x) \in R)$$

definition 5.2.4. Combination of a Relations

$$S \circ R = \{u \mid \exists x, y (u = (x, y) \wedge \exists z ((x, z) \in R \wedge (z, y) \in S))\}$$

or i.e.

$$S \circ R = \{(x, y) \mid \exists z ((x, z) \in R \wedge (z, y) \in S)\}$$

therefore.

$$\forall x, y ((x, y) \in S \circ R \leftrightarrow \exists z ((x, z) \in R \wedge (z, y) \in S))$$

definition 5.2.5. Image

$$R[A] = \{y \mid \exists x \in A (x, y) \in R\}$$

you can also see by definition of “inverse of a Relation” and “image” this equation holds.

$$R^{-1}[B] = \{x \mid \exists y \in B (x, y) \in R\}$$

we cal this “**Pre-image**”

5.3 Functions

definition 5.3.1. Function we call a relation “**f**” a “**function**” when every input has exactly one output i.e.

$$\text{Func}(f) : \forall x, y_1, y_2 \left((x, y_1) \in f \wedge (x, y_2) \in f \rightarrow y_1 = y_2 \right).$$

definition 5.3.2. Output of a function assume a function **f** and $(x, y) \in f$, we define “ $f(x) := y$ ” and we call “ $f(x)$ ” a “**output of a function**”. and the reverse is also true, if we use “ $f(x)$ ”, we have a function “**f**” and we can say “ $x \in \text{Dom}(f)$ ”. “output of a function” is object too, so we can have $f(f(x))$ as well.

Attention 5.3.1. input of a function can be a ordered pair itself, so we can have $f((x, y))$ and it is a “**output of a function**”. we define “ $f(x, y) := f((x, y))$ ” for simplicity. this is true for tuples as well, so we can have “ $f((x_1, x_2, \dots, x_n))$ ” and we define “ $f(x_1, x_2, \dots, x_n) := f((x_1, x_2, \dots, x_n))$ ” for simplicity.

definition 5.3.3. Term we define term like this.

1. every obj is a term.
2. if $t_0, \dots, t_n$ are terms and “ $\text{Func}(f)$ ” and “ $(t_0, \dots, t_n) \in \text{Dom}(f)$ ”. then “ $f(t_0, \dots, t_n)$ ” is also a term.

we denote terms with small letters like $t, s, r, \dots$.

definition 5.3.4. Substitution for term we define Substitution an term with variable like this.

1. for variables if $x = y$ then $y[t/x] := t$ and if $x \neq y$ then $y[t/x] := y$.
2. for constants $c[t/x] := c$.
3. for terms like $f(t_0, \dots, t_n)$

$$f(t_0, \dots, t_n)[t/x] := f(t_0[t/x], \dots, t_n[t/x])$$

definition 5.3.5. Substitution for formulas we define Substitution for a formula like this.

1. for atomic formulas we define it like this. if it's a single term we do it as we explain in “**Substitution for term**”. if it's like $t_0 = t_1$ or $t_0 \in t_1$ (t_0, t_1 are terms). we define it like this.

$$(t_0 = t_1)[t/x] := (t_0[t/x] = t_1[t/x])$$

$$(t_0 \in t_1)[t/x] := (t_0[t/x] \in t_1[t/x])$$

2. if formula A is complex we define it like this.

$$(B \bigcirc C)[t/x] := (B[t/x] \bigcirc C[t/x])$$

($\bigcirc$ can be $\rightarrow, \wedge, \vee$).

$$(\neg B)[t/x] := (\neg B[t/x])$$

Theorem 5.3.1. (schema) $(\forall I, \forall E, \exists I, \exists E)$

Rules	Terms and Conditions
$\frac{\phi(x)}{\therefore \forall x\phi(x)} \quad \forall I$	1. x is a variable. 2. "$\phi(x)$" must not be assumed.
$\frac{\forall x\phi(x)}{\therefore \phi[t/x]} \quad \forall E$	1. t is a term.
$\frac{\phi[t/x]}{\therefore \exists x\phi(x)} \quad \exists I$	1. t is a term.
$\begin{array}{l} \exists x\phi(x) \\ \left \begin{array}{l} \phi(x) \\ \vdots \\ \theta \end{array} \right. \\ \therefore \theta \end{array} \quad \exists E$	1. x is a variable. 2. "$\phi(x)$" must not be assumed in previous lines. (except ones that are closed) 3. x is not free in θ

$(\forall I, \exists E)$ is the same as $(\forall I, \exists I)$ in chapter three so we have to prove $(\forall E, \exists I)$. so assume $t = f(x)$ so to it's easy to see $\text{Func}(f)$ and $x \in \text{Dom}(f)$.

1	$\forall x\phi(x)$	ass	1	$\phi(f(x))$	ass
2	$\text{Func}(f)$	ass	2	$\text{Func}(f)$	ass
3	$\exists y((x, y) \in f)$	ass	3	$\exists y((x, y) \in f)$	ass
4	$\left \begin{array}{l} (x, y) \in f \end{array} \right.$	acp	4	$\left \begin{array}{l} (x, y) \in f \end{array} \right.$	acp
5	$\left \begin{array}{l} f(x) = y \end{array} \right.$	def	5	$\left \begin{array}{l} f(x) = y \end{array} \right.$	def
6	$\left \begin{array}{l} \phi(y) \end{array} \right.$	$\forall E$	6	$\left \begin{array}{l} \phi(y) \end{array} \right.$	$= E$
7	$\left \begin{array}{l} \phi(f(x)) \end{array} \right.$	$= E$	7	$\left \begin{array}{l} \exists x\phi(x) \end{array} \right.$	$\exists I$
8	$\phi(f(x))$	$\exists E$	8	$\exists x\phi(x)$	$\exists E$

Attention 5.3.2. in all rules and theorems you can replace all constants like $a, b, c, \dots$ with terms like $t, s, r, \dots$ and rewrite them.

definition 5.3.6. Function from set to set we call a function f from A to B when it's domain is A and it's range is subset of B i.e.

$$(f : A \rightarrow B) : \text{Func}(f) \wedge \text{Dom}(f) = A \wedge \text{Range}(f) \subseteq B$$

definition 5.3.7. Injective function we call a function f injective when every output has at

most one input i.e.

$$\text{Injective}(f) : \forall y, x_1, x_2 \left((x_1, y) \in f \wedge (x_2, y) \in f \rightarrow x_1 = x_2 \right).$$

we call a function f injective from A to B when it's function from A to B and it's injective i.e.

$$(f : A \xrightarrow{\text{inj}} B) : (f : A \rightarrow B) \wedge \text{Injective}(f)$$

definition 5.3.8. Surjective function we call a function f from A to B when it's domain is A and it's range is equal to B i.e.

$$(f : A \xrightarrow[\text{surj}]{} B) : (f : A \rightarrow B) \wedge \text{Range}(f) = B$$

or i.e.

$$(f : A \xrightarrow[\text{surj}]{} B) : (f : A \rightarrow B) \wedge (\forall y \in B \quad \exists x \in A \quad (x, y) \in f)$$

definition 5.3.9. Bijection function we call a function f from A to B when it's domain is A and it's range is equal to B i.e.

$$(f : A \xrightarrow[\text{surj}]{\text{inj}} B) : (f : A \xrightarrow{\text{inj}} B) \wedge (f : A \xrightarrow[\text{surj}]{} B)$$

5.4 Family of sets

Attention 5.4.1. Indexes are objects (like numbers) allows us to give a unique "tag" to every object. we use $i, j, k, \dots$ for variable indexes and we use $m, n, l, \dots$ for constant indexes.

definition 5.4.1. Family of sets assume $A : I \rightarrow \mathcal{F}$ is a function from a nonempty set of indices I to a family of sets $\mathcal{F}$, then we call $(\text{Range}[A])$ an indexed family of sets. we can write A_i instead of $A(i)$ and we can write $(A_i)_{i \in I}$ instead of " A " and $\{A_i\}_{i \in I}$ instead of " $\text{Range}[A]$ " or " $A[I]$ ".

$$(A_i)_{i \in I} := A$$

and

$$\{A_i\}_{i \in I} := A[I]$$

definition 5.4.2. Intersection and union of an family of sets

$$\bigcap_{i \in I} A_i := \bigcap \{A_i\}_{i \in I}$$

$$\bigcup_{i \in I} A_i := \bigcup \{A_i\}_{i \in I}$$

Attention 5.4.2. we can write $\forall i \in I \quad P(i)$ instead of $\forall i (i \in I \rightarrow P(i))$ and we can write $\exists i \in I \quad P(i)$ instead of $\exists i (i \in I \wedge P(i))$.

Lemma 5.4.1.

$$/ \therefore \bigcap_{i \in I} A_i = \{x \mid \forall i \in I \quad x \in A_i\}$$

1	$x \in \bigcap_{i \in I} A_i$	<i>acp</i>
2	$x \in \bigcap \{A_i\}_{i \in I}$	<i>def.1</i>
3	$x \in \bigcap A[I]$	<i>def.1</i>
4	$\forall Y \in A[I] \quad x \in Y$	<i>def</i> $\bigcap$
5	$\forall Y ((\exists i \in I \quad A_i = Y) \rightarrow x \in Y)$	<i>def image</i>
6	$(\exists i \in I \quad A_i = A_{i'}) \rightarrow x \in A_{i'}$	$\forall E$
7	$i' \in I$	<i>acp</i>
8	$A_{i'} = A_{i'}$	$= I$
9	$i' \in I \wedge A_{i'} = A_{i'}$	$\wedge I$
10	$\exists i \in I \quad A_i = A_{i'}$	$\exists I$
11	$x \in A_{i'}$	$\rightarrow E$
12	$i' \in I \rightarrow x \in A_{i'}$	$\rightarrow I$
13	$\forall i \in I \quad x \in A_i$	$\forall I$
14	$x \in \{x \mid \forall i \in I \quad x \in A_i\}$	<i>def</i>
15	$x \in \{x \mid \forall i \in I \quad x \in A_i\}$	<i>acp</i>
16	$\forall i \in I \quad x \in A_i$	<i>def</i>
17	$\exists i \in I \quad A_i = Y$	<i>acp</i>
18	$i \in I$	<i>acp</i>
19	$x \in A_i$	$\forall E$
20	$A_i = Y$	$\wedge E$
21	$x \in Y$	$= E$
22	$x \in Y$	$\exists E$
23	$(\exists i \in I \quad A_i = Y) \rightarrow x \in Y$	$\rightarrow I$
24	$\forall Y ((\exists i \in I \quad A_i = Y) \rightarrow x \in Y)$	$\forall I$
25	$\forall Y \in A[I] \quad x \in Y$	<i>def</i>
26	$x \in \bigcap A[I]$	<i>def</i>
27	$x \in \bigcap \{A_i\}_{i \in I}$	<i>def</i>
28	$x \in \bigcap_{i \in I} A_i$	<i>def</i>
29	$\bigcap_{i \in I} A_i = \{x \mid \forall i \in I \quad x \in A_i\}$	$= I$

$$/ \therefore \bigcup_{i \in I} A_i = \{x \mid \exists i \in I \quad x \in A_i\}$$

1	$x \in \bigcup_{i \in I} A_i$	<i>acp</i>
2	$x \in \bigcup \{A_i\}_{i \in I}$	<i>def.1</i>
3	$x \in \bigcup A[I]$	<i>def.1</i>
4	$\exists Y \in A[I] \quad x \in Y$	<i>def</i> $\bigcup$
5	$\exists Y((\exists i \in I \quad A_i = Y) \wedge x \in Y)$	<i>def.image</i>
6	$x \in Y$	<i>acp</i>
7	$\exists i \in I \quad A_i = Y$	<i>acp</i>
8	$i \in I$	<i>acp</i>
9	$A_i = Y$	<i>acp</i>
10	$x \in A_i$	$= E$
11	$\exists i \in I \quad x \in A_i$	$\exists I$
12	$\exists i \in I \quad x \in A_i$	$\exists E$
13	$\exists i \in I \quad x \in A_i$	$\exists E$
14	$x \in \{x \mid \exists i \in I \quad x \in A_i\}$	<i>def</i>
15	$x \in \{x \mid \exists i \in I \quad x \in A_i\}$	<i>acp</i>
16	$\exists i \in I \quad x \in A_i$	<i>def</i>
17	$i' \in I$	<i>acp</i>
18	$x \in A_{i'}$	<i>acp</i>
19	$A_{i'} = A_{i'}$	$= I$
20	$\exists i \in I \quad A_i = A_{i'}$	$\exists I$
21	$(\exists i \in I \quad A_i = A_{i'}) \wedge x \in A_{i'}$	$\wedge I$
22	$\exists Y((\exists i \in I \quad A_i = Y) \wedge x \in Y)$	$\exists I$
23	$\exists Y((\exists i \in I \quad A_i = Y) \wedge x \in Y)$	$\exists E$
24	$\exists Y \in A[I] \quad x \in Y$	<i>def</i>
25	$x \in \bigcup A[I]$	<i>def</i>
26	$x \in \bigcup \{A_i\}_{i \in I}$	<i>def</i>
27	$x \in \bigcup_{i \in I} A_i$	<i>def</i>
28	$\bigcup_{i \in I} A_i = \{x \mid \exists i \in I \quad x \in A_i\}$	$= I$

Theorem 5.4.1. (*schema*) $(\bigcap_I I, \bigcap_I E, \bigcup_I I, \bigcup_I E)$

$\left \begin{array}{l} i \in I \\ \vdots \\ t \in A_i \end{array} \right.$ $\therefore t \in \bigcap_{i \in I} A_i \quad \bigcap_I I$	$t \in \bigcap_{i \in I} A_i$ $\frac{n \in I}{\therefore t \in A_n} \quad \bigcap_I E$
$t \in A_n$ $\frac{n \in I}{\therefore t \in \bigcup_{i \in I} A_i} \quad \bigcup_I I$	$t \in \bigcup_{i \in I} A_i$ $\left \begin{array}{l} i \in I \\ t \in A_i \\ \vdots \\ \theta \end{array} \right.$ $\therefore \theta \quad \bigcup_I E$

Proof Is Left As An Exercise To The Reader. 😊