(

LOGIN

SIGNUP

STUDY MODULES

Lectures

Doubts

Tests

STUDY TOOLS

Bytes

© toppr • About • Guides • Be an Educator • Blog • Terms

EXAMS > SCIENCE > SCHOOL COLLEGE & CAREER PEOPLE GENERAL Search posts, tags or authors

# **Dot Product of Two Vectors**



PUBLISHED O March 13th, 201

TAGS

MATHEMATICS

SCHOOL

#### **Dot Product of Two Vectors**

Many mathematical operations are usable on vectors. In this article, we will take a look at the dot product of two vectors.

Let's understand first that vectors can be multiplied by two methods:

- · scalar product of vectors or dot product
- vector product of vectors or cross product

The main differentiation between these two methods is the fact that we get a scalar value as resultant through the first method and the resultant obtained is also a vector in nature by using the second technique.

Note that in a two-dimensional Cartesian plane, vectors are denoted in terms of the x-coordinates and y-coordinates of their end points, assuming they start at the origin (x, y) = (0,0).

When it comes to polar coordinates, vectors can be denoted in terms of length (magnitude) and direction. But when they are written in this format, the dot product of two vectors is equal to the product of their lengths, multiplied by the cosine of the angle between them.

*Geometric interpretation.* Dot product of two vectors  $\bar{a}$  and b can be defined as a scalar quantity that is equal to the product of magnitudes of vectors multiplied by the cosine of the angle between vectors:

$$\bar{a} \cdot b = |\bar{a}| \cdot |b| \cos \alpha$$

OTHER MODILIES

Lectures

Doubts

Tests

STUDY TOOLS

Bytes

© toppr • About • Guides • Be an Educator • Blog • Terms

*Algebraic interpretation.* Dot product of two vectors ā and b is basically a scalar quantity that is equal to the sum of pair-wise products of coordinate vectors a and b.

**Dot product** is also known as *scalar product* or *inner product*.

## **Dot Product - Formulae**

### Dot product formula for plane problems

When it comes to the plane problem, the dot product of vectors  $\bar{a} = \{a_x; a_y\}$  and  $b = \{b_x; b_y\}$  can be determined with the help of the following formula:

$$\bar{\mathbf{a}} \cdot \mathbf{b} = \mathbf{a}_{\mathbf{x}} \cdot \mathbf{b}_{\mathbf{x}} + \mathbf{a}_{\mathbf{y}} \cdot \mathbf{b}_{\mathbf{y}}$$

## Dot product formula for spatial problems

The dot product of vectors  $\bar{a} = \{a_x; a_y; a_z\}$  and  $b = \{b_x; b_y; b_z\}$  can be figured out by using the following formula for the spatial problem:

$$\bar{\mathbf{a}} \cdot \mathbf{b} = \mathbf{a}_{\mathbf{x}} \cdot \mathbf{b}_{\mathbf{x}} + \mathbf{a}_{\mathbf{v}} \cdot \mathbf{b}_{\mathbf{v}} + \mathbf{a}_{\mathbf{z}} \cdot \mathbf{b}_{\mathbf{z}}$$

### Dot product formula for n dimensional space problems

Talking about the n dimensional space problem, the dot product of vectors  $\bar{a} = \{a_1; a_2; ...; a_n\}$  and  $b = \{b_1; b_2; ...; b_n\}$  can be easily determined with the following formula:

$$\bar{\mathbf{a}} \cdot \mathbf{b} = \mathbf{a}_1 \cdot \mathbf{b}_1 + \mathbf{a}_2 \cdot \mathbf{b}_2 + \dots + \mathbf{a}_n \cdot \mathbf{b}_n$$

# **Properties of Dot Product of Vector**

1. The dot product is always more than zero or equal to zero when it comes to a vector with itself:

$$\bar{a} \cdot \bar{a} > 0$$

**2.** The dot product of a vector with itself is zero only if the vector is the zero vector:

$$\bar{\mathbf{a}} \cdot \bar{\mathbf{a}} = 0 \Longleftrightarrow \bar{\mathbf{a}} = 0$$

**3.** The dot product of a vector with itself is equal to the square of its magnitude:

$$\bar{\mathbf{a}} \cdot \bar{\mathbf{a}} = |\bar{\mathbf{a}}|^2$$

**4.** The dot product operation is communicative:

$$\bar{a} \cdot b = b \cdot \bar{a}$$

**5.** If the dot product of two not zero vectors is zero, then note that these vectors are orthogonal:

$$\bar{a} \neq 0$$
,  $b \neq 0$ ,  $\bar{a} \cdot b = 0 \iff \bar{a} \perp$ 

**6.** 
$$(\alpha \bar{a}) \cdot b = \alpha(\bar{a} \cdot b)$$

OTUDY MODULES

Lectures

Doubts

Tests

STUDY TOOLS

Bytes

© toppr • About • Guides • Be an Educator • Blog • Terms

**7.** The dot product operation is distributive:

$$(\bar{a} + b) \cdot c = \bar{a} \cdot c + \cdot c$$

# **Solved Examples**

*Example 1:* Let's assume there are two vectors [6,2,-1] and [5,-8,2]. Now, determine the dot product of the vectors.

*Solution:* Given vectors: [6,2,-1] and [5,-8,2] be a and b respectively.

ā. 
$$b = (6)(5)+(2)(-8)+(-1)(2)$$

$$\bar{a}$$
.  $b = 30-16-2$ 

*Example 2:* Find the dot product of the two vectors |a|=4 and |b|=2 and  $\Theta=60$ .

*Solution:* a.b=|a||b|cos 60

a.b=4.2cos60

a.b=4

*Example 3.* What is the dot product of vectors  $\bar{a} = \{1, 2\}$  and  $b = \{4, 8\}$ ?

**Solution:**  $\bar{a} \cdot b = 1 \cdot 4 + 2 \cdot 8 = 4 + 16 = 20$ .

**Example 4.** What is the dot product of vectors a and b, if their magnitudes is |a| = 3, |b| = 6, and the angle between the vectors is equal to  $60^{\circ}$ .

**Solution:**  $\bar{a} \cdot b = |\bar{a}| \cdot |b| \cos \alpha = 3 \cdot 6 \cdot \cos 60^\circ = 9$ .

**Example 5.** If their magnitudes is |a| = 3, |b| = 2, and the angle between the vectors a and b is equal to  $60^{\circ}$ , what is the dot product of vectors p = a + 3b and q = 5a - 3b?

Solution:

p. q = (
$$\bar{a}$$
 + 3b ) · (5 $\bar{a}$  - 3b ) = 5 $\bar{a}$  ·  $\bar{a}$  - 3 $\bar{a}$  · b + 15 b ·  $\bar{a}$  - 9 b · b  
= 5 |  $\bar{a}$ |<sup>2</sup> + 12  $\bar{a}$  · b - 9 |b|<sup>2</sup> = 5 · 3<sup>2</sup> + 12 · 3 · 2 · cos 60° - 9 · 2<sup>2</sup> = 45 + 36 - 36 = 45.

Example of the dot product of vectors for spatial problems

*Example 6.* Find the dot product of vectors  $a = \{1; 2; -5\}$  and  $b = \{4; 8; 1\}$ .

**Solution:**  $\bar{a} \cdot b = 1 \cdot 4 + 2 \cdot 8 + (-5) \cdot 1 = 4 + 16 - 5 = 15$ .

Example of the dot product of vectors for n dimensional space problems

**Example 7.** What is the dot product of vectors  $a = \{1; 2; -5; 2\}$  and  $b = \{4; 8; 1; -2\}$ ?

STUDY MODULE

Lectures

Doubts
Tests

STUDY TOOLS

Bytes

© toppr • About • Guides • Be an Educator • Blog • Terms

**Solution:**  $\bar{a} \cdot b = 1 \cdot 4 + 2 \cdot 8 + (-5) \cdot 1 + 2 \cdot (-2) = 4 + 16 - 5 - 4 = 11$ .

Hope you are now thorough with the concept of the dot product of two vectors. To know more about vectors, visit our website Toppr.

## You may also like:

# Basics of Vector (Mathematics and Physics) Explained

Vedic Maths: Advantages Of Knowing the Basics

#### Order of Matrix and Matrix Multiplication


#### SHARE THIS POST

f y s

#### ABOUT THE AUTHOR



# 1 Comment Sort by Oldest





#### Aridio Silva

I think is missing a graphical visualization of the dot product in both 2-D (plane problems) and 3 dimensions (spatial problems).

Post

Reply 1m

Facebook Comments Plugin