

Cached quotients and lookups

Ariel Gabizon
Zeta Function Technologies

29. august 2023

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraint approach:

Prover sends b_0, \dots, b_{n-1} . Shows:

- ▶ $\forall i, b_i \in \{0, 1\}$
- ▶ $\sum_i b_i 2^i = x.$

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraint approach:

Prover sends b_0, \dots, b_{n-1} . Shows:

► $\forall i, b_i \in \{0, 1\}$

► $\sum_i b_i 2^i = x.$

Requires $n + 1$ constraints.

Lookup approach

Preprocess table $T = \{0, \dots, 2^n - 1\}$. Let $N := |T|$.
Devise protocol to check $x \in T$.

Lookup approach

Preprocess table $T = \{0, \dots, 2^n - 1\}$. Let $N := |T|$.
Devise protocol to check $x \in T$.

Old results - good when amortized:

Thm[plookup]: Can check m different x 's are in T
in $O(m + N)$ constraints.

Lookup approach

Preprocess table $T = \{0, \dots, 2^n - 1\}$. Let $N := |T|$.
Devise protocol to check $x \in T$.

Old results - good when amortized:

Thm[plookup]: Can check m different x 's are in T
in $O(m + N)$ constraints.

New results - prover doesn't pay for table size!!

Thm [Caulk...cq]: After $O(N \log N)$
preprocessing, can check $x \in T$, in $O(1)$
constraints.

Rest of talk: explain main technical component
of new works - *cached quotients*

Rest of talk: explain main technical component
of new works - *cached quotients*

*First - a brief recap of polynomial commitment
schemes..*

The KZG Polynomial commitment scheme

\mathbf{G} - generator of pairing friendly elliptic curve group.

$\text{srs} := 1 \cdot \mathbf{G}, x \cdot \mathbf{G}, \dots, x^d \cdot \mathbf{G}$, for random $x \in \mathbb{F}$.

The KZG Polynomial commitment scheme

\mathbf{G} - generator of pairing friendly elliptic curve group.

$\mathbf{srs} := \mathbf{1} \cdot \mathbf{G}, \mathbf{x} \cdot \mathbf{G}, \dots, \mathbf{x}^d \cdot \mathbf{G}$, for random $\mathbf{x} \in \mathbb{F}$.

For $\mathbf{f} \in \mathbb{F}[\mathbf{X}]$ of degree \mathbf{d} :

$$\mathbf{cm}(\mathbf{f}) := \mathbf{f}(\mathbf{x}) \cdot \mathbf{G}$$

The KZG Polynomial commitment scheme

\mathbf{G} - generator of pairing friendly elliptic curve group.

$\mathbf{srs} := \mathbf{1} \cdot \mathbf{G}, \mathbf{x} \cdot \mathbf{G}, \dots, \mathbf{x}^d \cdot \mathbf{G}$, for random $\mathbf{x} \in \mathbb{F}$.

For $\mathbf{f} \in \mathbb{F}[\mathbf{X}]$ of degree \mathbf{d} :

$$\mathbf{cm}(\mathbf{f}) := \mathbf{f}(\mathbf{x}) \cdot \mathbf{G}$$

Central Feature: Given $\mathbf{cm}(\mathbf{f})$ and any $\mathbf{a} \in \mathbb{F}$;
there is short proof for correctness of $\mathbf{z} = \mathbf{f}(\mathbf{a})$.

The KZG Polynomial commitment scheme

srs $:= 1 \cdot \mathbf{G}, x \cdot \mathbf{G}, \dots, x^d \cdot \mathbf{G}$, for random $x \in \mathbb{F}$.

cm(f) $:= f(x) \cdot \mathbf{G}$

Nice features:

The KZG Polynomial commitment scheme

srs $:= 1 \cdot \mathbf{G}, \mathbf{x} \cdot \mathbf{G}, \dots, \mathbf{x}^d \cdot \mathbf{G}$, for random $\mathbf{x} \in \mathbb{F}$.

cm(f) $:= f(\mathbf{x}) \cdot \mathbf{G}$

Nice features:

► **Linearity:** **cm**(f + g) = **cm**(f) + **cm**(g)

The KZG Polynomial commitment scheme

$\text{srs} := \mathbf{1} \cdot \mathbf{G}, \mathbf{x} \cdot \mathbf{G}, \dots, \mathbf{x}^d \cdot \mathbf{G}$, for random $\mathbf{x} \in \mathbb{F}$.

$\text{cm}(\mathbf{f}) := \mathbf{f}(\mathbf{x}) \cdot \mathbf{G}$

Nice features:

- ▶ **Linearity:** $\text{cm}(\mathbf{f} + \mathbf{g}) = \text{cm}(\mathbf{f}) + \text{cm}(\mathbf{g})$
- ▶ **Product checks:** Given $\text{cm}(\mathbf{f}_1), \text{cm}(\mathbf{f}_2), \text{cm}(\mathbf{g}_1), \text{cm}(\mathbf{g}_2)$ can check $\mathbf{f}_1(\mathbf{X})\mathbf{f}_2(\mathbf{X}) \stackrel{?}{=} \mathbf{g}_1(\mathbf{X})\mathbf{g}_2(\mathbf{X})$ via pairings.
(Secure in the Algebraic Group Model)

Cached Quotients - a motivating example
from **Caulk**[ZBKMN]

Cached Quotients - a motivating example from **Caulk**[ZBKMN]

$Z_T(X) = \prod_{a \in T} (X - a)$ a vanishing polynomial of
a subset $T \subset \mathbb{F}$.

Cached Quotients - a motivating example from **Caulk**[ZBKMN]

$Z_T(X) = \prod_{a \in T} (X - a)$ a vanishing polynomial of
a subset $T \subset \mathbb{F}$.

$\mathbf{cm}(Z_T), \mathbf{cm}(f)$ given to verifier.

Cached Quotients - a motivating example from **Caulk**[ZBKMN]

$Z_T(X) = \prod_{a \in T} (X - a)$ a vanishing polynomial of
a subset $T \subset \mathbb{F}$.

$\mathbf{cm}(Z_T), \mathbf{cm}(\mathbf{f})$ given to verifier.

Prover wants to show $\mathbf{f} = Z_S$ for some $S \subset T$.

Cached Quotients - a motivating example from **Caulk**[ZBKMN]

$Z_T(X) = \prod_{a \in T} (X - a)$ a vanishing polynomial of
a subset $T \subset \mathbb{F}$.

$\text{cm}(Z_T), \text{cm}(f)$ given to verifier.

Prover wants to show $f = Z_S$ for some $S \subset T$.

Can we do this in $O(|S|)$ prover operations?(think
 $|S| \ll |T|$)

Cached quotients idea:

The quotient $\mathbf{Z}_{T \setminus S}(\mathbf{X}) = \frac{\mathbf{Z}_T(\mathbf{X})}{\mathbf{Z}_S(\mathbf{X})}$ is a “witness” to $S \subset T$.

Cached quotients idea:

The quotient $\mathbf{Z}_{T \setminus S}(\mathbf{X}) = \frac{\mathbf{Z}_T(\mathbf{X})}{\mathbf{Z}_S(\mathbf{X})}$ is a “witness” to $S \subset T$.

- Enough to compute **commitment** to $\mathbf{Z}_{T \setminus S}$.

Cached quotients idea:

The quotient $\mathbf{Z}_{T \setminus S}(\mathbf{X}) = \frac{\mathbf{Z}_T(\mathbf{X})}{\mathbf{Z}_S(\mathbf{X})}$ is a “witness” to $S \subset T$.

- ▶ Enough to compute **commitment** to $\mathbf{Z}_{T \setminus S}$.
- ▶ This commitment is a **sparse combination** of commitments we can **precompute**.

details in next slide..

For each $i \in T$, let $g_i(X) := Z_{T \setminus \{i\}}(X)$.

For each $\mathbf{i} \in \mathbf{T}$, let $\mathbf{g}_{\mathbf{i}}(\mathbf{X}) := \mathbf{Z}_{\mathbf{T} \setminus \{\mathbf{i}\}}(\mathbf{X})$.

We have [Tomescu et. al]

$$\mathbf{Z}_{\mathbf{T} \setminus \mathbf{S}}(\mathbf{X}) = \sum_{\mathbf{i} \in \mathbf{S}} \mathbf{c}_{\mathbf{i}} \cdot \mathbf{g}_{\mathbf{i}}(\mathbf{X})$$

for some $\mathbf{c}_{\mathbf{i}} \in \mathbb{F}$.

For each $i \in T$, let $g_i(X) := Z_{T \setminus \{i\}}(X)$.

We have [Tomescu et. al]

$$Z_{T \setminus S}(X) = \sum_{i \in S} c_i \cdot g_i(X)$$

for some $c_i \in \mathbb{F}$.

We precompute $cm(Z_T), \{cm(g_i)\}_{i \in T}$.

Prover then computes in $|S|$ operations:

$$\pi := \mathbf{cm}(Z_{T \setminus S}) = \sum_{i \in S} c_i \cdot \mathbf{cm}(g_i)$$

Prover then computes in $|S|$ operations:

$$\pi := \mathbf{cm}(Z_{T \setminus S}) = \sum_{i \in S} c_i \cdot \mathbf{cm}(g_i)$$

Verifier checks with pairing that:

$$e(\mathbf{cm}(f), \pi) = e(\mathbf{cm}(Z_T), 1 \cdot G)$$

Historical perspective: A trilogy of pairing-based SNARKs

Historical perspective: A trilogy of pairing-based SNARKs

1. **A new hope (for SNARKs, not the universe) - [Groth10, **GGPR**, ..., Groth16]**

Historical perspective: A trilogy of pairing-based SNARKs

1. **A new hope (for SNARKs, not the universe) - [Groth10, **GGPR**, ..., Groth16]**
2. **The polynomial commitment scheme strikes back - [vsq1, **Sonic**, Plonk, Marlin, ...]**

Historical perspective: A trilogy of pairing-based SNARKs

1. **A new hope (for SNARKs, not the universe) - [Groth10, **GGPR**, ..., Groth16]**
2. **The polynomial commitment scheme strikes back - [vsq1, **Sonic**, Plonk, Marlin, ...]**
3. **Return of the pairing - [Caulk, ..., cq, ...]**

Other Application in Chapter 3: lincheck

Fixed $n \times n$ matrix M .

Other Application in Chapter 3: lincheck

Fixed $\mathbf{n} \times \mathbf{n}$ matrix \mathbf{M} .

Prover has poly $\mathbf{f} \in \mathbb{F}_{\langle \mathbf{n} \rangle}[\mathbf{X}]$. Verifier $\mathbf{cm}(\mathbf{f})$.

$\mathbf{a} := \mathbf{f}|_{\mathbf{H}}$ for subgroup \mathbf{H} of size \mathbf{n} .

Other Application in Chapter 3: lincheck

Fixed $n \times n$ matrix M .

Prover has poly $f \in \mathbb{F}_{<n}[X]$. Verifier $cm(f)$.

$\alpha := f|_H$ for subgroup H of size n .

cq-lin: After preprocessing of M , prover can show $M \cdot \alpha = 0$ in $O(n)$ operations.

Other Application in Chapter 3: lincheck

Fixed $n \times n$ matrix M .

Prover has poly $f \in \mathbb{F}_{<n}[X]$. Verifier $cm(f)$.

$\alpha := f|_H$ for subgroup H of size n .

cq-lin: After preprocessing of M , prover can show $M \cdot \alpha = 0$ in $O(n)$ operations.