

cq: Cached quotients for fast lookups

Liam Eagen Dario Fiore **Ariel Gabizon**

13. januar 2023


Outline

- ▶ PCS/KZG review

Outline

- ▶ PCS/KZG review
- ▶ KZG shenanigans
 1. Committing to sparse polys.
 2. “cached quotients”

Outline

- ▶ PCS/KZG review
- ▶ KZG shenanigans
 1. Committing to sparse polys.
 2. “cached quotients”
- ▶ Lookups
 1. Motivation
 2. log derivative protocol[Eagen, Haböck,..]
 3. 

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\mathbf{cm}(\mathbf{f})$ to polynomial.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\mathbf{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $i \in \mathbb{F}$.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\mathbf{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $\mathbf{i} \in \mathbb{F}$.
- ▶ Prover sends back $\mathbf{z} = \mathbf{f}(\mathbf{i})$; together with proof $\mathbf{open}(\mathbf{f}, \mathbf{i})$ that \mathbf{z} is correct.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\mathbf{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $\mathbf{i} \in \mathbb{F}$.
- ▶ Prover sends back $\mathbf{z} = \mathbf{f}(\mathbf{i})$; together with proof $\mathbf{open}(\mathbf{f}, \mathbf{i})$ that \mathbf{z} is correct.

KZG give us PCS with commitments and openings are practically 32-48 bytes.

Notation: $[\mathbf{x}] = \mathbf{x} \cdot \mathbf{g}$ where \mathbf{g} generator of (an additive) elliptic curve group.

KZG10:

srs := $[1], [\mathfrak{x}], \dots, [\mathfrak{x}^d]$, for random $\mathfrak{x} \in \mathbb{F}$.

KZG10:

srs $:= [\mathbf{1}], [\mathbf{x}], \dots, [\mathbf{x}^d]$, for random $\mathbf{x} \in \mathbb{F}$.

cm(**f**) $:= [\mathbf{f}(\mathbf{x})]$

KZG10:

srs $:= [\mathbf{1}], [\mathbf{x}], \dots, [\mathbf{x}^d]$, for random $\mathbf{x} \in \mathbb{F}$.

cm(\mathbf{f}) $:= [\mathbf{f}(\mathbf{x})]$

open(\mathbf{f}, \mathbf{i}) $:= [\mathbf{h}(\mathbf{x})]$, where $\mathbf{h}(\mathbf{X}) := \frac{\mathbf{f}(\mathbf{X}) - \mathbf{f}(\mathbf{i})}{\mathbf{X} - \mathbf{i}}$

KZG10:

srs $:= [\mathbf{1}], [\mathbf{x}], \dots, [\mathbf{x}^d]$, for random $\mathbf{x} \in \mathbb{F}$.

cm(\mathbf{f}) $:= [\mathbf{f}(\mathbf{x})]$

open(\mathbf{f}, \mathbf{i}) $:= [\mathbf{h}(\mathbf{x})]$, where $\mathbf{h}(\mathbf{X}) := \frac{\mathbf{f}(\mathbf{X}) - \mathbf{f}(\mathbf{i})}{\mathbf{X} - \mathbf{i}}$

verify($\mathbf{cm}, \pi, \mathbf{z}, \mathbf{i}$) :

$$e(\mathbf{cm} - [\mathbf{z}], [\mathbf{1}]) \stackrel{?}{=} e(\pi, [\mathbf{x} - \mathbf{i}])$$

Shenanigan #1: Committing to sparse polys

notation: parameters $n \ll N$, $d := N - 1$.

$\mathbb{V} = \{\omega, \dots, \omega^N\} \subset \mathbb{F}$ subgroup of size N .

Shenanigan #1: Committing to sparse polys

notation: parameters $n \ll N$, $d := N - 1$.

$\mathbb{V} = \{\omega, \dots, \omega^N\} \subset \mathbb{F}$ subgroup of size N .

Say $A \in \mathbb{F}_{<N}[X]$ is n -sparse if has at most n non-zeroes on \mathbb{V} .

Shenanigan #1: Committing to sparse polys

notation: parameters $n \ll N$, $d := N - 1$.

$\mathbb{V} = \{\omega, \dots, \omega^N\} \subset \mathbb{F}$ subgroup of size N .

Say $\mathbf{A} \in \mathbb{F}_{<N}[\mathbf{X}]$ is n -sparse if has at most n non-zeroes on \mathbb{V} .

For $i \in [N]$, denote $\mathbf{A}_i := \mathbf{A}(\omega^i)$

$\mathbf{L}_1(\mathbf{X}), \dots, \mathbf{L}_N(\mathbf{X})$ - Lagrange basis of \mathbb{V} -
 $(\mathbf{L}_i)_j = \mathbf{0}$ when $i \neq j$.

Committing to sparse polys

From $\mathbf{srs} := [\mathbf{1}], [\mathbf{x}], \dots, [\mathbf{x}^d]$, we can precompute in $\mathbf{O}(\mathbf{N} \log \mathbf{N})$ operations the KZG commitments of \mathbb{V} 's Lagrange Base:

$$\mathbf{srs}_L := \{[\mathbf{L}_1(\mathbf{x})], \dots, [\mathbf{L}_N(\mathbf{x})]\}$$

Committing to sparse polys

From $\mathbf{srs} := [\mathbf{1}], [\mathbf{x}], \dots, [\mathbf{x}^d]$, we can precompute in $O(N \log N)$ operations the KZG commitments of \mathbb{V} 's Lagrange Base:

$$\mathbf{srs}_L := \{[\mathbf{L}_1(\mathbf{x})], \dots, [\mathbf{L}_N(\mathbf{x})]\}$$

Now for n -sparse $\mathbf{A}(\mathbf{X})$ of degree $< N$ compute

$$\mathbf{cm}(\mathbf{A}) = [\mathbf{A}(\mathbf{x})] = \sum_{i \in [N], \mathbf{A}_i \neq 0} \mathbf{A}_i \cdot [\mathbf{L}_i(\mathbf{x})]$$

Shenanigan #2: “Cached quotients” method

Scenario: $\mathbf{T}(\mathbf{X}) \in \mathbb{F}_{<\mathbf{N}}[\mathbf{X}]$ preprocessed poly.

$\mathbf{Z}_{\mathbb{V}}(\mathbf{X})$ -vanishing poly of \mathbb{V} .

Input: \mathbf{n} -sparse $\mathbf{A}(\mathbf{X}) \in \mathbb{F}_{<\mathbf{N}}[\mathbf{X}]$.

Shenanigan #2: “Cached quotients” method

Scenario: $\mathbf{T}(\mathbf{X}) \in \mathbb{F}_{<\mathbf{N}}[\mathbf{X}]$ preprocessed poly.

$Z_{\mathbb{V}}(\mathbf{X})$ -vanishing poly of \mathbb{V} .

Input: \mathbf{n} -sparse $\mathbf{A}(\mathbf{X}) \in \mathbb{F}_{<\mathbf{N}}[\mathbf{X}]$.

\mathbf{V} has $\mathbf{cm}(\mathbf{A})$. Want to prove to \mathbf{V} that:

$Z_{\mathbf{V}}(\mathbf{X})$ divides $\mathbf{A}(\mathbf{X})\mathbf{T}(\mathbf{X})$ *using $\mathbf{O}(\mathbf{n})$ prover operations.*

“Cached quotients” method

There exists quotient $Q_A(X)$ such that
 $A \cdot T \equiv Z_V \cdot Q_A.$

“Cached quotients” method

There exists quotient $\mathbf{Q}_A(\mathbf{X})$ such that
 $\mathbf{A} \cdot \mathbf{T} \equiv \mathbf{Z}_V \cdot \mathbf{Q}_A.$

We'll compute $[\mathbf{Q}_A(\mathbf{x})]$ in $\mathbf{O}(\mathbf{n})$ operations:

“Cached quotients” method

There exists quotient $Q_A(\mathbf{X})$ such that
 $\mathbf{A} \cdot \mathbf{T} \equiv \mathbf{Z}_V \cdot Q_A$.

We'll compute $[Q_A(\mathbf{x})]$ in $O(n)$ operations:

preprocessing: For each $i \in [N]$, compute $[Q_i(\mathbf{x})]$
such that for some $\mathbf{R}_i(\mathbf{X}) \in \mathbb{F}_{<N}[\mathbf{X}]$

$$\mathbf{L}_i(\mathbf{X}) \cdot \mathbf{T}(\mathbf{X}) = Q_i(\mathbf{X}) \cdot \mathbf{Z}_V(\mathbf{X}) + \mathbf{R}_i(\mathbf{X})$$

“Cached quotients” method

There exists quotient $Q_A(\mathbf{X})$ such that
 $\mathbf{A} \cdot \mathbf{T} \equiv \mathbf{Z}_V \cdot Q_A$.

We'll compute $[Q_A(\mathbf{x})]$ in $\mathbf{O}(\mathbf{n})$ operations:

preprocessing: For each $\mathbf{i} \in [\mathbf{N}]$, compute $[Q_i(\mathbf{x})]$
such that for some $\mathbf{R}_i(\mathbf{X}) \in \mathbb{F}_{\prec \mathbf{N}}[\mathbf{X}]$

$$\mathbf{L}_i(\mathbf{X}) \cdot \mathbf{T}(\mathbf{X}) = Q_i(\mathbf{X}) \cdot \mathbf{Z}_V(\mathbf{X}) + \mathbf{R}_i(\mathbf{X})$$

Also precompute $[\mathbf{Z}_V(\mathbf{x})], [\mathbf{T}(\mathbf{x})]$

“Cached quotients” method

After preprocessing, prover can compute

$$[Q_A(x)] = \sum_{i \in [N], A_i \neq 0} A_i \cdot [Q_i(x)]$$

“Cached quotients” method

After preprocessing, prover can compute

$$[Q_A(\mathbf{x})] = \sum_{i \in [N], A_i \neq 0} A_i \cdot [Q_i(\mathbf{x})]$$

Verifier can check:

$$e([A(\mathbf{x})], [T(\mathbf{x})]) = e([Q_A(\mathbf{x})], [Z_V(\mathbf{x})])$$

“Cached quotients” method

After preprocessing, prover can compute

$$[Q_A(x)] = \sum_{i \in [N], A_i \neq 0} A_i \cdot [Q_i(x)]$$

Verifier can check:

$$e([A(x)], [T(x)]) = e([Q_A(x)], [Z_V(x)])$$

In algebraic group model [FKL] can prove this is sound.

Lookup protocols

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraint approach: \mathcal{P} sends x_0, \dots, x_{n-1} Proves

- ▶ $\forall i, x_i \in \{0, 1\}$
- ▶ $\sum_i x_i 2^i = x.$

Constraints vs Lookups

Example: Check $0 \leq x \leq 2^n - 1$

Constraint approach: \mathcal{P} sends x_0, \dots, x_{n-1} Proves

- ▶ $\forall i, x_i \in \{0, 1\}$
- ▶ $\sum_i x_i 2^i = x.$

Requires $n + 1$ “gates”.

Lookup approach

Preprocess table $\mathbf{T} = \{\mathbf{0}, \dots, \mathbf{2}^n - \mathbf{1}\}$ Devise protocol to check $\mathbf{x} \in \mathbf{T}$.

Lookup approach

Preprocess table $\mathbf{T} = \{\mathbf{0}, \dots, \mathbf{2}^n - \mathbf{1}\}$ Devise protocol to check $\mathbf{x} \in \mathbf{T}$.

Thm-informal [Arya..plookup]: Check can be done in amortized $\mathbf{O}(\mathbf{1})$ constraints per check, when have $\mathbf{O}(|\mathbf{T}|)$ checks.

Lookup approach

Preprocess table $\mathbf{T} = \{0, \dots, 2^n - 1\}$ Devise protocol to check $x \in \mathbf{T}$.

Thm-informal [Arya..lookup]: Check can be done in amortized $\mathbf{O}(1)$ constraints per check, when have $\mathbf{O}(|\mathbf{T}|)$ checks.

Thm [Caulk..cq]: Can be done in $\mathbf{O}(1)$ constraints without need of amortization!

The question in polynomials:

Preprocessed: $V \subset \mathbb{F}$ subgroup of size N . $H \subset \mathbb{F}$ subgroup of size n . $T \in \mathbb{F}_{<N}[X]$.

The question in polynomials:

Preprocessed: $V \subset \mathbb{F}$ subgroup of size N . $H \subset \mathbb{F}$ subgroup of size n . $T \in \mathbb{F}_{<N}[X]$.

Input: $f \in \mathbb{F}_{<n}[X]$. $\mathbf{cm}(f)$ given to V .

The question in polynomials:

Preprocessed: $V \subset \mathbb{F}$ subgroup of size N . $H \subset \mathbb{F}$ subgroup of size n . $T \in \mathbb{F}_{<N}[X]$.

Input: $f \in \mathbb{F}_{<n}[X]$. $\mathbf{cm}(f)$ given to V .

Want to convince V that $f|_H \subset T|_V$ in $O(n)$ prover operations.

Log-derivative approach:

Lemma[Haböck]: $f|_H \subset T|_V$ if and only if there exists $m(X) \in \mathbb{F}_{<N}[X]$ s.t. as rational functions

$$\sum_{i \in [N]} \frac{m_i}{X + T_i} = \sum_{a \in H} \frac{1}{X + f(a)}$$

Log-derivative approach:

Lemma[Haböck]: $f|_H \subset T|_V$ if and only if there exists $\mathbf{m}(\mathbf{X}) \in \mathbb{F}_{\langle \mathbf{N} \rangle}[\mathbf{X}]$ s.t. as rational functions

$$\sum_{i \in [\mathbf{N}]} \frac{\mathbf{m}_i}{\mathbf{X} + \mathbf{T}_i} = \sum_{\mathbf{a} \in H} \frac{1}{\mathbf{X} + \mathbf{f}(\mathbf{a})}$$

Strategy: check this identity at random $\beta \in \mathbb{F}$.

Main prover task: Compute polynomial $\mathbf{A}(\mathbf{X})$ that interpolates RHS on \mathbb{V} , and prove it correct:

$$\mathbf{A}_i = \frac{\mathbf{m}_i}{\beta + \mathbf{T}_i}, \forall i \in [\mathbf{N}]$$

Main prover task: Compute polynomial $\mathbf{A}(\mathbf{X})$ that interpolates RHS on \mathbb{V} , and prove it correct:

$$A_i = \frac{m_i}{\beta + T_i}, \forall i \in [N]$$

Can be done via the “KZG shenanigans” we described before.

Main prover task: Compute polynomial $\mathbf{A}(\mathbf{X})$ that interpolates RHS on \mathbb{V} , and prove it correct:

$$\mathbf{A}_i = \frac{\mathbf{m}_i}{\beta + \mathbf{T}_i}, \forall i \in [\mathbf{N}]$$

Can be done via the “KZG shenanigans” we described before.

Must compute $[\mathbf{Q}_\mathbf{A}(\mathbf{x})]$ where

$$\mathbf{A}(\mathbf{X})(\beta + \mathbf{T}(\mathbf{X})) - \mathbf{m}(\mathbf{X}) = \mathbf{Q}_\mathbf{A}(\mathbf{X})\mathbf{Z}_\mathbb{V}(\mathbf{X}).$$