

fflonk: cheaply opening many polynomials using the fast-fourier equation

Ariel Gabizon (based on work with Zac Williamson)



Motivation: Save gas on Ethereum

- ▶ Recent polynomial IOP based snarks (Sonic, Marlin, Plonk,...) verification consists of two pairings and some number t of \mathbf{G}_1 scalar multiplications.

Motivation: Save gas on Ethereum

- ▶ Recent polynomial IOP based snarks (Sonic, Marlin, Plonk,..) verification consists of two pairings and some number t of \mathbf{G}_1 scalar multiplications.
- ▶ Each \mathbf{G}_1 scalar mult - 6000 gas 5usd.
- ▶ That's 76800usd per scalar mult when doing an on chain proof once in half an our per scalar mult.

Motivation: Save gas on Ethereum

- ▶ Recent polynomial IOP based snarks (Sonic, Marlin, Plonk,...) verification consists of two pairings and some number t of G_1 scalar multiplications.
- ▶ Each G_1 scalar mult - 6000 gas 5usd.
- ▶ That's 76800usd per scalar mult when doing an on chain proof once in half an our per scalar mult.
- ▶ **This work:** Getting t down from 16 to 5 in plonk (at the cost of tripling prover time) .

*snark verification reduces to polynomial
commitment scheme (PCS) opening verification -*

a 3 minute reminder on the
Kate-Zaverucha-Goldberg PCS

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\text{cm}(\mathbf{f})$ to polynomial.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\text{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $s \in \mathbb{F}$.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\text{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $\mathbf{s} \in \mathbb{F}$.
- ▶ Prover sends back $\mathbf{z} = \mathbf{f}(\mathbf{s})$; together with proof $\text{open}(\mathbf{f}, \mathbf{s})$ that \mathbf{z} is correct.

Polynomial commitment schemes [KZG, 10]

- ▶ Prover send short commitment $\text{cm}(\mathbf{f})$ to polynomial.
- ▶ Later Verifier can choose value $\mathbf{s} \in \mathbb{F}$.
- ▶ Prover sends back $\mathbf{z} = \mathbf{f}(\mathbf{s})$; together with proof $\text{open}(\mathbf{f}, \mathbf{s})$ that \mathbf{z} is correct.

KZG give us PCS with commitments and openings are practically 32 bytes.

Notation: $[\mathbf{x}]_1 = \mathbf{x} \cdot \mathbf{g}$ where \mathbf{g} generator of (first source group of) elliptic curve group with pairing.

Setup: $[1]_1, [\mathbf{x}]_1, \dots, [\mathbf{x}^d]_1$, for random $\mathbf{x} \in \mathbb{F}$.

Setup: $[1]_1, [\mathbf{x}]_1, \dots, [\mathbf{x}^d]_1$, for random $\mathbf{x} \in \mathbb{F}$.

$$\text{cm}(\mathbf{f}) := [\mathbf{f}(\mathbf{x})]_1$$

Setup: $[1]_1, [\mathbf{x}]_1, \dots, [\mathbf{x}^d]_1$, for random $\mathbf{x} \in \mathbb{F}$.

$$\text{cm}(\mathbf{f}) := [\mathbf{f}(\mathbf{x})]_1$$

$$\text{open}(\mathbf{f}, \mathbf{s}) := [\mathbf{h}(\mathbf{x})]_1, \text{ where } \mathbf{h}(\mathbf{X}) := \frac{\mathbf{f}(\mathbf{X}) - \mathbf{f}(\mathbf{s})}{\mathbf{X} - \mathbf{s}}$$

Setup: $[1]_1, [\mathbf{x}]_1, \dots, [\mathbf{x}^d]_1$, for random $\mathbf{x} \in \mathbb{F}$.

$$\text{cm}(\mathbf{f}) := [\mathbf{f}(\mathbf{x})]_1$$

$$\text{open}(\mathbf{f}, \mathbf{s}) := [\mathbf{h}(\mathbf{x})]_1, \text{ where } \mathbf{h}(\mathbf{X}) := \frac{\mathbf{f}(\mathbf{X}) - \mathbf{f}(\mathbf{s})}{\mathbf{X} - \mathbf{s}}$$

$$\text{verify}(\text{cm}, \boldsymbol{\pi}, \mathbf{z}, \mathbf{s}) :$$

$$\mathbf{e}(\text{cm} - [\mathbf{z}]_1, [1]_1) \stackrel{?}{=} \mathbf{e}(\boldsymbol{\pi}, [\mathbf{x} - \mathbf{s}]_1)$$

Opening many polynomials at s

Input: f_0, \dots, f_{d-1} , $z_0 = f_0(s), \dots, z_{d-1} = f_{d-1}(s)$.

Verifier has commitments cm_i to f_i 's wants to
verify correctness of z 's.

Opening many polynomials at s

Input: f_0, \dots, f_{d-1} , $z_0 = f_0(s), \dots, z_{d-1} = f_{d-1}(s)$.

Verifier has commitments cm_i to f_i 's wants to verify correctness of z 's.

Naive solution: Run KZG for each f_i .

Cost: d group elements in proof, d pairings for verifier

Batched opening (Sonic, Marlin, Plonk)

- ▶ Verifier sends random $\gamma \in \mathbb{F}$

Batched opening (Sonic, Marlin, Plonk)

- ▶ Verifier sends random $\gamma \in \mathbb{F}$
- ▶ Prover computes combination
$$\mathbf{f}(\mathbf{X}) := \sum_{i < d} \gamma^i \mathbf{f}_i(\mathbf{X})$$
- ▶ Verifier computes commitment to \mathbf{f} as
$$\text{cm}(\mathbf{f}) := \sum_{i < d} \gamma^i \text{cm}_i$$
- ▶ Prover and verifier use KZG to verify $\mathbf{f}(\mathbf{s}) = \mathbf{z}$
for $\mathbf{z} = \sum_{i < d} \gamma^i \mathbf{z}_i$

cost: $d - 1$ verifier scalar muls to compute $\text{cm}(\mathbf{f})$

Punchline of this work: can get rid of this d dependence when \mathbf{s} is a d 'th power

A useful tool from BDFG/Halo infinite

thm[BDFG]: We can open a polynomial f at points s_0, \dots, s_{d-1} with 2 verifier scalar mults no matter how large d is.

A useful tool from BDFG/Halo infinite

thm[BDFG]: We can open a polynomial f at points s_0, \dots, s_{d-1} with 2 verifier scalar mults no matter how large d is.

*If we can reduce opening many polys at s to opening *one poly* at many points, we can use BDFG to get our desired result*

fflonk central idea: reduce opening many polys to one using the FFT identity

polys f_0, f_1 , $\mathbf{a} = f_0(\mathbf{s})$, $\mathbf{b} = f_1(\mathbf{s})$

fflonk central idea: reduce opening many polys to one using the FFT identity

polys f_0, f_1 , $\mathbf{a} = f_0(\mathbf{s})$, $\mathbf{b} = f_1(\mathbf{s})$

First attempt: Only open the sum -
 $F(\mathbf{X}) := f_0(\mathbf{X}) + f_1(\mathbf{X})$. Prove that
 $F(\mathbf{s}) = \mathbf{c} = \mathbf{a} + \mathbf{b}$.

fflonk central idea: reduce opening many polys to one using the FFT identity

polys f_0, f_1 , $\mathbf{a} = f_0(\mathbf{s})$, $\mathbf{b} = f_1(\mathbf{s})$

First attempt: Only open the sum -
 $F(\mathbf{X}) := f_0(\mathbf{X}) + f_1(\mathbf{X})$. Prove that
 $F(\mathbf{s}) = \mathbf{c} = \mathbf{a} + \mathbf{b}$.

problem: Doesn't constrain \mathbf{a}, \mathbf{b} individually - for any \mathbf{a}' , $(\mathbf{a}', \mathbf{c} - \mathbf{a}')$ will also verify

fflonk central idea: reduce opening many polys to one using the FFT identity

polys f_0, f_1 , $\mathbf{a} = f_0(\mathbf{s})$, $\mathbf{b} = f_1(\mathbf{s})$

First attempt: Only open the sum -
 $F(\mathbf{X}) := f_0(\mathbf{X}) + f_1(\mathbf{X})$. Prove that
 $F(\mathbf{s}) = \mathbf{c} = \mathbf{a} + \mathbf{b}$.

problem: Doesn't constrain \mathbf{a}, \mathbf{b} individually - for any \mathbf{a}' , $(\mathbf{a}', \mathbf{c} - \mathbf{a}')$ will also verify

Solution: Use “FFT equation in reverse direction”:

$$\mathbf{F}(\mathbf{X}) = \mathbf{f}_0(\mathbf{X}^2) + \mathbf{X}\mathbf{f}_1(\mathbf{X}^2)$$

To commit to $\mathbf{f}_0, \mathbf{f}_1$ send $\text{cm}(\mathbf{F})$.

Solution: Use “FFT equation in reverse direction”:

$$\mathbf{F}(\mathbf{X}) = \mathbf{f}_0(\mathbf{X}^2) + \mathbf{X}\mathbf{f}_1(\mathbf{X}^2)$$

To commit to $\mathbf{f}_0, \mathbf{f}_1$ send $\text{cm}(\mathbf{F})$.

Assume $\mathbf{s} = \mathbf{t}^2$.

To open $\mathbf{f}_0(\mathbf{s}), \mathbf{f}_1(\mathbf{s})$ open \mathbf{F} at $\{\mathbf{t}, -\mathbf{t}\}$:

Solution: Use “FFT equation in reverse direction”:

$$\mathbf{F}(\mathbf{X}) = \mathbf{f}_0(\mathbf{X}^2) + \mathbf{X}\mathbf{f}_1(\mathbf{X}^2)$$

To commit to $\mathbf{f}_0, \mathbf{f}_1$ send $\text{cm}(\mathbf{F})$.

Assume $\mathbf{s} = \mathbf{t}^2$.

To open $\mathbf{f}_0(\mathbf{s}), \mathbf{f}_1(\mathbf{s})$ open \mathbf{F} at $\{\mathbf{t}, -\mathbf{t}\}$:

$$\mathbf{b}_0 := \mathbf{F}(\mathbf{t}) = \mathbf{f}_0(\mathbf{s}) + \mathbf{t}\mathbf{f}_1(\mathbf{s}) = \mathbf{a} + \mathbf{t}\mathbf{b}$$

$$\mathbf{b}_1 := \mathbf{F}(-\mathbf{t}) = \mathbf{f}_0(\mathbf{s}) - \mathbf{t}\mathbf{f}_1(\mathbf{s}) = \mathbf{a} - \mathbf{t}\mathbf{b}$$

i.e. $(\mathbf{b}_0, \mathbf{b}_1)$ give two independent constraints on (\mathbf{a}, \mathbf{b}) !

Assume $\mathbf{s} = \mathbf{t}^2$.

To open $\mathbf{f}_0(\mathbf{s}), \mathbf{f}_1(\mathbf{s})$ open \mathbf{F} at $\{\mathbf{t}, -\mathbf{t}\}$:

Assume $\mathbf{s} = \mathbf{t}^2$.

To open $\mathbf{f}_0(\mathbf{s}), \mathbf{f}_1(\mathbf{s})$ open \mathbf{F} at $\{\mathbf{t}, -\mathbf{t}\}$:

$$\mathbf{b}_0 := \mathbf{F}(\mathbf{t}) = \mathbf{f}_0(\mathbf{s}) + \mathbf{t}\mathbf{f}_1(\mathbf{s}) = \mathbf{a} + \mathbf{t}\mathbf{b}$$

$$\mathbf{b}_1 := \mathbf{F}(-\mathbf{t}) = \mathbf{f}_0(\mathbf{s}) - \mathbf{t}\mathbf{f}_1(\mathbf{s}) = \mathbf{a} - \mathbf{t}\mathbf{b}$$

i.e. $(\mathbf{b}_0, \mathbf{b}_1)$ give two independent constraints on (\mathbf{a}, \mathbf{b}) !

- ▶ Similar construction can open \mathbf{d} polys at any $\mathbf{s} = \mathbf{t}^{\mathbf{d}}$
- ▶ *Important:* poly-iop based snarks work fine with a PCS that can only open \mathbf{d} 'th powers.

Cheaply opening a poly at d points

[BDFG]

Given poly f with commitment cm ,

$s_0, \dots, s_{d-1} \in \mathbb{F}$, suppose $z_i = f(s_i)$ for $i < d$.

- Define poly r of degree $< d$ with $r(s_i) = z_i$ for $i < d$.

Cheaply opening a poly at d points

[BDFG]

Given poly f with commitment cm ,

$s_0, \dots, s_{d-1} \in \mathbb{F}$, suppose $z_i = f(s_i)$ for $i < d$.

- ▶ Define poly r of degree $< d$ with $r(s_i) = z_i$ for $i < d$.
- ▶ Define $Z(X) = \prod_{i < d} (X - s_i)$. Then we have $Z|(f - r)$.

Cheaply opening a poly at d points

[BDFG]

Given poly f with commitment cm ,

$s_0, \dots, s_{d-1} \in \mathbb{F}$, suppose $z_i = f(s_i)$ for $i < d$.

- ▶ Define poly r of degree $< d$ with $r(s_i) = z_i$ for $i < d$.
- ▶ Define $Z(X) = \prod_{i < d} (X - s_i)$. Then we have $Z | (f - r)$.
- ▶ Let $h(X) := (f(X) - r(X)) / (Z(X))$. Prover sends $\pi = [h(x)]_1$.

Cheaply opening a poly at d points

[BDFG]

Given poly f with commitment cm ,

$s_0, \dots, s_{d-1} \in \mathbb{F}$, suppose $z_i = f(s_i)$ for $i < d$.

- ▶ Define poly r of degree $< d$ with $r(s_i) = z_i$ for $i < d$.
- ▶ Define $Z(X) = \prod_{i < d} (X - s_i)$. Then we have $Z | (f - r)$.
- ▶ Let $h(X) := (f(X) - r(X)) / (Z(X))$. Prover sends $\pi = [h(x)]_1$.
- ▶ From [KZG]: Verifier can now check

$$e(cm - [r(x)]_1, 1) = e(\pi, [Z(x)]_2)$$

Cheaply opening a poly at d points [BDFG]

Given poly f with commitment cm ,

$s_0, \dots, s_{d-1} \in \mathbb{F}$, suppose $z_i = f(s_i)$ for $i < d$.

- ▶ Define poly r of degree $< d$ with $r(s_i) = z_i$ for $i < d$.
- ▶ Define $Z(X) = \prod_{i < d} (X - s_i)$. Then we have $Z | (f - r)$.
- ▶ Let $h(X) := (f(X) - r(X)) / (Z(X))$. Prover sends $\pi = [h(x)]_1$.
- ▶ From [KZG]: Verifier can now check

$$e(cm - [r(x)]_1, 1) = e(\pi, [Z(x)]_2)$$

This requires d G1 and G2 verifier scalar muls!

In [BDFG] we trade these scalar mults for an extra group element in the proof:

- ▶ Verifier chooses random $\alpha \in \mathbb{F}$ and sends to prover.
- ▶ Define the polynomial

$$\mathbf{L}(\mathbf{X}) := \mathbf{f}(\mathbf{X}) - \mathbf{r}(\alpha) - \mathbf{Z}(\alpha)\mathbf{h}(\mathbf{X})$$

In [BDFG] we trade these scalar mults for an extra group element in the proof:

- ▶ Verifier chooses random $\alpha \in \mathbb{F}$ and sends to prover.
- ▶ Define the polynomial

$$\mathbf{L}(\mathbf{X}) := \mathbf{f}(\mathbf{X}) - \mathbf{r}(\alpha) - \mathbf{Z}(\alpha)\mathbf{h}(\mathbf{X})$$

- ▶ If evals are correct, $\mathbf{L}(\mathbf{X})$ should be zero at α

In [BDFG] we trade these scalar mults for an extra group element in the proof:

- ▶ Verifier chooses random $\alpha \in \mathbb{F}$ and sends to prover.
- ▶ Define the polynomial

$$\mathbf{L}(\mathbf{X}) := \mathbf{f}(\mathbf{X}) - \mathbf{r}(\alpha) - \mathbf{Z}(\alpha)\mathbf{h}(\mathbf{X})$$

- ▶ If evals are correct, $\mathbf{L}(\mathbf{X})$ should be zero at α
- ▶ Verifier can compute $\text{cm}(\mathbf{L})$ with only two scalar mults:

$$\text{cm}(\mathbf{L}) = \text{cm} - [\mathbf{r}(\alpha)]_1 - \mathbf{Z}(\alpha)\pi$$

- ▶ Prover and verifier can now use KZG to check $\mathbf{L}(\alpha) = 0$.