

Developing Circadian Rhythmicity in Infants

Scott A. Rivkees, MD

ABSTRACT. Circadian rhythms are endogenously generated rhythms with a period length of approximately 24 hours. Evidence gathered during the past decade indicates that the circadian timing system develops prenatally and the suprachiasmatic nuclei, the site of a circadian clock, is present by midgestation in primates. Recent evidence also shows that the circadian system of primate infants is responsive to light at very premature stages and that low-intensity lighting can regulate the developing clock. After birth, there is progressive maturation of the circadian system outputs, with pronounced rhythms in sleep-wake and hormone secretion generally developing after 2 months of age. Showing the importance of photic regulation of circadian phase in infants, exposure of premature infants to low-intensity cycled lighting results in the early establishment of rest-activity patterns that are in phase with the 24-hour light-dark cycle. With the continued elucidation of circadian system development and influences on human physiology and illness, it is anticipated that consideration of circadian biology will become an increasingly important component of neonatal care. *Pediatrics* 2003;112:373–381; *circadian rhythms, infant, human, baboon, suprachiasmatic nuclei*.

ABBREVIATIONS. SCN, suprachiasmatic nuclei; RHT, retinohypothalamic tract.

THE CIRCADIAN TIMING SYSTEM

Circadian rhythms are endogenously driven rhythms with a period length of approximately 24 hours.¹ Notable examples of circadian rhythms include the sleep-wake cycle and daily rhythms in hormone production. Circadian rhythms are also involved in the pathogenesis of illnesses, such as reactive airway disease and myocardial infarction.^{2,3}

The system responsible for the generation and regulation of circadian rhythms is the circadian timing system. This neural system consists of a biological clock, input pathways, and output pathways¹ (Fig 1). The paired suprachiasmatic nuclei (SCN) in the anterior hypothalamus are the site of a biological clock. The SCN are located above the optic chiasm at the base of the third ventricle⁴ (Fig 2). The SCN exhibit

endogenous rhythmicity and have a period of oscillation close to 24 hours.

Lesion studies in rodents provided the initial evidence that the SCN are the site of a circadian pacemaker.⁴ In vivo and in vitro studies have since shown day-night rhythms in electrical activity, metabolic activity, and gene expression. Transplantation of fetal SCN cells into SCN-lesioned animals restores rhythmicity to the recipient, further supporting that the SCN contain a biological clock.⁴ Circadian oscillations have been seen in individual rodent SCN cells, and expressed rhythmicity reflects the collective oscillations of many SCN cells.^{5,6}

Evidence gathered during the past several years has yielded new insights into the molecular mechanisms resulting in clock oscillations.^{1,7,8} Twenty-four-hour rhythmicity seems to result from rhythmic expression of transcriptional activating factors, including *per*, *clock*, and *tim*, which form long feedback loops in SCN cells. Cryptochromes (*cry*) also play a critical role in regulating clock function by interacting with *per*.^{1,7,8}

Because SCN oscillations are not exactly 24 hours, it is necessary to reset the circadian pacemaker each day to prevent endogenous clock oscillations from drifting (or free-running) out of phase with the external light-dark cycle. Input pathways relay photic information from the retina to the SCN to synchronize (or entrain) the oscillations of the clock to the 24-hour light-dark cycle.⁹ A direct pathway from the retina to the SCN, the retinohypothalamic tract (RHT; Figs 1 and 3), has been shown to be both necessary and sufficient for photic entrainment.⁹ The raphe nucleus also influences SCN function via serotonergic projections.⁹

Glutamate is the neurotransmitter that mediates RHT action, and the phase shifting actions of glutamate involve the nitric oxide system.¹⁰ Another pathway from the intergeniculate leaflet to the SCN modulates clock entrainment and involves the release of neuropeptide Y.⁹

Recent evidence suggests that the products of intermediate-early genes, which include the DNA-binding proteins c-Fos and Jun-B, may mediate entrainment.¹¹ Light exposure at times that reset (phase shift) the clock induces increases in SCN metabolic activity and robust expression of c-Fos and Jun-B in the SCN.¹¹

Output pathways are responsible for the overt expression of circadian rhythms. Several discrete neural pathways projecting from the SCN to several

From the Yale Child Health Research Center, Department of Pediatrics, Yale University School of Medicine, New Haven, Connecticut.

Received for publication Feb 12, 2003; accepted Feb 12, 2003.

Reprint requests to (S.A.R.) Yale Pediatrics, Yale Child Health Research Center, Box 208081, 333 Cedar St, New Haven, CT 06510. E-mail: scott.rivkees@yale.edu

PEDIATRICS (ISSN 0031 4005). Copyright © 2003 by the American Academy of Pediatrics.

The Circadian Timing System




Fig 1. Schematic representation of the circadian timing system.

hypothalamic and nonhypothalamic sites have been defined.^{11–13} Via these pathways, the circadian system acts to influence neural physiology broadly. Output pathways of the circadian system also regulate the rhythmic production of several hormones, including melatonin and cortisol.^{2,3,11–13}

THE PRIMATE CIRCADIAN TIMING SYSTEM

Several lines of evidence support that the paired SCN are the site of a biological clock in primates. Similar to rodents, the primate SCN are located above the optic chiasm at the base of the third ventricle.¹⁴ In contrast to rodents, human SCN cells are not densely clustered, making the nuclei less visually apparent.^{14–16} However, with the use of probes for melatonin receptors and SCN peptides, the human SCN can be identified.^{14–16} With the use of 2-deoxyglucose, day-night oscillations in SCN metabolic activity have been detected in squirrel monkeys and baboons.^{17–19}

Lesion studies performed in the early 1980s suggested the presence of a circadian pacemaker outside the SCN in monkeys.²⁰ However, analysis of these reports revealed that either the completeness of the lesions was not verified or monkeys were not studied

in constant conditions.²¹ Reexamination of this issue challenges the existence of primate circadian pacemakers outside the SCN. Squirrel monkeys with total SCN lesions show a complete absence of circadian rhythmicity when animals are monitored in constant conditions.²¹ Supporting that the SCN are the site of a circadian pacemaker in humans, tumors and congenital lesions in the SCN region result in the loss of temperature rhythms and organized sleep-wake patterns.^{22,23}

The RHT has been anatomically characterized in prosimian (lemurs and shrews) and simian (squirrel monkeys, rhesus macaques, baboons, chimpanzees, and apes) species.¹⁴ This tract also has been identified in studies of postmortem human specimens using techniques that label degenerating retinal axons.^{24,25} Although it was suggested that cutaneous light exposure can influence circadian function,²⁶ there is little support for the notion that there is extraretinal photoreception in mammals.^{27–29} Furthermore, other investigators have failed to reproduce phase-shifting effects of cutaneous light exposure.¹⁷

Outputs of the primate circadian system have been widely characterized in human clinical studies. Many day-night rhythms have been documented.^{2,3,28} Several of these rhythms have been shown to persist in constant conditions, indicating that they are true endogenously generated circadian rhythms. Notable examples of circadian rhythms include the sleep-wake cycle, daily rhythms in body temperature, and day-night rhythms in cortisol and melatonin production.^{2,3,28} Day-night differences in gonadotropin, testosterone, growth hormone, and thyrotropin secretion are also present.^{30,31}





Fig 2. The human SCN. Reproduced with permission from Moore.¹⁴

Fig 3. Innervation of the SCN by the RHT in a newborn baboon infant. A, Low-power image showing labeling of retinal fibers in the optic chiasm by horseradish peroxidase. B, Adjacent tissue section showing the location of the SCN. C, High-power image showing projections of the RHT into the right SCN. D, Autoradiographic image generated from [^{14}C]2-deoxyglucose uptake studies showing that light exposure at night induces increases in SCN metabolic activity. Areas of increased uptake are dark. Arrows identify the SCN. Scale bar = 5 mm. Reproduced with permission from Rivkees et al.¹⁹



DEVELOPMENT OF THE PRIMATE SCN

Although rodent studies have led to our understanding of developmental circadian physiology,^{30–33} notable differences between rodents and primates have prevented the extension of rodent data to clinical care. In general, rodents are more immature at birth than humans. Differences in the sensitivity to light and other aspects of circadian physiology between humans and rodents also have been observed. However, on the basis of evidence gathered during the past decade, it seems that the circadian clock in the SCN forms and begins oscillating in utero in primates.


In squirrel monkeys, SCN neurogenesis occurs

early in gestation during days 27 to 48.³⁴ Because monkey and human embryonic development is very similar during the first 100 days of gestation,³⁵ it therefore is likely that the human SCN neurons form early in gestation.

It is not known when the primate SCN are first apparent morphologically, yet with the use of [^{125}I]melatonin and [^{125}I]SKF38393 to label the nuclei, the human SCN have been detected at gestation week 18^{36,37} (Fig 4).

Functional studies suggest that the primate SCN oscillate prenatally. Studies of squirrel monkeys reveal day-night differences in SCN metabolic activity at the end of gestation.³⁸ It is not known whether

Fig 4. A, Localization of [^{125}I]melatonin binding to the SCN of an 18-week gestation human fetus. Specific labeling is shown in black. B, The stained section used to generate the autoradiograph in A. Reproduced with permission from Reppert et al.³⁶ C, Localization of [^{125}I]SKF38393 binding to D1 dopamine receptors in the SCN of a 20-week postconceptional human infant. Specific labeling is shown in black. D, Nonspecific labeling. Reproduced with permission from Rivkees et al.³⁷ OC, optic chiasm; ST, striatum. Arrows identify the SCN.



SCN oscillations are present at earlier ages. The physiologic processes influenced by the fetal clock have yet to be elucidated in primates.

Similar to rodents, the timing of the onset of labor and birth in humans is influenced by the circadian cycle with peak incidences between midnight and the early morning.³⁹ However, we do not know whether the fetal clock plays a role in the circadian gating of birth in humans.

Immunocytochemistry studies show that SCN maturation continues after birth.⁴⁰ The SCN contain distinct populations of neurons that express arginine vasopressin or vasoactive intestinal polypeptide.⁴⁰ In term infants, the number of vasopressinergic neurons is 20% of the number present in adults.⁴⁰ It is not until 1 year of age that infants and adults have comparable vasopressin neuron numbers.⁴⁰ The number of vasoactive intestinal polypeptide-containing SCN cells also increase after birth.⁴⁰

DEVELOPMENT OF PRIMATE PHOTIC ENTRAINMENT

A critical issue in knowing whether environmental cycles need to be considered in the care of infants is knowing when the primate circadian system becomes functionally responsive to light. The RHT has been identified in a 36-week-gestation human newborn.⁴¹ However, because of human study limitations, it has not been possible to determine whether the circadian clock of human infants is functionally responsive to light at birth.

Noninvasive methods used to examine regional changes in brain activity, such as function magnetic resonance imaging or positron emission tomography, hold promise in being able to examine SCN function directly. In human adults, we have been able to observe acute increases in SCN metabolic activity after light exposure at night using ¹⁸F-2-deoxyglucose in positron emission tomography studies.⁴² However, because of the small size of the SCN, consistent visualization of SCN activity is difficult to achieve and these methods have not been applied to infants.

Because of human study limitations, we have studied baboons, which are excellent models for human infants, to provide insights into the developing human clock. By monitoring changes in SCN metabolic

activity and gene expression (Fig 3), light responsiveness can be demonstrated at birth in term baboon infants.¹⁹ The presence of the RHT can also be demonstrated (Fig 3).¹⁹

By monitoring the effects of different lighting conditions on newborn baboon activity patterns, we have been able to show that newborn baboons are entrained by low-intensity (200 lux) lighting.¹⁹ These findings are similar to those seen in human adults showing that circadian phase can be regulated by low-intensity (ca. 180 lux) lighting.^{43,44} Thus, it is likely that low-intensity lighting, similar to that found indoors, can regulate the developing primate clock.


To determine when photic responsiveness first occurs in primates, we have also studied premature baboon infants.⁴⁵ To our surprise, we find that the SCN are functionally innervated by the retina at stages equivalent to human infants at 25 weeks postconception⁴⁵ (Fig 5). The primate circadian system therefore is sensitive to light in very premature infants when postnatal survival with intensive support becomes possible.

DEVELOPMENT OF EXPRESSED RHYTHMICITY IN PRIMATES

The development of expressed rhythmicity has received attention in both human and nonhuman primates. During pregnancy, day-night rhythms are observed for a variety of hormones (estrogen and progesterone) and physiologic parameters (uterine contractility) in mothers.^{46,47} In human fetuses, day-night rhythms in heart rate, respiratory rate, and adrenal steroidogenesis have been detected.^{46,47} However, these rhythms seem to be driven by the mother.

When term human infants are examined, day-night rhythms are difficult to detect in the neonatal period.^{42,48–50} Consolidated periods of activity and rest generally are not observed until after the first or second month of life. Activity plots of human newborns reveal that sleep is generally distributed over the 24-hour day during the first few weeks of life (Fig 6). At 6 weeks of age, infants are awake more during the daytime than at night. By 12 weeks of age, daytime sleep duration decreases further and much more sleep occurs at night. Importantly, although

Fig 5. Autoradiograph images of preterm baboon brain sections showing SCN 2-deoxyglucose uptake after light exposure at night. Animals shown were 160 or 125 weeks of postconceptional age and were either blindfolded or directly exposed to light. The images are obtained from mid-SCN levels. Arrows identify the SCN image. Reproduced with permission from Hao and Rivkees.⁴⁵ Human equivalent ages are provided in weeks.






Fig 6. Rest-activity patterns of a newborn human (A) and a baboon (B) infant (right). Double-plotted actograms are shown. In A, dark bars represent sleep. In B, dark bars represent activity. Please note that the circadian phase of the baboon infant was shifted by exposure to a 200-lux reversed light-dark cycle at 30 days of age and much less so by 50 lux of exposure at 55 days of age. Reproduced with permission from Rivkees et al.¹⁹

consolidated periods of rest and activity are not apparent until >1 to 2 months after birth, day-night differences in activity can be detected as early as 1 week of age in some infants.

At the age when day-night differences in infant activity become clearly apparent, day-night rhythms in hormone production are observed. Day-night rhythms in melatonin production can be detected at 12 weeks of age.^{51,52} Circadian variation in cortisol levels appears between 3 and 6 months of age.^{53–55} With advancing age, circadian rhythms have been detected for a variety of other hormones and circulating factors.⁵⁶

Because infant care influences activity patterns, it is possible that patterns of developing circadian rhythmicity in human infants reflects influences of caregivers rather than endogenous rhythmicity. Thus, to characterize the development of expressed rhythmicity in primates, we have examined the development of expressed rhythmicity in newborn baboons raised in constant conditions (continuous dim lighting, evenly spaced care).¹⁹ Similar to human infants, baboon infants do not manifest clear day-night differences in activity patterns in the early neonatal period (Fig 6), yet at 1 month of age, day-night differences in activity patterns are observed. Developing primate rhythmicity thus reflects maturation from a state of relative arrhythmicity to rhythmicity during the first few months of life.

As in rodents, it seems that infant circadian phase is synchronous with that of the mother in baboon and human infants. However, in some humans and baboons,^{19,57} infant phase may be out of synchrony with that of the mother at birth. Thus, whereas there is mother-infant synchrony of circadian phase in most primates, it may not be universal.

RHYTHMICITY IN PRETERM INFANTS

The large number of premature infants who are hospitalized for extended periods has greatly facilitated studies of rhythmicity in preterm infants. During the past decade, studies of patterns of infant activity, heart rate, temperature, and sleep state surprisingly have not flourished.⁵⁸ Several of these studies have revealed the presence of ultradian rhythms (rhythms with period lengths of much less than 24 hours). Endogenously driven circadian rhythms, however, are not clearly apparent.

When temperature and heart rate are studied beginning at a postconceptional age of 24 to 29 weeks, circadian rhythmicity is generally not apparent even at 17 weeks after birth.⁵⁹ Studies of preterm infants at 32 weeks' postconceptional age have failed to detect day-night differences in sleep patterns, whereas some differences are noted in term infants.⁶⁰ Analysis of temperature, heart rate, and activity patterns at 35 weeks' postconceptional age has revealed ultradian rhythms but no clear-cut circadian rhythms.^{60–62} Because feeding and physical contact influence infant temperature, heart rates, and activity patterns, it is likely that infant care schedules drive the ultradian rhythms seen in preterm infants. These interventions may also mask the detection of circadian rhythms.

YALE NEONATAL ENTRAINMENT STUDY

After the discovery that the primate circadian clock is responsive to light in very premature infants, we next assessed the effects of photic entrainment on premature infants.⁶³ In these studies, the development of rest-activity patterns was examined in human preterm infants who were exposed to continuous dim lighting or low-intensity cycled lighting before discharge from hospital to home.

In general, day-night differences in rest and activity are not apparent in hospitalized control infants (Fig 7), whereas day-night differences in rest and activity are seen in experimental infants (Fig 8). During the first 10 days at home, distinct day-night differences in activity are not seen in control infants, but experimental infants are more active during the day than at night. It was not until 21 to 30 days after discharge that day-night activity ratios in control infants matched those seen in experimental infants shortly after discharge, yet even at this age, experimental infants are considerably more active during the day as compared with control infants. Despite the differences in rest-activity patterns among groups, no differences in weight gain or change in head circumference are seen.

These observations show that exposure to low-intensity cycled lighting for 10 days before discharge induces distinct patterns of rest/activity in preterm infants that are in synchrony with the light-dark cycle that they will encounter at home. These effects are even more pronounced as soon as the child is discharged to home. In contrast, the appearance of rest-activity patterns in synchrony with the solar light-dark cycle is delayed in infants who have been exposed to continuous dim lighting in the hospital.

Potential influences of cycled lighting on premature infants have been the subject of a few previous studies. In the Stanford Cycled Lighting Study, differences in circadian rhythms in temperature were not detected among infants who were exposed either to continuous dim lighting or to cycled lighting before discharge.^{61,62,64} These infants were studied 1 and 3 months after discharge. Because we observed that infants in both groups manifest similar circadian

phase by 30 days of age, treatment effects on the rhythm of core body temperature may no longer be distinct after 1 month of age.

Other investigators have suggested that exposing infants to light-dark cycles improves infant weight gain. Mann et al⁶⁵ found that exposure to light-dark cycles before discharge resulted in better weight gain and more sleep during the 24-hour day than did chaotic lighting patterns. These effects were seen 6 weeks after discharge and not sooner.⁶⁵ Because of this lag period, it has been suggested that the observed effects were not a direct result of cycled lighting exposure on the infant.⁶⁴ More recently, it has been suggested that exposing infants to light-dark cycles improves the in-hospital growth of infants if exposure occurs before 36 weeks of age,⁶⁶ yet the infants in the near-darkness group in this study seemed more ill than the other groups. Considering that it is difficult to detect circadian activity in premature infants,^{42,67} the potential mechanisms by which lighting could directly influence the growth of premature infants is not clear. By studying infants that were closely matched at enrollment, we failed to observe influences of lighting on growth either in hospital or at home.

Previous studies have suggested that day-night rhythmicity is not apparent in prematurely born infants until nearly 1 month after term-birth age equivalence is reached (>42 weeks' postmenstrual age).^{61,62,64,68} These conclusions have been based on 24- to 48-hour assessments of rectal temperature and/or sleep patterns. However, using actigraphy to monitor rest-activity patterns continuously, we found that circadian phase can be detected in infants who were exposed to cycled lighting as early as a




Fig 7. Actograms of rest-activity in a representative infant exposed to constant dim light. Dark bars represent activity. Arrow depicts the date of discharge. The number of days relative to discharge is shown on the left. The time of day is shown on top. Note that day-night differences in rest-activity patterns in synchrony with the light-dark cycle are not apparent until approximately 20 days after discharge.

Cycled Lighting




Fig 8. Actograms of rest-activity in a representative infant exposed to cycled lighting. Note the regular patterns of rest and activity in the infant exposed to cycled lighting before and after discharge.

postmenstrual age of 34 weeks. In our previous studies of nonhuman primate infants reared in constant conditions, we also found that day-night differences in rest and activity were apparent shortly after term birth.¹⁹ Most important, we found that day-night differences in activity could be detected several weeks before it was possible to detect circadian rhythms in core temperature using internal telemetry devices.¹⁹ Thus, analysis of rest-activity patterns may provide the earliest index of developing circadian rhythmicity in infants.

Although continuous dim lighting is the current practice in most nurseries in the United States, the scientific basis for this practice is not clear.⁶⁹ It has been suggested that ambient lighting may contribute to eye disease in premature infants,⁷⁰ yet rigorous clinical studies have failed to show adverse effects of low-intensity lighting on premature infants.^{71–73} Investigators who propose a Neonatal Individualized Developmental Care Assessment Program have suggested that because the womb is dark, infants should be reared in the dark.⁷⁴ This approach overlooks that prenatally the infant is exposed to maternal time-of-day cues that synchronize the fetal clock with the external light-dark cycle.³² Rearing premature infants in the dark thus deprives infants of the time-of-day information that they would have received during full gestation. Data also show that the Neonatal Individualized Developmental Care Assessment Program approach does not improve developmental outcome or sleep of premature infants.⁷⁵

CONCLUSIONS

Increasing evidence indicates that the circadian timing system is a fundamental homeostatic system

that potently influences human behavior and physiology throughout development (Fig 9). After birth, there is progressive maturation of the circadian system with day-night rhythms in activity and hormone secretion developing between 1 and 3 months of age. Recent evidence shows that the circadian system of primate infants is responsive to light at very premature stages and that low-intensity lighting can regulate the developing clock. With the continued elucidation of circadian system development and influences on human physiology and illness, it is anticipated that consideration of circadian biology will become an increasingly important component of neonatal care.




Fig 9. Schematic representation of primate circadian system development based on studies of nonhuman primates. Estimated human ages are given.

ACKNOWLEDGMENT

This work was supported by National Institutes of Health grant NS32624.

REFERENCES

- Panda S, Hogenesch JB, Kay SA. Circadian rhythms from flies to human. *Nature*. 2002;417:329–335
- Moore-Ede MC, Czeisler CA, Richardson GS. Circadian timekeeping in health and disease. Part 1. Basic properties of circadian pacemakers. *N Engl J Med*. 1983;309:469–476
- Moore-Ede MC, Czeisler CA, Richardson GS. Circadian timekeeping in health and disease. Part 2. Clinical implications of circadian rhythmicity. *N Engl J Med*. 1983;309:530–536
- Weaver DR. The suprachiasmatic nucleus: a 25-year retrospective. *J Biol Rhythms*. 1998;13:100–112
- Welsh DK, Logothetis DE, Meister M, Reppert SM. Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron*. 1995;14:697–706
- Liu C, Weaver DR, Strogatz SH, Reppert SM. Cellular construction of a circadian clock: period determination in the suprachiasmatic nuclei. *Cell*. 1997;91:855–860
- Albrecht U. Invited review: regulation of mammalian circadian clock genes. *J Appl Physiol*. 2002;92:1348–1355
- Reppert SM. Cellular and molecular basis of circadian timing in mammals. *Semin Perinatol*. 2000;24:243–246
- Morin LP. The circadian visual system. *Brain Res Brain Res Rev*. 1994;19:102–127
- Ding JM, Buchanan GF, Tischkau SA, et al. A neuronal ryanodine receptor mediates light-induced phase delays of the circadian clock. *Nature*. 1998;394:381–384
- Kornhauser JM, Mayo KE, Takahashi JS. Light, immediate-early genes, and circadian rhythms. *Behav Genet*. 1996;26:221–240
- Watts AG, Swanson LW, Sanchez-Watts G. Efferent projections of the suprachiasmatic nucleus: I. Studies using anterograde transport of Phaseolus vulgaris leucoagglutinin in the rat. *J Comp Neurol*. 1987;258:204–229
- Watts AG, Swanson LW. Efferent projections of the suprachiasmatic nucleus: II. Studies using retrograde transport of fluorescent dyes and simultaneous peptide immunohistochemistry in the rat. *J Comp Neurol*. 1987;258:230–252
- Moore RY. Organization of the primate circadian system. *J Biol Rhythms*. 1993;(8 suppl):S3–S9
- Lydic R, Schoene WC, Czeisler CA, Moore-Ede MC. Suprachiasmatic region of the human hypothalamus: homolog to the primate circadian pacemaker? *Sleep*. 1980;2:355–361
- Lydic R, Albers HE, Tepper B, Moore-Ede MC. Three-dimensional structure of the mammalian suprachiasmatic nuclei: a comparative study of five species. *J Comp Neurol*. 1982;204:225–237
- Lushington K, Galka R, Sassi LN, Kennaway DJ, Dawson D. Extraocular light exposure does not phase shift saliva melatonin rhythms in sleeping subjects. *J Biol Rhythms*. 2002;17:377–386
- Schwartz WJ, Reppert SM, Eagan SM, Moore-Ede MC. In vivo metabolic activity of the suprachiasmatic nuclei: a comparative study. *Brain Res*. 1983;274:184–187
- Rivkees SA, Hofman PL, Fortman J. Newborn primate infants are entrained by low intensity lighting. *Proc Natl Acad Sci U S A*. 1997;94:292–297
- Reppert SM, Perlow MJ, Ungerleider LG, et al. Effects of damage to the suprachiasmatic area of the anterior hypothalamus on the daily melatonin and cortisol rhythms in the rhesus monkey. *J Neurosci*. 1981;1:1414–1425
- Edgar DM, Dement WC, Fuller CA. Effect of SCN lesions on sleep in squirrel monkeys: evidence for opponent processes in sleep-wake regulation. *J Neurosci*. 1993;13:1065–1079
- Schwartz WJ, Busis NA, Hedley-Whyte ET. A discrete lesion of ventral hypothalamus and optic chiasm that disturbed the daily temperature rhythm. *J Neurol*. 1986;233:1–4
- Rivkees S. Arrhythmicity in a child with septo-optic dysplasia and establishment of sleep-wake cyclicity with melatonin. *J Pediatr*. 2001;139:463–465
- Friedman DI, Johnson JK, Chorsky RL, Stopa EG. Labeling of human retinohypothalamic tract with the carbocyanine dye, DiI. *Brain Res*. 1991;560(1–2):297–302
- Sadun AA, Schaechter JD, Smith LE. A retinohypothalamic pathway in man: light mediation of circadian rhythms. *Brain Res*. 1984;302:371–377
- Campbell SS, Murphy PJ. Extraocular circadian phototransduction in humans. *Science*. 1998;279:396–399
- Hebert M, Martin SK, Eastman CI. Nocturnal melatonin secretion is not suppressed by light exposure behind the knee in humans. *Neurosci Lett*. 1999;274:127–130
- Foster RG. Shedding light on the biological clock. *Neuron*. 1998;20:829–832
- Jean-Louis G, Kripke DF, Cole RJ, Elliot JA. No melatonin suppression by illumination of popliteal fossae or eyelids. *J Biol Rhythms*. 2000;15:265–269
- Weitzman ED, Czeisler CA, Zimmerman JC, Moore-Ede MC. Biological rhythms in man: relationship of sleep-wake, cortisol, growth hormone, and temperature during temporal isolation. *Adv Biochem Psychopharmacol*. 1981;28:475–499
- Czeisler CA, Klerman EB. Circadian and sleep-dependent regulation of hormone release in humans. *Recent Prog Horm Res*. 1999;54:97–130
- Reppert SM, Weaver DR, Rivkees SA. Maternal communication of circadian phase to the developing mammal. *Psychoneuroendocrinology*. 1988;13(1–2):63–78
- Reppert SM. Interaction between the circadian clocks of mother and fetus. *Ciba Found Symp*. 1995;183:198–207; discussion 207–211
- van Eerdenburg FJ, Rakic P. Early neurogenesis in the anterior hypothalamus of the rhesus monkey. *Brain Res Dev Brain Res*. 1994;79:290–296
- Nowell-Morris L, Fahrenbruch CE. *Practical and Evolutionary Considerations for the Use of Nonhuman Primate Model in Prenatal Research*. New York, NY: Alan R. Liss; 1985
- Reppert SM, Weaver DR, Rivkees SA, Stopa EG. Putative melatonin receptors in a human biological clock. *Science*. 1988;242:78–81
- Rivkees SA, Lachowicz JE. Functional D1 and D5 dopamine receptors are expressed in the suprachiasmatic, supraoptic, and paraventricular nuclei of primates. *Synapse*. 1997;26:1–10
- Reppert SM, Schwartz WJ. Functional activity of the suprachiasmatic nuclei in the fetal primate. *Neurosci Lett*. 1984;46:145–149
- Jolly A. Hour of birth in primates and man. *Folia Primatol (Basel)*. 1972;18:108–121
- Swaab DF. Development of the human hypothalamus. *Neurochem Res*. 1995;20:509–519
- Glutzbach SF, Sollars P, Pagano M. Development of the human retino-hypothalamic tract. *Soc Neurosci*. 1992;18:857
- Rivkees SA, Hao H. Developing circadian rhythmicity. *Semin Perinatol*. 2000;24:232–242
- Boivin DB, Duffy JF, Kronauer RE, Czeisler CA. Dose-response relationships for resetting of human circadian clock by light. *Nature*. 1996;379:540–542
- Shanahan TL, Czeisler CA. Physiological effects of light on the human circadian pacemaker. *Semin Perinatol*. 2000;24:299–320
- Hao H, Rivkees SA. The biological clock of very premature primate infants is responsive to light. *Proc Natl Acad Sci U S A*. 1999;96:2426–2429
- Seron-Ferre M, Ducsay CA, Valenzuela GJ. Circadian rhythms during pregnancy. *Endocr Rev*. 1993;14:594–609
- Seron-Ferre M, Torres-Farfan C, Forcelledo ML, Valenzuela GJ. The development of circadian rhythms in the fetus and neonate. *Semin Perinatol*. 2001;25:363–370
- Parmelee AH Jr. Sleep cycles in infants. *Dev Med Child Neurol*. 1969;11:794–795
- Meier-Koll A, Hall U, Hellwig U, Kott G, Meier-Koll V. A biological oscillator system and the development of sleep-waking behavior during early infancy. *Chronobiologia*. 1978;5:425–440
- Kleitman J, Engelman K. Sleep characteristics of infants. *J Appl Physiol*. 1953;6:127–134
- Kennaway DJ, Stamp GE, Goble FC. Development of melatonin production in infants and the impact of prematurity. *J Clin Endocrinol Metab*. 1992;75:367–369
- Kennaway DJ, Goble FC, Stamp GE. Factors influencing the development of melatonin rhythmicity in humans. *J Clin Endocrinol Metab*. 1996;81:1525–1532
- Beitins IZ, Kowarski A, Migeon CJ, Graham GG. Adrenal function in normal infants and in marasmus and kwashiorkor. Cortisol secretion, diurnal variation of plasma cortisol, and urinary excretion of 17-hydroxycorticoids, free corticoids, and cortisol. *J Pediatr*. 1975;86:302–308
- Onishi S, Miyazawa G, Nishimura Y, et al. Postnatal development of circadian rhythm in serum cortisol levels in children. *Pediatrics*. 1983;72:399–404
- Price DA, Close GC, Fielding BA. Age of appearance of circadian rhythm in salivary cortisol values in infancy. *Arch Dis Child*. 1983;58:454–456
- Hellbrügge T, Lange JE, Rutenfranz J. Circadian periodicity of physio-

- logical functions in different stages of infancy and childhood. *Ann N Y Acad Sci.* 1964;117:361–373
57. Parmelee AH Jr, Wenner WH, Akiyama Y, Schultz M, Stern E. Sleep states in premature infants. *Dev Med Child Neurol.* 1967;9:70–77
 58. Rivkees SA. Developing circadian rhythmicity. Basic and clinical aspects. *Pediatr Clin North Am.* 1997;44:467–487
 59. D'Souza SW, Tenreiro S, Minors D, Chiswick ML, Sims DG, Waterhouse J. Skin temperature and heart rate rhythms in infants of extreme prematurity. *Arch Dis Child.* 1992;67(Spec No):784–788
 60. Anders TF, Keener MA, Kraemer H. Sleep-wake state organization, neonatal assessment and development in premature infants during the first year of life. II. *Sleep.* 1985;8:193–206
 61. Glotzbach SF, Edgar DM, Boeddiker M, Ariagno RL. Biological rhythmicity in normal infants during the first 3 months of life. *Pediatrics.* 1994;94:482–488
 62. Glotzbach SF, Edgar DM, Ariagno RL. Biological rhythmicity in preterm infants prior to discharge from neonatal intensive care. *Pediatrics.* 1995;95:231–237
 63. Rivkees SA, Gross I, Mayes L. Influence of cycled lighting on the development of rest-activity patterns in premature infants [abstract]. *Pediatr Res.* 2002;51:375
 64. Mirmiran M, Ariagno RL. Influence of light in the NICU on the development of circadian rhythms in preterm infants. *Semin Perinatol.* 2000;24:247–257
 65. Mann NP, Haddow R, Stokes L, Goodley S, Rutter N. Effect of night and day on preterm infants in a newborn nursery: randomised trial. *Br Med J (Clin Res Ed).* 1986;293:1265–1267
 66. Brandon DH, Holditch-Davis D, Belyea M. Preterm infants born at less than 31 weeks' gestation have improved growth in cycled light compared with continuous near darkness. *J Pediatr.* 2002;140:192–199
 67. Mirmiran M, Lunshof S. Perinatal development of human circadian rhythms. *Prog Brain Res.* 1996;111:217–226
 68. McGraw K, Hoffmann R, Harker C, Herman JH. The development of circadian rhythms in a human infant. *Sleep.* 1999;22:303–310
 69. Ariagno RL, Mirmiran M. Shedding light on the very low birth weight infant. *J Pediatr.* 2001;139:476–477
 70. Glass P, Avery GB, Subramanian KN, Keys MP, Sostek AM, Friendly DS. Effect of bright light in the hospital nursery on the incidence of retinopathy of prematurity. *N Engl J Med.* 1985;313:401–404
 71. Fielder AR, Moseley MJ. Environmental light and the preterm infant. *Semin Perinatol.* 2000;24:291–298
 72. Reynolds JD, Hardy RJ, Kennedy KA, Spencer R, van Heuven WA, Fielder AR. Lack of efficacy of light reduction in preventing retinopathy of prematurity. Light Reduction in Retinopathy of Prematurity (LIGHT-ROP) Cooperative Group. *N Engl J Med.* 1998;338:1572–1576
 73. Kennedy KA, Fielder AR, Hardy RJ, Tung B, Gordon DC, Reynolds JD. Reduced lighting does not improve medical outcomes in very low birth weight infants. *J Pediatr.* 2001;139:527–531
 74. Als H, Lawhon G, Duffy FH, McAnulty GB, Gibes-Grossman R, Blickman JG. Individualized developmental care for the very low-birth-weight preterm infant. Medical and neurofunctional effects. *JAMA.* 1994;272:853–858
 75. Ariagno RL, Thoman EB, Boeddiker MA, et al. Developmental care does not alter sleep and development of premature infants. *Pediatrics.* 1997;100(6). Available at: <http://www.pediatrics.org/cgi/content/full/100/6/e9>

SURPRISE! DIRECT-TO-CONSUMER (DTC) ADVERTISING WORKS!

“... Direct-to-consumer (DTC) advertising is perhaps the most visible facet of marketing efforts by drug companies. It is just one portion of an explosion in pharmaceutical promotional advertising. In 2000, the industry spent \$15.7 billion on promotion, with \$2.5 billion spent of DTC advertising, \$4 billion on office-based promotions, and \$8 billion on free samples. In 2001, overall promotional spending jumped to \$19.1 billion, with \$2.7 billion on DTC advertising, \$4.8 billion on office-based promotions, and \$10.5 billion on free samples, according to IMS, a company based in Plymouth Meeting, Pennsylvania, offering economic information and analysis to the pharmaceutical industry... The Food and Drug Administration's numbers imply that DTC advertising is having the effect desired by drug manufacturers. In the survey, 211 of 359 physicians (59%) said a patient had asked them to prescribe a specific brand name drug. And of those 211 physicians, 57% prescribed the brand name drug requested.”

Mitka M. Survey suggesting that prescription drug ads help public is met with skepticism. *JAMA.* 2003;289:827–828

Noted by JFL, MD

Developing Circadian Rhythmicity in Infants

Scott A. Rivkees

Pediatrics 2003;112;373

DOI: 10.1542/peds.112.2.373

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/112/2/373>

References

This article cites 73 articles, 11 of which you can access for free at:
<http://pediatrics.aappublications.org/content/112/2/373#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Endocrinology
http://www.aappublications.org/cgi/collection/endocrinology_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Developing Circadian Rhythmicity in Infants

Scott A. Rivkees

Pediatrics 2003;112;373

DOI: 10.1542/peds.112.2.373

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/112/2/373>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2003 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

