Neap

VCE Specialist Mathematics Units 3&4

Formula Sheet

2024 Trial Examination

You may keep this Formula Sheet.

Neap® Education (Neap) Trial Exams are licensed to be photocopied or placed on the school intranet and used only within the confines of the school purchasing them, for the purpose of examining that school's students only for a period of 12 months from the date of receiving them. They may not be otherwise reproduced or distributed. The copyright of Neap Trial Exams remains with Neap. No Neap Trial Exam or any part thereof is to be issued or passed on by any person to any party inclusive of other schools, non-practising teachers, coaching colleges, tutors, parents, students, publishing agencies or websites without the express written consent of Neap.

Copyright © 2024 Neap Education Pty Ltd ABN 43 634 499 791 Level 1 223 Hawthorn Rd Caulfield North VIC 3161 Tel: (03) 9639 4318 VCE_SpecMaths_3&4_FS_2024

Mensuration

area of a circle segment	$\frac{r^2}{2}(\theta - \sin(\theta))$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc\sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$	sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
volume of a pyramid	$\frac{1}{3}Ah$	cosine rule	$c^2 = a^2 + b^2 - 2ab\cos(C)$

Algebra, number and structure (complex numbers)

$z = x + iy = r(\cos(\theta) + i\sin(\theta)) = r\text{cis}(\theta)$	$\lvert z\rvert = \sqrt{x^2 + y^2} = r$	
$-\pi < \text{Arg}(z) \le \pi$	$z_1 z_2 = r_1 r_2 \text{cis}(\theta_1 + \theta_2)$	
$\frac{z_1}{z_2} = \frac{r_1}{r_2}\text{cis}(\theta_1 - \theta_2)$	de Moivre's theorem	$z^n = r^n \text{cis}(n\theta)$

Data analysis, probability and statistics

for independent random variables $X_1, X_2 \ldots X_n$	$\text{E}(aX_1 + b) = a\,\text{E}(X_1) + b$ $\text{E}(a_1X_1 + a_2X_2 + \ldots + a_nX_n)$ $= a_1\text{E}(X_1) + a_2\,\text{E}(X_2) + \ldots + a_n\text{E}(X_n)$	
	$\text{Var}(aX_1 + b) = a^2\text{Var}(X_1)$ $\text{Var}(a_1X_1 + a_2X_2 + \ldots + a_nX_n)$ $= a_1^2\text{Var}(X_1) + a_2^2\text{Var}(X_2) + \ldots + a_n^2\text{Var}(X_n)$	
for independent identically distributed variables $X_1, X_2 \ldots X_n$	$\text{E}(X_1 + X_2 + \ldots + X_n) = n\mu$	
	$\text{Var}(X_1 + X_2 + \ldots + X_n) = n\sigma^2$	
approximate confidence interval for μ	$\left(\bar{x} - z\frac{s}{\sqrt{n}}, \bar{x} + z\frac{s}{\sqrt{n}}\right)$	
distribution of sample mean $\bar{X}$	mean	$\text{E}(\bar{X}) = \mu$
	variance	$\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$

Copyright © 2024 Neap Education Pty Ltd

Calculus

$\frac{d}{dx}\left(x^n\right) = n\,x^{n-1}$	$\int x^n dx = \frac{1}{n+1}x^{n+1} + c,\ n \neq -1$
$\frac{d}{dx}\left(e^{ax}\right) = a\,e^{ax}$	$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$
$\frac{d}{dx}\left(\log_e(x)\right) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e\lvert x\rvert + c$
$\frac{d}{dx}\left(\sin(ax)\right) = a\cos(ax)$	$\int \sin(ax)\,dx = -\frac{1}{a}\cos(ax) + c$
$\frac{d}{dx}\left(\cos(ax)\right) = -a\sin(ax)$	$\int \cos(ax)\,dx = \frac{1}{a}\sin(ax) + c$
$\frac{d}{dx}\left(\tan(ax)\right) = a\sec^2(ax)$	$\int \sec^2(ax)\,dx = \frac{1}{a}\tan(ax) + c$
$\frac{d}{dx}\left(\cot(ax)\right) = -a\,\mathrm{cosec}^2(ax)$	$\int \mathrm{cosec}^2(ax)\,dx = -\frac{1}{a}\cot(ax) + c$
$\frac{d}{dx}\left(\sec(ax)\right) = -a\sec(ax)\tan(ax)$	$\int \sec(ax)\tan(ax)\,dx = \frac{1}{a}\sec(ax) + c$
$\frac{d}{dx}\left(\mathrm{cosec}(ax)\right) = -a\,\mathrm{cosec}(ax)\cot(ax)$	$\int \mathrm{cosec}(ax)\cot(ax)\,dx = -\frac{1}{a}\mathrm{cosec}(ax) + c$
$\frac{d}{dx}\left(\sin^{-1}(ax)\right) = \frac{a}{\sqrt{1-(ax)^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}}dx = \sin^{-1}\left(\frac{x}{a}\right) + c,\ a > 0$
$\frac{d}{dx}\left(\cos^{-1}(ax)\right) = \frac{-a}{\sqrt{1-(ax)^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}}dx = \cos^{-1}\left(\frac{x}{a}\right) + c,\ a > 0$
$\frac{d}{dx}\left(\tan^{-1}(ax)\right) = \frac{a}{1+(ax)^2}$	$\int \frac{a}{a^2+x^2}dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c,\ n \neq -1$
	$\int \frac{1}{ax+b}dx = \frac{1}{a}\log_e\lvert ax+b\rvert + c$

Copyright © 2024 Neap Education Pty Ltd

Calculus – continued

product rule	$\frac{d}{dx}(uv) = u\frac{dv}{dx} + v\frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
integration by parts	$\int u\frac{dv}{dx}dx = uv - \int v\frac{du}{dx}dx$
Euler's method	If $\frac{dy}{dx} = f(x, y)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + h \times f(x_n, y_n)$.
arc length parametric	$\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt$
surface area Cartesian about x-axis	$\int_{x_1}^{x_2} 2\pi y\sqrt{1 + \left(\frac{dy}{dx}\right)^2}\,dx$
surface area Cartesian about y-axis	$\int_{y_1}^{y_2} 2\pi x\sqrt{1 + \left(\frac{dx}{dy}\right)^2}\,dy$
surface area parametric about x-axis	$\int_{t_1}^{t_2} 2\pi y\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt$
surface area parametric about y-axis	$\int_{t_1}^{t_2} 2\pi x\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}\,dt$

Kinematics

acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v\frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$	
constant acceleration formulas	$v = u + at$	$s = ut + \frac{1}{2}at^2$
	$v^2 = u^2 + 2as$	$s = \frac{1}{2}(u + v)t$

Copyright © 2024 Neap Education Pty Ltd

Vectors in two or three dimensions

$\underset{\sim}{r}(t) = x(t)\underset{\sim}{i} + y(t)\underset{\sim}{j} + z(t)\underset{\sim}{k}$	$\lvert\underset{\sim}{r}(t)\rvert = \sqrt{x(t)^2 + y(t)^2 + z(t)^2}$
	$\dot{\underset{\sim}{r}}(t) = \frac{d\underset{\sim}{r}}{dt} = \frac{dx}{dt}\underset{\sim}{i} + \frac{dy}{dt}\underset{\sim}{j} + \frac{dz}{dt}\underset{\sim}{k}$
for $\underset{\sim}{r}_1 = x_1\underset{\sim}{i} + y_1\underset{\sim}{j} + z_1\underset{\sim}{k}$ and $\underset{\sim}{r}_2 = x_2\underset{\sim}{i} + y_2\underset{\sim}{j} + z_2\underset{\sim}{k}$	vector scalar product $\underset{\sim}{r}_1 \cdot \underset{\sim}{r}_2 = \lvert\underset{\sim}{r}_1\rvert\lvert\underset{\sim}{r}_2\rvert\cos(\theta) = x_1x_2 + y_1y_2 + z_1z_2$
	vector cross product $\underset{\sim}{r}_1 \times \underset{\sim}{r}_2 = \begin{vmatrix} \underset{\sim}{i} & \underset{\sim}{j} & \underset{\sim}{k} \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = (y_1z_2 - y_2z_1)\underset{\sim}{i} + (x_2z_1 - x_1z_2)\underset{\sim}{j} + (x_1y_2 - x_2y_1)\underset{\sim}{k}$
vector equation of a line	$\underset{\sim}{r}(t) = \underset{\sim}{r}_1 + t\underset{\sim}{r}_2 = (x_1 + x_2t)\underset{\sim}{i} + (y_1 + y_2t)\underset{\sim}{j} + (z_1 + z_2t)\underset{\sim}{k}$
parametric equation of a line	$x(t) = x_1 + x_2t \qquad y(t) = y_1 + y_2t \qquad z(t) = z_1 + z_2t$
vector equation of a plane	$\underset{\sim}{r}(s, t) = \underset{\sim}{r}_0 + s\underset{\sim}{r}_1 + t\underset{\sim}{r}_2$ $= (x_0 + x_1s + x_2t)\underset{\sim}{i} + (y_0 + y_1s + y_2t)\underset{\sim}{j} + (z_0 + z_1s + z_2t)\underset{\sim}{k}$
parametric equation of a plane	$x(s, t) = x_0 + x_1s + x_2t, \quad y(s, t) = y_0 + y_1s + y_2t, \quad z(s, t) = z_0 + z_1s + z_2t$
Cartesian equation of a plane	$ax + by + cz = d$

Circular functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x - y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x - y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x + y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x - y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\sin(2x) = 2\sin(x)\cos(x)$	
$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$
$\sin^2(ax) = \frac{1}{2}(1 - \cos(2ax))$	$\cos^2(ax) = \frac{1}{2}(1 + \cos(2ax))$

End of Formula Sheet

Copyright © 2024 Neap Education Pty Ltd