


KINABALU — NW SABAH SECTION · ASAS-MALAM PUBLIC v1.0

Φ GEOX · L0 measured · L1 calculated · L2 interpreted


Three blocks · two margins · one suture — ages from GEOX event spine v0.1 · public sources only · internal well data excluded

Not seismic picks (F7) · no volumetrics


A · TRUE SCALE 1:1 — the whole basin system is a thin skin on a ~30 km crust: THREE BLOCKS, TWO MARGINS



B · SEDIMENT SECTION · VE ≈ 11.4× — geometry schematic (L2); horizons placed by published ages, not seismic picks



PHYSICS CHECK · flexure



EVIDENCE-TIERED LEGEND · every body = L2

post-rift / Stage IV	granite sheets
syn-rift	lower/upper crust
Stage III (Setap/Temburong)	topography/bathymetry · L0
accreted pre-BMU slices	BU/BMU ~24 Ma · L2
Rajang/Crocker	DRU ~13-12 Ma · L2
ophiolite	SRU ~8.5-9 Ma · L2
mélange	bending-extension fault (predicted)
carbonate	thrust (W-vergent)

MEASURED vs MODELLED

- L0** Topography, bathymetry, Tepat-1 public result
- L1** Granite U-Pb 7.85-7.22 Ma · Moho 26-33 km (non-unique) · flexure curves (this figure)
- L2** Every surface, body, fault, closure — depths uncalibrated (no velocity model)
- L2** Bending-extension faults: PREDICTED here — verify soles + strike || trough on 3D
- L2** Forebulge position: untested — measuring it fixes Te

Te ≈ 8-17 km fits a 110-150 km load distance — consistent with weak rifted lower plate.
BUT 2.9 km of moat relief needs sediment load PLUS underthrust downdrag — a pure sag cannot do it.
Extension at the Trough = lower-plate bending (outer-rise analog) + up-dip gravity collapse — NO spreading (5-vector falsification 2026-09-30).