

Negation and contrapositive

Guided notes and practice

Arij Asad

The question to ask: what would make the statement false? A negation is true exactly when the original statement is false. Keep the same domain (real numbers, integers, students, and so on). Write the negation before deciding whether either statement is actually true.

1. Start with a single condition

For a real number x , the statement $x > 3$ fails when $x \leq 3$. Equality must be included: $x = 3$ does not satisfy $x > 3$. Similarly,

$$\neg(x \geq 3) \iff x < 3, \quad \neg(x \neq 3) \iff x = 3.$$

2. Make “and” or “or” fail

If both parts are required, one failure is enough. If either part is allowed, both must fail:

$$\neg(P \wedge Q) \iff \neg P \vee \neg Q, \quad \neg(P \vee Q) \iff \neg P \wedge \neg Q.$$

For an integer n , “ n is positive and even” is false when $n \leq 0$ or n is odd. Ask yourself which part you could break to make the whole sentence fail.

3. Turn “every” into “some”

One exception disproves a claim about every member of a set. A claim that *some* member works fails only if none works:

$$\neg(\forall x \in D, P(x)) \iff \exists x \in D \text{ such that } \neg P(x),$$

$$\neg(\exists x \in D \text{ such that } P(x)) \iff \forall x \in D, \neg P(x).$$

Example: “Every integer n has $n^2 \geq 0$ ” negates to “There is an integer n with $n^2 < 0$.” That negation is false, but it is still the correct negation. With two quantifiers, change each in its original order: $\neg(\forall x \exists y R(x, y))$ becomes $\exists x \forall y \neg R(x, y)$.

4. An implication has two different relatives

“If P , then Q ” fails only when P happens but Q does not. Its *negation* is $P \wedge \neg Q$. Its *contrapositive* is another if-then statement, $\neg Q \Rightarrow \neg P$, with the same truth value as the original:

$$\neg(P \Rightarrow Q) \iff P \wedge \neg Q, \quad (P \Rightarrow Q) \iff (\neg Q \Rightarrow \neg P).$$

For every integer n : “If $4 \mid n$, then n is even.” Its negation is “There is an integer divisible by 4 that is odd.” Its contrapositive is “If n is odd, then $4 \nmid n$.” The converse, “If n is even, then $4 \mid n$,” is different and fails at $n = 2$.

Before you answer: name the domain, identify the condition that would fail, and check whether you need a negation (a way the statement fails) or a contrapositive (an equivalent if-then statement). “ P only if Q ” means $P \Rightarrow Q$.

Practice questions

First write the logical form, then check it with a simple example where useful.

A. Basic negations

Question 1

For a real number x , negate each statement: (a) $x > 7$; (b) $x \neq 2$.

Question 2

For an integer n , negate: “ n is positive and even.”

Question 3

For real numbers a, b , negate: “ $a < 1$ or $b \geq 4$.”

Question 4

For an integer n , negate: “ n is divisible by 3 but not by 5.”

B. Every, some and quantifiers

Question 5

Negate: “Every integer $n \geq 1$ satisfies $n^2 > n$.” If the negation is true, give a value of n that shows it.

Question 6

Negate: “There is a real number x such that $x^2 + 1 = 0$.” Decide which statement is true.

Question 7

Negate: “For every integer n , there is an integer $m > n$.” Decide which statement is true.

Question 8

Negate: “Some student solved every question on the worksheet.”

C. Implications and contrapositives**Question 9**

For every integer n : “If $8 \mid n$, then n is even.” Write (a) the negation and (b) the contrapositive.

Question 10

For every real x : “If $x^2 > 9$, then $x > 3$.” (a) Find a counterexample. (b) Negate the statement. (c) Write its contrapositive.

Question 11

“An integer is divisible by 12 only if it is divisible by 3.” (a) Rewrite this as an if-then statement. (b) Give its contrapositive. (c) Is its converse true?

D. Short proofs**Question 12**

For an integer n , prove by contrapositive: “If n^2 is odd, then n is odd.”

Question 13

For integers a, b , prove by contrapositive: “If ab is odd, then both a and b are odd.”