

LOW DOSE ASPIRIN- aspirin tablet, delayed release
Pinnacle RX LLC

MDR 981

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID-nonsteroidal anti-inflammatory drug)

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- the stomach bleeding warning applies to you

- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

These can be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. it is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose , get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose
- adults and children 12 years and older: take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- children under 12: ask a doctor

Other information

- store at 20°-25°C (68°-77°F) ; excursions permitted between 15°-30C° (59°-86°F)

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate, wax.

QUESTIONS OR COMMENTS?

package label

SODIUM BICARBONATE (UNII: 8MDF5V390O)

SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87438-0003-1	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/01/2026	

Labeler - Pinnacle RX LLC (136361430)

Registrant - Geri-Care Pharmaceutical Corp (611196254)