

ASPIRIN ADULT LOW DOSE- acetylsalicylic acid tablet, delayed release
Medline Industries, LP

985 Aspirin 81mg EC tablets

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- if you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or loss of hearing occurs.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and older: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless otherwise directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 25°C(77°F) excursions permitted between 15°-30°C (59°-68°F)
- use by expiration date on package

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Manufacturing Information

Manufactured by Medline Industries, LP
Three Lakes Drive, Northfield, IL 60093 USA

Made in the USA with U.S. and Foreign Components

www.medline.com

1-800-MEDLINE (633-5463)

REF: OTCM00003

V2 RH22HND

Package Label



NDC: 53329-985-67

Adult Low Dose
Aspirin
Pain Reliever (NSAID)*

**Compare to the active ingredient in Bayer® Low Dose*

120 ENTERIC COATED TABLETS



81 mg

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*

Purpose
Pain reliever

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach

REF: OTCM00003 (CONTINUED ON BACK OF LABEL)

*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Low Dose.

www.medline.com Made in the USA with U.S. and Foreign Components. Manufactured for Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA 1-800-MEDLINE V2-RH22HND



8 88277 69599 6

LOT
EXP

Coating Free

PEEL HERE

Drug Facts (continued)

ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for: ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and older: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless otherwise directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F) excursions permitted between 15°–30°C (59°–86°F) ■ use by expiration date on package

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

ASPIRIN ADULT LOW DOSE

acetylsalicyclic acid tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53329-985
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics			
Color	yellow	Score	score with uneven pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	embossed;heart
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-985-67	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/2018	04/30/2027

Marketing Information				
-----------------------	--	--	--	--

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/25/2018	04/30/2027

Labeler - Medline Industries, LP (025460908)

Registrant - Medline Industries, LP (025460908)

Revised: 6/2026

Medline Industries, LP