

IBUPROFEN- ibuprofen tablet, film coated
L.N.K. International, Inc.

Quality Plus 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin reddening
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - weakness in one part or side of body
 - leg swelling
 - slurred speech
 - trouble breathing
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**Quality
+Plus**

NDC 50844-291-16

†Compare to active ingredient
in Advil® Tablets

**IBUPROFEN TABLETS
USP, 200 mg**

PAIN RELIEVER/
FEVER REDUCER (**NSAID**)

1000 Coated Tablets

ACTUAL

SIZE

QUALITY PLUS

NDC 50844-291-16

†Compare to active ingredient in Advil® Tablets

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER/
FEVER REDUCER (NSAID)

1000 Coated Tablets

ACTUAL SIZE

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each brown tablet)

Ibuprofen USP 200 mg (NSAID)*. *Nonsteroidal anti-inflammatory drug.

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:
headache toothache backache menstrual cramps
the common cold muscular aches minor pain of arthritis
temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling
asthma (wheezing) skin reddening shock rash blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

if you have ever had an allergic reaction to any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if

stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers or fever reducers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
under a doctor's care for any serious condition
taking any other drug

Drug Facts (continued)

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:
feel faint vomit blood have bloody or black stools
have stomach pain that does not get better
you have symptoms of heart problems or stroke:
chest pain weakness in one part or side of body leg swelling
slurred speech trouble breathing
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days any new symptoms appear
redness or swelling is present in the painful area

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed
the smallest effective dose should be used
adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
if pain or fever does not respond to 1 tablet, 2 tablets may be used
do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years: ask a doctor

Other information

store between 20°-25°C (68°-77°F)
avoid excessive heat 40°C (104°F)
use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

This product is not manufactured or distributed by PF Consumer Healthcare LLC, owner of the registered trademark Advil® Tablets. NDC 50844-291-16 REV1221C29116

Distributed by
LWK INTERNATIONAL, INC.
60 Akar Drive, Hauppauge, NY 11788
USA

Quality Plus 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-291
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Basis of Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN
		200 mg

Inactive Ingredients	
Ingredient Name	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWL1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-291-50	50 in 1 CARTON	05/24/1988	
1	NDC:50844-291-24	2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:50844-291-03	10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
3	NDC:50844-291-02	1 in 1 CARTON	05/24/1988	
3		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:50844-291-08	1 in 1 CARTON	05/24/1988	
4		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:50844-291-01	1 in 1 CARTON	05/24/1988	
5		30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:50844-291-10	1 in 1 CARTON	05/24/1988	
6		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:50844-291-15	1 in 1 CARTON	05/24/1988	
7		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:50844-291-11	1 in 1 CARTON	05/24/1988	
8		60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
9	NDC:50844-291-12	1 in 1 CARTON	05/24/1988	
9		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
10	NDC:50844-291-29	1 in 1 CARTON	05/24/1988	
10		150 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
11	NDC:50844-291-99	250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
12	NDC:50844-291-13	1 in 1 CARTON	05/24/1988	
12		250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

13	NDC:50844-291-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
14	NDC:50844-291-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
15	NDC:50844-291-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-291) , pack(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117597853	pack(50844-291)