

**ALKA SELTZER COLD COLD FLU CONGESTION SEVERE FAST HOT LIQUID DAY NIGHT- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate
Bayer HealthCare LLC.**

**ALKA SELTZER COLD Cold Flu Congestion Severe Fast Hot Liquid Day Night UI
1611758/1611711**

Alka-Seltzer COLD

Cold & Flu + Congestion Severe Day Powder Packets

<i>Active ingredients (in each packet)</i>	<i>Purposes</i>
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - sore throat
 - cough
 - sinus congestion and pressure
 - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and makes cough more productive
- temporarily reduces fever

Warnings

.

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 packets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening

- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that is accompanied by excessive phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take every 4 hours; do not exceed 6 packets in 24 hours or as directed by a doctor
- dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

Other information

- **each packet contains:** potassium 5 mg; sodium 5 mg
- store between 20 °C to 25 °C (68 °F to 77 °F)

Inactive ingredients

acesulfame potassium, anhydrous citric acid, compressible sugar, D&C yellow #10, dental-type silica, FD&C red #40, flavors, maltodextrin, povidone, pregelatinized starch, silicon dioxide, sodium citrate, stearic acid, sucralose, tartaric acid, tribasic calcium phosphate

Questions or comments?

1-800-986-0369 (Mon-Fri 8AM - 8PM EST)

Alka-Seltzer COLD

Cold & Flu + Congestion Severe Night Powder Packets

<i>Active ingredients (in each packet)</i>	<i>Purposes</i>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - headache
 - minor aches and pains
 - cough
 - sore throat

- nasal congestion
- sinus congestion and pressure
- runny nose
- sneezing
- temporarily reduces fever

Warnings

.

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 5 packets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that is accompanied by excessive phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- may cause marked drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over : take every 4 hours; do not exceed 5 packets in 24 hours or as directed by a doctor
- dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

Other information

- **each packet contains:** potassium 5 mg; sodium 5 mg

- ALKA SELTZER COLD COLD FLU CONGESTION SEVERE FAST HOT
LIQUID DAY NIGHT**

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-8284

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-8284-12	1 in 1 CARTON; Type 0: Not a Combination Product	06/01/2026	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	6 PACKET	6
Part 2	6 PACKET	6

Part 1 of 2

ALKA SELTZER COLD COLD FLU AND CONGESTION SEVERE FAST HOT LIQUID DAY

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride powder, for solution

Product Information	
Item Code (Source)	NDC:0280-8273
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients	
Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	
SILICA (UNII: ETJ7Z 6XBU4)	

STARCH, CORN (UNII: O8232NY3SJ)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POVIDONE (UNII: FZ989GH94E)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TARTARIC ACID (UNII: W4888I119H)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	

Product Characteristics

Color	yellow	Score	
Shape		Size	
Flavor	LEMON, HONEY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2026	

Part 2 of 2

ALKA SELTZER COLD COLD AND FLU PLUS CONGESTION SEVERE FAST HOT LIQUID NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride powder, for solution

Product Information

Item Code (Source)	NDC:0280-8272
Route of Administration	ORAL

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	650 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	20 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)			DOXYLAMINE SUCCINATE	12.5 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)			PHENYLEPHRINE HYDROCHLORIDE	10 mg
Inactive Ingredients				
Ingredient Name				Strength
SILICA (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
SUCROSE (UNII: C151H8M554)				
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
TARTARIC ACID (UNII: W4888I119H)				
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)				
Product Characteristics				
Color	yellow		Score	
Shape			Size	
Flavor	HONEY, LEMON		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/01/2026	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC Monograph Drug	M012	06/01/2026	
--------------------	------	------------	--

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 6/2026

Bayer HealthCare LLC.