

IBUPROFEN- ibuprofen tablet, film coated
L.N.K. International, Inc.

Quality Plus 44-392

Active ingredient (in each orange tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin reddening
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - weakness in one part or side of body
 - trouble breathing
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**QUALITY
+PLUS**

NDC 50844-392-16

†Compare to active ingredient
in Motrin® IB

**IBUPROFEN
TABLETS
USP, 200 mg**

**PAIN RELIEVER/FEVER REDUCER
(NSAID)**

1000 Coated Tablets

ACTUAL SIZE

NDC 50844-392-16

†Compare to active ingredient in Motrin® IB

QUALITY PLUS

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

1000 Coated Tablets

ACTUAL SIZE

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each orange tablet) Purpose

Ibuprofen USP, 200 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache toothache backache menstrual cramps

the common cold muscular aches minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives skin reddening shock rash blisters

asthma (wheezing) skin reddening shock rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

have 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

if you have ever had an allergic reaction to any other pain reliever/fever reducer

right before or after heart surgery

Ask a doctor before use if

stomach bleeding warning applies to you

you have problems or serious side effects from taking pain relievers or fever reducers

you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

under a doctor's care for any serious condition

taking any other drug

Drug Facts (continued)

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint vomit blood have bloody or black stools

have stomach pain that does not get better

you have symptoms of heart problems or stroke:

chest pain weakness in one part or side of body trouble breathing

slurred speech leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

any new symptoms appear

redness or swelling is present in the painful area

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed

the smallest effective dose should be used

adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist.

if pain or fever does not respond to 1 tablet, 2 tablets may be used

do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

Other information

avoid excessive heat 40°C (104°F)

store between 20°-25°C (68°-77°F)

use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB.

50844 REV12/12/13/216

Distributed by

LINK INTERNATIONAL, INC.

60 Arroyo Drive, Hauppauge, NY 11788

USA

0 50844 39216 8

Quality Plus 44-392

IBUPROFEN

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-392
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			
Ingredient Name			Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POLYDEXTROSE (UNII: VH2XOU12IE)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;392	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-392-50	50 in 1 CARTON	03/01/1999	
1	NDC:50844-392-24	2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:50844-392-02	1 in 1 CARTON	03/01/1999	
2		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:50844-392-08	1 in 1 CARTON	03/01/1999	
3		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:50844-392-15	1 in 1 CARTON	03/01/1999	
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:50844-392-12	1 in 1 CARTON	03/01/1999	
5		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:50844-392-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
7	NDC:50844-392-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA075139		03/01/1999	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-392)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-392)