


Ligjerata 10ç

Bymimi termik linear i trupave

$\alpha \rightarrow$ koeficienti termik i
shtimit linear

$$\Delta l_r = \frac{\Delta l}{l_0}$$

$$\Delta l = l - l_0$$



$$\alpha = \frac{\Delta l_r}{\Delta t} = \frac{\Delta l}{l_0 \cdot \Delta t} \Rightarrow \Delta l = l_0 \cdot \alpha \cdot \Delta t$$

$$l - l_0 = l_0 \cdot \alpha \cdot \Delta t$$

$$l = l_0 + l_0 \cdot \alpha \cdot \Delta t$$

$$l = l_0 (1 + \alpha \Delta t)$$

$$\Delta t = t - \cancel{t_0}^\circ = t$$

$$l = l_0 (1 + \alpha t)$$

$$\Delta t = \Delta T$$


$$\alpha = \frac{1}{\Delta t} \cdot \frac{\Delta l}{l_0} [\Rightarrow] \frac{1}{^\circ\text{C}} = \frac{\mu\text{m}}{\mu^\circ\text{C}} = \frac{1}{^\circ\text{C}} = \frac{1}{\text{K}}$$

Bymimi termik sipërfaqësor i trupave

$$S = a \cdot b$$

$$a = a_0 (1 + \alpha t)$$

$$b = b_0 (1 + \alpha t)$$



$$S = a_0 (1 + \alpha t) \cdot b_0 (1 + \alpha t)$$

$$S = S_0 (1 + \alpha t + \alpha t + \alpha^2 t^2)$$

$$S_0 = a_0 \cdot b_0$$

$$\alpha \approx 10^{-3} \quad \alpha^2 \rightarrow 0$$

$$S = S_0 (1 + 2\alpha t)$$

$$\gamma = 2\alpha$$

$$S = S_0 (1 + \gamma t)$$