

# Probability Review and Statistical Estimation

Instructor: Alan Ritter

Many slides from Tom Mitchell, Pedro Domingos

# Random Variables

- Informally,  $A$  is a random variable if
  - $A$  denotes something about which we are uncertain
  - perhaps the outcome of a randomized experiment
- Examples
  - $A$  = True if a randomly drawn person from our class is female
  - $A$  = The hometown of a randomly drawn person from our class
  - $A$  = True if two randomly drawn persons from our class have same birthday
- Define  $P(A)$  as “the fraction of possible worlds in which  $A$  is true” or “the fraction of times  $A$  holds, in repeated runs of the random experiment”
  - the set of possible worlds is called the sample space,  $S$
  - A random variable  $A$  is a function defined over  $S$ 
$$A: S \rightarrow \{0,1\}$$

# A little formalism

More formally, we have


- a sample space  $S$  (e.g., set of students in our class)
  - aka the set of possible worlds
- a random variable is a function defined over the sample space
  - Gender:  $S \rightarrow \{m, f\}$
  - Height:  $S \rightarrow \text{Reals}$
- an event is a subset of  $S$ 
  - e.g., the subset of  $S$  for which Gender=f
  - e.g., the subset of  $S$  for which (Gender=m) AND (eyeColor=blue)
- we're often interested in probabilities of specific events
- and of specific events conditioned on other specific events

# Visualizing A

Sample space  
of all possible  
worlds



Its area is 1



$P(A)$  = Area of  
reddish oval

# The Axioms of Probability

- $0 \leq P(A) \leq 1$
- $P(\text{True}) = 1$
- $P(\text{False}) = 0$
- $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$

[di Finetti 1931]:


when gambling based on “uncertainty formalism A” you can be exploited by an opponent

iff

your uncertainty formalism A violates these axioms


# Elementary Probability in Pictures

- $P(\sim A) + P(A) = 1$




# Elementary Probability in Pictures

- $P(A) = P(A \wedge B) + P(A \wedge \sim B)$



# Definition of Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$





# Definition of Conditional Probability


$$P(A|B) = \frac{P(A \wedge B)}{P(B)}$$

## Corollary: The Chain Rule

$$P(A \wedge B) = P(A|B) P(B)$$

# Bayes Rule

- let's write 2 expressions for  $P(A \wedge B)$



$$P(A|B) = \frac{P(B|A) * P(A)}{P(B)}$$
 Bayes' rule

we call  $P(A)$  the “prior”

and  $P(A|B)$  the “posterior”



**Bayes, Thomas (1763)** An essay towards solving a problem in the doctrine of chances. *Philosophical Transactions of the Royal Society of London*, **53:370-418**

...by no means merely a curious speculation in the doctrine of chances, but necessary to be solved in order to a sure foundation for all our reasonings concerning past facts, and what is likely to be hereafter.... necessary to be considered by any that would give a clear account of the strength of *analogical* or *inductive reasoning*...

## Other Forms of Bayes Rule

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

$$P(A|B \wedge X) = \frac{P(B|A \wedge X)P(A \wedge X)}{P(B \wedge X)}$$

# Applying Bayes Rule

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

A = you have the flu, B = you just coughed

Assume:

$$P(A) = 0.05$$

$$P(B|A) = 0.80$$

$$P(B|\sim A) = 0.2$$

what is  $P(\text{flu} | \text{cough}) = P(A|B)$ ?

what does all this have to do with  
function approximation?

# The Joint Distribution

*Example: Boolean  
variables  $A, B, C$*

Recipe for making a joint  
distribution of  $M$  variables:

# The Joint Distribution

*Example: Boolean variables A, B, C*

Recipe for making a joint distribution of M variables:

1. Make a truth table listing all combinations of values of your variables (if there are M Boolean variables then the table will have  $2^M$  rows).

<b>A</b>	<b>B</b>	<b>C</b>
0	0	0
0	0	1
0	1	0
0	1	1
1	0	0
1	0	1
1	1	0
1	1	1



# The Joint Distribution

*Example: Boolean variables A, B, C*

Recipe for making a joint distribution of M variables:

1. Make a truth table listing all combinations of values of your variables (if there are M Boolean variables then the table will have  $2^M$  rows).
2. For each combination of values, say how probable it is.


<b>A</b>	<b>B</b>	<b>C</b>	<b>Prob</b>
0	0	0	0.30
0	0	1	0.05
0	1	0	0.10
0	1	1	0.05
1	0	0	0.05
1	0	1	0.10
1	1	0	0.25
1	1	1	0.10

# The Joint Distribution









Recipe for making a joint distribution of  $M$  variables:

1. Make a truth table listing all combinations of values of your variables (if there are  $M$  Boolean variables then the table will have  $2^M$  rows).
2. For each combination of values, say how probable it is.
3. If you subscribe to the axioms of probability, those numbers must sum to 1.

A	B	C	Prob
0	0	0	0.30
0	0	1	0.05
0	1	0	0.10
0	1	1	0.05
1	0	0	0.05
1	0	1	0.10
1	1	0	0.25
1	1	1	0.10



# Using the Joint Distribution

gender	hours_worked	wealth		
Female	v0:40.5-	poor	0.253122	
		rich	0.0245895	
	v1:40.5+	poor	0.0421768	
		rich	0.0116293	
Male	v0:40.5-	poor	0.331313	
		rich	0.0971295	
	v1:40.5+	poor	0.134106	
		rich	0.105933	

Once you have the JD  
you can ask for the  
probability of **any** logical  
expression involving  
these variables

$$P(E) = \sum_{\text{rows matching } E} P(\text{row})$$

# Using the Joint

gender	hours_worked	wealth		
Female	v0:40.5-	poor	0.253122	<div></div>
		rich	0.0245895	<div></div>
	v1:40.5+	poor	0.0421768	<div></div>
		rich	0.0116293	<div></div>
Male	v0:40.5-	poor	0.331313	<div></div>
		rich	0.0971295	<div></div>
	v1:40.5+	poor	0.134106	<div></div>
		rich	0.105933	<div></div>

$$P(\text{Poor Male}) = 0.4654$$

$$P(E) = \sum_{\text{rows matching } E} P(\text{row})$$

# Using the Joint

gender	hours_worked	wealth	
Female	v0:40.5-	poor	0.253122
		rich	0.0245895
	v1:40.5+	poor	0.0421768
		rich	0.0116293
Male	v0:40.5-	poor	0.331313
		rich	0.0971295
	v1:40.5+	poor	0.134106
		rich	0.105933

$$P(\text{Poor}) = 0.7604$$

$$P(E) = \sum_{\text{rows matching } E} P(\text{row})$$









# Inference with the Joint

gender	hours_worked	wealth	
Female	v0:40.5-	poor	0.253122
		rich	0.0245895
	v1:40.5+	poor	0.0421768
		rich	0.0116293
Male	v0:40.5-	poor	0.331313
		rich	0.0971295
	v1:40.5+	poor	0.134106
		rich	0.105933

$$P(E_1 | E_2) = \frac{P(E_1 \wedge E_2)}{P(E_2)} = \frac{\sum_{\text{rows matching } E_1 \text{ and } E_2} P(\text{row})}{\sum_{\text{rows matching } E_2} P(\text{row})}$$

$$P(\text{Male} | \text{Poor}) = 0.4654 / 0.7604 = 0.612$$

# Learning and the Joint Distribution

gender	hours_worked	wealth		
Female	v0:40.5-	poor	0.253122	
		rich	0.0245895	
	v1:40.5+	poor	0.0421768	
		rich	0.0116293	
Male	v0:40.5-	poor	0.331313	
		rich	0.0971295	
	v1:40.5+	poor	0.134106	
		rich	0.105933	

Suppose we want to learn the function  $f: \langle G, H \rangle \rightarrow W$

Equivalently,  $P(W \mid G, H)$

Solution: learn joint distribution from data, calculate  $P(W \mid G, H)$

e.g.,  $P(W=\text{rich} \mid G = \text{female}, H = 40.5- ) =$

sounds like the solution to  
learning  $F: X \rightarrow Y$ ,  
or  $P(Y | X)$ .

Are we done?



sounds like the solution to  
learning  $F: X \rightarrow Y$ ,  
or  $P(Y | X)$ .

Main problem: learning  $P(Y|X)$   
can require more data than we have

consider learning Joint Dist. with 100 attributes

# of rows in this table?

# of people on earth?

fraction of rows with 0 training examples?

# What to do?

1. Be smart about how we estimate probabilities from sparse data
  - maximum likelihood estimates
  - maximum a posteriori estimates
2. Be smart about how to represent joint distributions
  - Bayes networks, graphical models

1. Be smart about how we estimate probabilities

# Estimating Probability of Heads



- I show you the above coin  $X$ , and hire you to estimate the probability that it will turn up heads ( $X = 1$ ) or tails ( $X = 0$ )
- You flip it repeatedly, observing
  - it turns up heads  $\alpha_1$  times
  - it turns up tails  $\alpha_0$  times
- Your estimate for  $P(X = 1)$  is....?

# Estimating $\theta = P(X=1)$



X=1

X=0

Test A:

100 flips: 51 Heads (X=1), 49 Tails (X=0)

Test B:

3 flips: 2 Heads (X=1), 1 Tails (X=0)

# Estimating $\theta = P(X=1)$



X=1

X=0

Case C: (online learning)

- keep flipping, want single learning algorithm that gives reasonable estimate after each flip

# Principles for Estimating Probabilities

Principle 1 (maximum likelihood):

- choose parameters  $\theta$  that maximize  **$P(\text{data} \mid \theta)$**

- e.g., 
$$\hat{\theta}^{MLE} = \frac{\alpha_1}{\alpha_1 + \alpha_0}$$

Principle 2 (maximum a posteriori prob.):

- choose parameters  $\theta$  that maximize  **$P(\theta \mid \text{data})$**
- e.g.

$$\hat{\theta}^{MAP} = \frac{\alpha_1 + \# \text{hallucinated\_1s}}{(\alpha_1 + \# \text{hallucinated\_1s}) + (\alpha_0 + \# \text{hallucinated\_0s})}$$

# Maximum Likelihood Estimation



$$P(X=1) = \theta \quad P(X=0) = (1-\theta)$$

Data D:


Flips produce data D with  $\alpha_1$  heads,  $\alpha_0$  tails

- flips are independent, identically distributed 1's and 0's (Bernoulli)
- $\alpha_1$  and  $\alpha_0$  are counts that sum these outcomes (Binomial)

$$P(D|\theta) = P(\alpha_1, \alpha_0|\theta) = \theta^{\alpha_1} (1 - \theta)^{\alpha_0}$$



# Maximum Likelihood Estimate for $\Theta$


$$\begin{aligned}\hat{\theta} &= \arg \max_{\theta} \ln P(\mathcal{D} \mid \theta) \\ &= \arg \max_{\theta} \ln \theta^{\alpha_H} (1 - \theta)^{\alpha_T}\end{aligned}$$

■ Set derivative to zero:

$$\frac{d}{d\theta} \ln P(\mathcal{D} \mid \theta) = 0$$

$$\hat{\theta} = \arg \max_{\theta} \ln P(D|\theta)$$

■ Set derivative to zero:

$$\frac{d}{d\theta} \ln P(\mathcal{D} | \theta) = 0$$

$$= \arg \max_{\theta} \ln [\theta^{\alpha_1} (1 - \theta)^{\alpha_0}]$$

hint:  $\frac{\partial \ln \theta}{\partial \theta} = \frac{1}{\theta}$

# Summary:

## Maximum Likelihood Estimate



$X=1$     $X=0$

$$P(X=1) = \theta$$

$$P(X=0) = 1-\theta$$

(Bernoulli)

- Each flip yields boolean value for  $X$

$$X \sim \text{Bernoulli}: P(X) = \theta^X (1 - \theta)^{(1-X)}$$

- Data set  $D$  of independent, identically distributed (iid) flips produces  $\alpha_1$  ones,  $\alpha_0$  zeros (Binomial)

$$P(D|\theta) = P(\alpha_1, \alpha_0|\theta) = \theta^{\alpha_1} (1 - \theta)^{\alpha_0}$$

$$\hat{\theta}^{MLE} = \operatorname{argmax}_{\theta} P(D|\theta) = \frac{\alpha_1}{\alpha_1 + \alpha_0}$$

# Principles for Estimating Probabilities

Principle 1 (maximum likelihood):

- choose parameters  $\theta$  that maximize  $P(\text{data} \mid \theta)$

Principle 2 (maximum a posteriori prob.):

- choose parameters  $\theta$  that maximize


$$P(\theta \mid \text{data}) = \frac{P(\text{data} \mid \theta) P(\theta)}{P(\text{data})}$$

## Beta prior distribution – $P(\theta)$

- $$P(\theta) = \frac{\theta^{\beta_H-1}(1-\theta)^{\beta_T-1}}{B(\beta_H, \beta_T)} \sim \text{Beta}(\beta_H, \beta_T)$$
- Likelihood function:  $P(\mathcal{D} \mid \theta) = \theta^{\alpha_H}(1-\theta)^{\alpha_T}$
- Posterior:  $P(\theta \mid \mathcal{D}) \propto P(\mathcal{D} \mid \theta)P(\theta)$

# Beta prior distribution – $P(\theta)$

- $$P(\theta) = \frac{\theta^{\beta_H-1}(1-\theta)^{\beta_T-1}}{B(\beta_H, \beta_T)} \sim \text{Beta}(\beta_H, \beta_T)$$



## Eg. 1 Coin flip problem

Likelihood is  $\sim$  Binomial

$$P(\mathcal{D} | \theta) = \theta^{\alpha_H} (1 - \theta)^{\alpha_T}$$

If prior is Beta distribution,

$$P(\theta) = \frac{\theta^{\beta_H-1} (1 - \theta)^{\beta_T-1}}{B(\beta_H, \beta_T)} \sim \text{Beta}(\beta_H, \beta_T)$$

Then posterior is Beta distribution

$$P(\theta | D) \sim \text{Beta}(\alpha_H + \beta_H, \alpha_H + \beta_H)$$

and MAP estimate is therefore

$$\hat{\theta}^{MAP} = \frac{\alpha_H + \beta_H - 1}{(\alpha_H + \beta_H - 1) + (\alpha_T + \beta_T - 1)}$$



**Eg. 2** Dice roll problem (6 outcomes instead of 2)



Likelihood is  $\sim \text{Multinomial}(\theta = \{\theta_1, \theta_2, \dots, \theta_k\})$

$$P(\mathcal{D} | \theta) = \theta_1^{\alpha_1} \theta_2^{\alpha_2} \dots \theta_k^{\alpha_k}$$

If prior is Dirichlet distribution,

$$P(\theta) = \frac{\theta_1^{\beta_1-1} \theta_2^{\beta_2-1} \dots \theta_k^{\beta_k-1}}{B(\beta_1, \dots, \beta_k)} \sim \text{Dirichlet}(\beta_1, \dots, \beta_k)$$

Then posterior is Dirichlet distribution

$$P(\theta | D) \sim \text{Dirichlet}(\beta_1 + \alpha_1, \dots, \beta_k + \alpha_k)$$

and MAP estimate is therefore

$$\hat{\theta}_i^{MAP} = \frac{\alpha_i + \beta_i - 1}{\sum_{j=1}^k (\alpha_j + \beta_j - 1)}$$



## Some terminology

- Likelihood function:  $P(\text{data} \mid \theta)$
- Prior:  $P(\theta)$
- Posterior:  $P(\theta \mid \text{data})$
- Conjugate prior:  $P(\theta)$  is the conjugate prior for likelihood function  $P(\text{data} \mid \theta)$  if the forms of  $P(\theta)$  and  $P(\theta \mid \text{data})$  are the same.

# You should know

- Probability basics
  - random variables, conditional probs, ...
  - Bayes rule
  - Joint probability distributions
  - calculating probabilities from the joint distribution
- Estimating parameters from data
  - maximum likelihood estimates
  - maximum a posteriori estimates
  - distributions – binomial, Beta, Dirichlet, ...
  - conjugate priors