1 Reconstruct a connection once and sample at finer spacings



$U^{(0)} = U$



$U^{(1)} = F_{1/3}U$



$U^{(2)} = F_{1/9}U$ ...

2 Compare chiral projectors at adjacent spacings

$\widehat{P}_-[U^{(0)}]$ $\xrightarrow{B_0}$



$\widehat{P}_-[U^{(1)}]$ $\xrightarrow{B_1}$

$\widehat{P}_-[U^{(2)}]$ ...

$\mathrm{d}B_n = \Omega_n - \Omega_{n+1}$ $\qquad$ $e^{i\oint_\gamma B_n} = \dfrac{\mathcal{T}_n[\gamma]}{\mathcal{T}_{n+1}[\gamma]}$

3a Control the local current



Color-trace cancellation

$\Omega_n \longrightarrow 0$

Gauge variation cancels the anomaly

$\mathfrak{L} = \sum_{n\geq 0} B_n, \qquad \mathrm{d}\mathfrak{L} = \Omega$

3b Control the global phase



auxiliary connection $\qquad$ conjugate pairs

$\mathcal{T}[F_h\gamma] \longrightarrow 1$

$W[\gamma] = e^{i\oint_\gamma \mathfrak{L}} = \mathcal{T}[\gamma]$ for every closed γ

4 Reconstruct a path-independent measure



Gauge-invariant measure on the original lattice

$s(U) \longleftrightarrow \mathcal{D}[\psi]\,\mathcal{D}[\bar{\psi}] = e^{i\varphi_v(U)} \prod_k \mathrm{d}c_k \prod_l \mathrm{d}\bar{c}_l$

$\longrightarrow$ Restrict to the four Standard Model groups

$[\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)]/\Gamma_{\mathrm{SM}}, \qquad \Gamma_{\mathrm{SM}} = 1, \mathbb{Z}_2, \mathbb{Z}_3, \mathbb{Z}_6$