

Openings for the mass gap, and why it is not special to $SU(3)$

Recorded 2026-09-23 at the user's request and **parked**: the Planck-gap paper comes first, and none of these is being attacked now, except possibly the fourth, which is the link from the Planck-gap work. The note answers two questions the user asked and corrects one statement of the map.

The gap is expected for every compact simple non-abelian group. Lattice evidence shows gapped glueball spectra for $SU(N)$ from $N = 2$ to large N , with masses smooth in $1/N^2$, and for the other classical groups; the dividing line is between abelian and non-abelian, since compact $U(1)$ in four dimensions has a proven massless phase at weak coupling. $SU(3)$ fixes the constants; a method that works will almost surely be group-general.

A precision on the map. The position note says that the conjecture is T2' together with T3, given T4, and the obligations map says that T3 presupposes T2'. T3 needs less: a gap $\delta_\infty(g) > 0$ on some interval $(0, g_1)$ along the scaling curve. T2', a gap at every lattice coupling, is a property of a particular lattice action, and for the Wilson action it fails for $SU(N \geq 5)$, where a first-order bulk transition occurs at intermediate coupling. For $SU(3)$ with the Wilson action the bulk behaviour is a crossover, so T2' remains plausible there; the conditional theorem of the conditional-theorem note, stated along the trajectory, needs no T2' at all.

Four openings, ranked by how directly they reach the place where the map ends, the control of about six blocking steps at order-one coupling.

1. Curvature: the Bakry–Émery criterion and its multiscale form.
2. The centre-stabilized theory on $\mathbb{R}^3 \times S^1$ at small radius.
3. Three dimensions first.
4. The Planck-gap link: Simon's valley lifting, a floor that holds for every state.

1. Evidence across groups

- $SU(N)$. Glueball masses and string tensions computed for N from 2 to about 12 are finite and extrapolate smoothly in $1/N^2$ (Lucini and Panero, Phys. Rep. **526**, 93, 2013, metadata; Lucini and Teper, JHEP **2001**(06), 050, metadata).
- G_2 . The exceptional group has a trivial centre and still shows confinement behaviour and a first-order deconfinement transition (Pepe and Wiese, Nucl. Phys. B **768**, 21, 2007, metadata), so the centre is not what the gap rests on.
- $U(1)$. Four-dimensional compact $U(1)$ has a massless Coulomb phase at weak coupling (Guth, PRD **21**, 2291, 1980; Fröhlich and Spencer, CMP **83**, 411, 1982; metadata), which is why the abelian note separates the groups by the running, $b_0 = 0$ against $b_0 = 11N/(48\pi^2)$. In three dimensions compact $U(1)$ is gapped at every coupling (Polyakov, Nucl. Phys. B **120**, 429, 1977; proved by Göpfert and Mack, CMP **82**, 545, 1982; metadata), with a gap that vanishes in physical units in the continuum.

Bulk transitions depend on the action. With the standard Wilson plaquette action, “the bulk transition is a rapid cross-over if $N \leq 4$ but becomes an increasingly strong first order transition for $N \geq 5$ ”; for $N = 3$ a negative adjoint term avoids the bulk phase altogether (Rindlisbacher, Rummukainen and Salami, arXiv:2306.14319, passage). In the plane of fundamental and adjoint couplings, a line of first-order transitions ends at a critical point near the Wilson axis (Bhanot and Creutz, PRD **24**, 3212, 1981, metadata). By Corollary 3 of the gapped-set note, a first-order transition at g_b gives $\delta_\infty(g_b) = 0$ through vacuum coexistence, so the gapped set $\mathcal{G}$ has a hole there while the physical gap, which lives at $g \rightarrow 0$, is untouched. The statement the conjecture needs is $\mathcal{G} \supseteq (0, g_1)$ for some $g_1 > 0$, joined to T3 along the scaling curve.

2. The openings

Opening 1: curvature

The configuration space of lattice gauge theory is a product of copies of the group, and a compact simple group carries positive Ricci curvature: for $SU(N)$ with the metric $\langle X, Y \rangle = \text{Tr}(XY^*)$, $\text{Ric} = \frac{N}{2}g$. The Bakry–Émery criterion turns a positive lower bound on $\text{Ric} + \text{Hess } S$ into a log-Sobolev and a Poincaré inequality, that is into a spectral gap of the Langevin dynamics and exponential decay of correlations (Bakry and Émery, LNM 1123, 177, 1985, metadata). A flat group, $U(1)$, gets nothing from the first term, which lines up with the abelian failure.

Shen, Zhu and Zhu use exactly this to prove, for lattice Yang–Mills with ’t Hooft coupling βN in any dimension $d > 1$, uniqueness of the infinite-volume measure, log-Sobolev and Poincaré inequalities and exponential decay of correlations, “a strictly positive mass gap”, for $|\beta| < 1/(16(d-1))$ when the group is $SU(N)$ (CMP **400**, 2022; arXiv:2204.12737, abstract). In the Wilson normalization of this repository, $\beta_W = 2N/g^2$ with weight $e^{(\beta_W/N) \sum \text{Re Tr } U_p}$, their β is β_W/N^2 , so for $SU(3)$ in four dimensions the condition is $\beta_W < 3/16$, that is

$$g^2 > 32.$$

That conversion is ours and must be checked against their definitions. If it stands, the curvature method already beats this repository’s strong-coupling thresholds, $g^2 \geq 176$ for the Wilson transfer matrix (note), $g^2 \geq 388$ for Kogut–Susskind (note) and $g^2 > 444$ by Dobrushin (note), and narrows band B of the bands note from $\beta_W \in (0.0135, 5.7)$ to $\beta_W \in (0.19, 5.7)$.

The weak side is where the multiscale form enters. Bauerschmidt and Bodineau derive “a multiscale generalisation of the Bakry–Émery criterion” through the Polchinski renormalization equation, “effective for measures which are far from log-concave”, and prove optimal log-Sobolev inequalities for the continuum sine-Gordon model at $\beta < 6\pi$ (CPAM **74**, 2020; arXiv:1907.12308, abstract). Its output is a spectral gap controlled at every scale, which is the form of hypothesis H1 in the conditional theorem. It also sits outside the family closed in flow-before-decimation: that note closes every criterion that depends on the perturbation through a norm, and a curvature bound is a lower bound on a Hessian.

The risk is concrete. After a blocking step the effective action is a function on the coarse configuration manifold, and the criterion needs $\text{Ric} + \text{Hess } S_{\text{eff}}$ bounded below at every scale. At weak coupling the Hessian is degenerate along gauge orbits and along the abelian valleys, and the curvature of the group has to carry those directions. **First note if opened:** check the conversion to $g^2 > 32$; compute the curvature budget for $SU(2)$ and $SU(3)$ against the Hessian of the Wilson action; and state exactly what lower bound one blocking step must preserve.

Opening 2: the centre-stabilized theory at small radius

Ünsal and Yaffe add a double-trace deformation that keeps centre symmetry unbroken; on $\mathbb{R}^3 \times S^1$ at small radius “the deformed Yang–Mills theory has a mass gap and exhibits linear confinement”, by an analytic semiclassical treatment, and “there are no order parameters which distinguish the small and large radius regimes” (PRD **78**, 065035, 2008, abstract). The mechanism at small radius is Polyakov’s monopole plasma, which Göpfert and Mack made rigorous for three-dimensional compact $U(1)$ at every coupling. **Target if opened:** a rigorous mass gap for the deformed theory at small radius, with explicit constants. That would be a theorem about a deformed theory; carrying it to $\mathbb{R}^4$ is the adiabatic-continuity conjecture.

Opening 3: three dimensions first

In three dimensions g^2 has the dimensions of energy, the theory is superrenormalizable, and the coupling at lattice scale is $g^2 a / \hbar c$, which vanishes as a power of a . The weak-side deficit of twelve orders of magnitude found in the small-field threshold note comes from the logarithmic running

of four dimensions, and in three dimensions the ultraviolet steps shrink geometrically, so the constant-chasing that failed in four dimensions may close. The one-function note reduces the problem to a lower bound $f(x) \geq f_- > 0$ on one function of one variable. A three-dimensional gap uniform in the spacing is open and would be a major result by itself.

Opening 4: the Planck-gap link

The small-volume mechanism of G07 and G08 is Simon's: along the flat valleys of the commutator potential the transverse modes are oscillators whose frequency grows with the distance along the valley, and their zero-point energy lifts the valley (Simon, Ann. Phys. **146**, 209, 1983, metadata). In its simplest form, for fixed x ,

$$\frac{\hat{p}_y^2}{2m} + \frac{1}{2}m\Omega(x)^2\hat{y}^2 \geq \frac{1}{2}\hbar\Omega(x),$$

an operator inequality that holds in every state. The torus-valley note finds the field-theory version term by term: the linear term $2|a|$ of the valley potential is the zero-point energy of the $k = 0$ charged modes.

The Planck-gap work sharpens what kind of floor this is. Its floors that rest on second moments of a prepared state fall to squeezing and to grid states, and the ones that survive rest on the commutator alone: the phase between Newton's polygon and the parabola, and the cost of disturbance. Simon's inequality is of the surviving kind, a spectral statement, so the valley lifting is the mass-gap instance of the lesson the Planck paper draws. **First step if opened:** hypothesis (H3) of the Feshbach note, the gap of the zero-mode Hamiltonian h_3 plus the even valley potential $U(a)$, proved by a Simon-type operator inequality with every constant explicit. That is the lower side of the small-volume corner S at weak coupling, and it stays within reach of the Planck-gap methods.

3. Consequence for STATE

The mass-gap track stays paused. This note records the four openings and the answer on groups: the gap is expected for every compact simple non-abelian group, and $SU(3)$ is the group that fixes the constants. The map's statement that the conjecture is T2' with T3 is too strong: T3, with a gap on an interval $(0, g_1)$ along the scaling curve, is what is needed, and T2' fails for $SU(N \geq 5)$ with the Wilson action. If one opening is taken up next, it is the fourth, which uses the Planck-gap methods directly; the first carries the most promise, and its first check is the conversion of Shen, Zhu and Zhu's threshold to $g^2 > 32$.