

Same collisions, different transport: an exactly soluble two-speed gas

Abstract

Can a collision rate and a velocity variance determine a diffusion coefficient? An equal-mass gas of point particles on a line gives an explicit counterexample. We compare two stationary preparations with the same mass, speed, density, energy density and mean collision rate. Independent Poisson positions and random velocity signs give diffusive tagged motion. Two randomly shifted lattices carrying opposite velocities give bounded, periodic tagged motion. The collision rule is identical in both cases: particles exchange velocities. Following straight trajectories through successive label exchanges reduces the comparison to exponential versus constant waiting times. The construction provides a classroom example of the statistical information hidden in a mean-free-time description of transport.

1 A collision rate does not specify a velocity history

A particle that repeatedly collides need not diffuse. In elementary transport arguments a typical speed and a mean free time suggest a diffusion scale, but the coefficient also depends on correlations between successive displacements. The purpose of this example is to expose that extra input in a gas whose microscopic dynamics can be drawn with straight lines.

Preparation-dependent tagged transport is established in single-file systems, where particles cannot pass one another. Leibovich and Barkai compare random and equally spaced Brownian preparations at the same density and find different long-time transport coefficients.[2] Cividini and Kundu discuss arbitrary initial positions and the distinction between fixed and ensemble-averaged preparations.[3] Here the microscopic motion is ballistic between elastic collisions. Correlating positions with velocity signs yields a particularly simple contrast: a tag that diffuses and a tag that returns to its starting point every period, even though their mean collision rates agree. The Poisson endpoint is the known two-speed case of the Jepsen gas.[1]

The contribution is an elementary comparison at matched collision frequency: the same elastic mechanics and one-particle kinetic scales support either ordinary diffusion or exact periodic cancellation. Sections 3 and 4 derive the two trajectories and their variances; Table 1 collects what is held fixed and what changes. The required tools are equal-mass elastic collisions, exponential waiting times and integration of a velocity covariance.

2 The common mechanics and the measured quantity

All particles have mass $m > 0$ and velocity $+u$ or $-u$, with $u > 0$. They occupy the infinite line at mean number density $\rho > 0$. An elastic collision between equal masses exchanges their velocities, conserving momentum and kinetic energy. Thus physical particles retain their spatial order. Equivalently, draw freely crossing trajectories $x = x_0 \pm ut$ in a position–time diagram, and exchange particle labels at each crossing. A tagged physical particle follows the successive segments selected by those label exchanges (Fig. 1).

Place a typical particle at the origin at time zero and follow its position $X(t)$. This selection is often called a Palm preparation. It fixes a tagged origin without changing the macroscopic




Figure 1: One collision in the label-exchange construction. A physical tag turns onto the other free line. Both preparations use this same rule.

density. The velocity process $V(t)$ is stationary: its joint law is unchanged by shifting all observation times. The displacement increments $X(t+s) - X(t)$ consequently have laws independent of t . The position process pinned at $X(0) = 0$ is not stationary. Reflection symmetry gives $\mathbb{E}[X(t)] = 0$. We use the ordinary diffusion coefficient

$$D = \lim_{t \rightarrow \infty} \frac{\text{Var } X(t)}{2t}, \quad (1)$$

when the limit exists. Its units are length squared per time.

Both gases are defined on the infinite line before taking this limit. They have finite energy density $\rho mu^2/2$, but infinite total reservoir energy. On any finite time interval $[0, T]$, bounded speed confines the tag to $[-uT, uT]$. Only free lines starting in $[-2uT, 2uT]$ can affect it during that interval. Local finiteness of either preparation therefore gives a well-defined path with finitely many collisions. A finite container followed to arbitrarily late times would be a different problem.

3 Poisson gaps give an exponential collision clock

For the first preparation, put independent Poisson particles of density ρ on the line and give each an independent, equally probable velocity sign. For a typical tag at zero the remaining gas has that same Poisson law. The two velocity streams are independent Poisson processes of density $\rho/2$. Free evolution translates each stream, preserving this homogeneous marked configuration law.

Condition on an initially right-moving tag. Let B_j be the initial positions of successive left-moving lines to its right, and $-A_j$ those of successive right-moving lines to its left, with $A_0 = B_0 = 0$. The tag follows ut , then $B_1 - ut$, then $-A_1 + ut$, then $B_2 - ut$, and so on. Parallel lines never intersect. At each reversal the next opposing line is the next unused line in the corresponding ordered sequence. The collision times consequently satisfy

$$t_{2j-1} = \frac{B_j + A_{j-1}}{2u}, \quad t_{2j} = \frac{B_j + A_j}{2u}. \quad (2)$$

Poisson gaps $B_j - B_{j-1}$ and $A_j - A_{j-1}$ are independent exponentials with rate $\rho/2$. Dividing them by the relative speed $2u$ shows that all tagged waiting times are independent exponentials with rate

$$\lambda = \rho u. \quad (3)$$

Reflection handles the other initial sign. The tag is therefore a stationary symmetric two-state velocity process, reversing at the times of a Poisson process. This is the standard telegraph velocity law, now obtained from spatial gaps rather than prescribed as a random clock.

If $N(s)$ counts reversals in a duration s , then $V(s) = V(0)(-1)^{N(s)}$. Summing the Poisson probabilities gives

$$C(s) = \mathbb{E}[V(0)V(s)] = u^2 e^{-2\lambda s}. \quad (4)$$

Stationarity and $X(t) = \int_0^t V(r) dr$ imply

$$\begin{aligned}\text{Var } X(t) &= 2 \int_0^t (t-s)C(s) ds \\ &= \frac{u^2}{\lambda} \left[t - \frac{1 - e^{-2\lambda t}}{2\lambda} \right].\end{aligned}\tag{5}$$

Thus

$$D_P = \frac{u^2}{2\lambda} = \frac{u}{2\rho} > 0.\tag{6}$$

The covariance and Markov interpretation agree with the exact two-speed result of Balakrishnan, Bena and Van den Broeck.[1] The tag here starts at one of the two gas velocities; a tag initially at rest has a different transient law.

4 Ordered streams give periodic cancellation

For the second preparation set $a = 2/\rho$. Right-moving free lines start at $a\mathbb{Z} + U_+$ and left-moving lines at $a\mathbb{Z} + U_-$, where the two phases are independent and uniform on $[0, a)$. Each stream has density $1/a = \rho/2$. Space translations and free time evolution translate the two phases modulo a , preserving their uniform joint law. This is another stationary marked gas with the same one-particle velocity probabilities and energy density. It changes both spatial gaps and their correlations with velocity signs: the union is not a Poisson gas with independent marks.

A typical tag has either velocity sign with probability $1/2$. Conditional on positive sign and placing it at zero, its own stream is $a\mathbb{Z}$; the opposite stream retains an independent uniform phase $U \in (0, a)$. Equation (2) now has

$$A_j = ja, \quad B_j = U + (j-1)a.\tag{7}$$

The first waiting time $R = U/(2u)$ is uniform on $(0, d)$, where

$$d = \frac{a}{2u} = \frac{1}{\rho u}.\tag{8}$$

Every subsequent waiting time is exactly d . The uniform initial sign and uniform residual time make the velocity a stationary square wave of period $2d$. It reverses indefinitely at mean rate $1/d = \rho u$, yet the tag travels as far to the right as to the left in each full period. Every realization returns to its initial position after $2d$.

The variance can be computed without invoking a long-time correlation integral. For $0 \leq s \leq d$, there is no reversal with probability $1 - s/d$, giving a displacement of magnitude us . Otherwise a reversal at residual time $r \in (0, s)$ gives displacement of magnitude $u|2r - s|$. Hence

$$\begin{aligned}\text{Var } X(s) &= u^2 \left[(1 - s/d)s^2 + \frac{1}{d} \int_0^s (2r - s)^2 dr \right] \\ &= u^2 \left(s^2 - \frac{2s^3}{3d} \right).\end{aligned}\tag{9}$$

For any $t > 0$, write $\delta = t - 2d\lfloor t/(2d) \rfloor$ and $s = \min(\delta, 2d - \delta)$. Full periods contribute zero displacement. Stationarity and cancellation over a full period equate the variance over $\delta > d$ to that over its complementary duration $2d - \delta$. Equation (9) therefore gives $\text{Var } X(t)$ with this folded s . Its maximum, at $s = d$, is $u^2 d^2/3$, so

$$0 \leq \text{Var } X(t) \leq \frac{1}{3\rho^2}, \quad D_O = 0.\tag{10}$$

There is no disorder-driven diffusion hidden at a longer timescale: the paths are exactly periodic. Simultaneous collisions elsewhere in the lattice are disjoint binary events. Coincident initial phases have probability zero.

5 What the comparison teaches

Table 1 collects the controlled quantities and the changed preparation. Equal values of the mean complete collision interval and velocity variance do not determine the covariance history that enters Eq. (5). In the ordered gas the initial global phases persist. The velocity process is stationary but not mixing, and positive and negative displacements cancel exactly. Inferring an effective independent collision clock from the mean collision rate would discard precisely the information responsible for Eq. (10).

Quantity	Poisson gas	Ordered streams
Mass; speed; density	$m; u; \rho$	$m; u; \rho$
Energy per length	$\rho mu^2/2$	$\rho mu^2/2$
Velocity variance	u^2	u^2
Mean collision rate	ρu	ρu
Joint preparation	Poisson gaps; independent signs	Two shifted lattices; sign fixed by stream
Complete waiting intervals	Independent exponential, mean d	Constant d
First residual interval	Exponential, mean d	Uniform on $(0, d)$
Tagged displacement	Diffusive at long times	Bounded and periodic
Diffusion coefficient	$u/(2\rho)$	0

Table 1: Same mechanics and mean collision frequency, different transport. Here $d = 1/(\rho u)$. The residual interval is measured from a typical observation time, which need not coincide with a collision.

The distinction between a complete waiting interval and its residual provides a useful teaching check. In the ordered gas the latter has mean $d/2$, although collisions occur at long-run rate $1/d$. Taking the reciprocal of the mean forward waiting time would give the wrong collision rate. In the Poisson gas memorylessness makes those two means equal.

Three short classroom questions follow directly from the construction: (i) derive Eq. (2) from a position–time diagram; (ii) explain why the collision rate counts reversals rather than the rate 2λ in the velocity covariance; and (iii) use a uniformly chosen observation phase to derive the ordered residual law and verify cancellation over a full period. Together these separate collision mechanics, spatial statistics and transport, using only elastic collisions, exponential waiting times and elementary integration.

References

- [1] V. Balakrishnan, I. Bena, and C. Van den Broeck, “Velocity correlations, diffusion, and stochasticity in a one-dimensional system,” *Phys. Rev. E* **65**, 031102 (2002), doi:10.1103/PhysRevE.65.031102.
- [2] N. Leibovich and E. Barkai, “The lasting effect of initial conditions on single file diffusion,” *Phys. Rev. E* **88**, 032107 (2013), doi:10.1103/PhysRevE.88.032107.
- [3] J. Cividini and A. Kundu, “Tagged particle in single-file diffusion with arbitrary initial conditions,” arXiv:1704.04017 (2017), doi:10.1088/1742-5468/aa75de.