

Rivero's consistency conjecture: quartic paths and central-force angles

Result, 2026-09-27. The quartic oscillator supplies three precise tests of the user's 1998 proposal. Its real-time path integral exists as a strong operator limit for every supplied action constant $h > 0$. Its halved critical-point functional already fails to recover the classical Dirac measure on a two-cell partition with three stationary paths (Theorem 1). Branch selection repairs the classical iterated limit; in Euclidean time there is also a positive, non-quadratic construction in which both iterated limits and every joint zero-resolution limit agree (Theorems 2–3).

For Newton's central-force polygon, angular momentum and equal areas survive every central kick. Requiring a separated angular phase to descend to the angle circle gives $L = nh$, or $\oint L d\theta = 2\pi nh$ (Theorem 4). This condition uses an additional coherent-state premise. For one fixed orbit it permits $h = |L|/n \rightarrow 0$. Hooke and Kepler radial actions give the corresponding Keller–Maslov conditions explicitly; even a fixed torus satisfying both conditions admits a sequence $h \downarrow 0$ (Theorem 5). These are consistency relations between a phase constant and orbit data. Selection of one universal positive value requires a further premise.

The tests and proofs below develop the question in the fifth-postulate note, §7, using the time-graded groupoid and the insertion laws, §§2–3. The constructions assemble established stationary phase, product formulas and semiclassical quantization.

1. Two regulators and three meanings of consistency

Use mass $m > 0$, duration $T > 0$, fixed endpoints, and a partition $\pi = (0 = t_0 < \dots < t_N = T)$. Write $\tau_j = t_{j+1} - t_j$ and $\varepsilon' = |\pi| = \max_j \tau_j$. The symmetric discrete action is

$$S_\pi(q) = \sum_{j=0}^{N-1} \left\{ \frac{m(q_{j+1} - q_j)^2}{2\tau_j} - \frac{\tau_j}{2} [V(q_j) + V(q_{j+1})] \right\}. \quad (1)$$

The action resolution ε and the constant h have units of action. Throughout this note h **denotes a reduced phase constant**: when identified with quantum mechanics, $h = \hbar$ and the ordinary Planck constant is $h_P = 2\pi h$. Time steps always use τ or ε' .

Rivero 1998, eqs. (1)–(10) (full-read, three pages) proposes a critical-point distribution, its halved oscillatory functional, a discretization with two regulators, and a controlled limit with arbitrary positive h . The control map in eq. (6) remains unspecified. The conjectural equivalence therefore needs both a normalization and a topology. The following definitions supply those for the tests here.

For $d = N - 1$ internal coordinates, choose a reference length $b > 0$ and put $z_j = q_j/b$, so $\partial_{z_j} S_\pi$ has action units. With a smooth compact cutoff χ , supported away from degenerate critical points, define a genuine regularization

$$D_{\varepsilon,\pi}(O) = \int \frac{\chi(z)O(z)}{(2\pi\varepsilon^2)^{d/2}} \exp \left[-\frac{|\nabla_z S_\pi(z)|^2}{2\varepsilon^2} \right] dz.$$

The change-of-variables formula at the finitely many critical points z_a gives

$$D_{0,\pi}(O) = \sum_a \frac{\chi(z_a)O(z_a)}{|\det S''_\pi(z_a)|}. \quad (2)$$

The coordinate convention and determinant weights are part of the definition. A normalized single-branch version is the point measure δ_{z_a} . Define separately the halved functional

$$A_{\varepsilon,\pi}(O) = (2\pi\varepsilon)^{-d/2} \int \chi(z)O(z)e^{iS_\pi(z)/\varepsilon} dz.$$

If $\chi = 1$ near all retained critical points, stationary phase gives

$$A_{\varepsilon,\pi}(O) = \sum_a \frac{O(z_a) e^{iS_\pi(z_a)/\varepsilon + i\pi\sigma_a/4}}{\sqrt{|\det S''_\pi(z_a)|}} + O_\pi(\varepsilon), \quad (3)$$

where σ_a is the Hessian signature. For one critical point, $|A(O)|^2 \rightarrow D_{0,\pi}(|O|^2)$. The complex amplitude itself needs its critical phase removed, or the ratio $A(O)/A(1)$. For several critical points, (3) retains cross terms. Also, a squared amplitude is quadratic in O , whereas (2) is linear in its test function. Equations (2)–(3) state precisely the limited sense in which halving recovers a Dirac measure.

The two proposed routes are now:

1. **Classical:** take $\varepsilon \downarrow 0$ at fixed π in (2), or in a branch-normalized (3), then refine the resulting discrete stationary paths. Convergence means weak convergence on continuous paths, or convergence against specified cylinder observables.
2. **Controlled at fixed h :** use free-kernel normalization on each cell and integrate $e^{iS_\pi/h}$ over the new vertices. Refine with $h > 0$ fixed. In real time, convergence means strong operator convergence on L^2 , allowing distributional coordinate kernels. A finite complex measure of bounded variation on all continuous paths is a stronger requirement and is not assumed here.

The second route implements Rivero’s eqs. (7)–(8). Calling it a joint limit of the *bare* regulators requires an explicit control map. For example, with $\eta = \varepsilon'/T_0$ dimensionless, a prescribed rescaling $S_\pi^{\text{bare}} = Z_S S_\pi$ has effective phase constant $h_{\text{eff}} = \varepsilon/Z_S$. Taking $Z_S = \eta$ and $\varepsilon = h\eta$ keeps $h_{\text{eff}} = h$. Simply substituting $\varepsilon = h\eta$ into $e^{iS_\pi/\varepsilon}$ changes the phase to $S_\pi/(h\eta)$; obtaining S_π/h requires the action/control transformation as well. This specifies the dimensional form of a possible control map; deriving that map remains an obligation.

Conjecture R (a sharpened version to test). For a specified action class, boundary data, blocking maps and normalizations, existence of a refinement-compatible classical critical-point distribution is equivalent to existence of its controlled oscillatory continuum functional; the classical limit of the latter recovers the former on a stated observable class. Every quantifier here matters. A stronger *selection conjecture* would also require these conditions to exclude all zero-action limits, or to select a common $h > 0$. The latter claim goes beyond existence.

There are three concrete theorem forms for “consistency”:

- **Classical elimination:** on a specified noncaustic branch, $S_{t+s}(x, y) = \text{stat}_z[S_t(x, z) + S_s(z, y)]$ for the exact endpoint action. A minimum replaces stationarity when the branch is a genuine global minimizing problem. For real-time quartic mechanics, the unrestricted action is unbounded below; the $(\min, +)$ law in the groupoid note needs this qualification. Euclidean action restores a minimizing problem.
- **Oscillatory composition:** $U_h(t + s) = U_h(t)U_h(s)$, with intermediate coordinates integrated. Eliminating a vertex by stationary phase produces its Hessian factor and higher corrections as well as its stationary action. Exact integration carries those amplitudes along.
- **Compatibility of limits:** after specifying a common observable functional $F_{h,\pi}$, an estimate $|F_{h,\pi}(O) - F_{h,0}(O)| \leq r(|\pi|, h)$ plus a controlled classical limit is sufficient along routes where $r \rightarrow 0$. Uniformity in h would justify interchanging limits. The summable-defect theorem of the refinement note supplies such a criterion when its norm hypotheses hold. Determinant weights in (2) also need compatible pushforwards; meshwise existence alone does not supply them.

The determinant issue has an exact local formula. Write an insertion as $S(z, y)$ and assume $S_y(z, y_*(z)) = 0$ defines one branch with invertible S_{yy} . Set $S_{\text{eff}}(z) = S(z, y_*(z))$. Integration of the two Euler–Lagrange delta factors gives, locally on this branch,

$$\int \delta(S_y(z, y)) \delta(S_z(z, y)) dy = \frac{\delta(\nabla S_{\text{eff}}(z))}{|\det S_{yy}(z, y_*(z))|}.$$

Indeed the first delta fixes $y = y_*(z)$ with the displayed Jacobian, and $\nabla S_{\text{eff}} = S_z$ there. At a joint critical point the same identity follows from the Schur complement $\det S'' = \det S_{yy} \det S''_{\text{eff}}$. Classical refinement therefore has an explicit weight to transport; stationary phase transports its square root, signature and higher corrections.

Thus even quadratic exactness presupposes a Gaussian normalization and appropriate topology; focal times require distributional kernels. The quartic test below separates the three theorem forms explicitly.

2. Quartic real time: a finite interference obstruction

Fix $V(q) = \lambda q^4$, $\lambda > 0$, so $F(q) = -4\lambda q^3$. Take endpoints zero, two cells of length $T/2$, and internal coordinate y . Work in the physical y coordinate; choosing $z = y/b$ multiplies the following functionals by common coordinate factors and leaves the failure of convergence unchanged. Equation (1) becomes

$$f(y) = \frac{2my^2}{T} - \frac{\lambda T}{2} y^4.$$

Theorem 1. A compact cutoff equal to one near the three critical points gives a finite $\delta(f')$, while the squared halved functional fails to converge as $\varepsilon \downarrow 0$.

Proof. The critical points and their data are

$$\begin{aligned} y_0 = 0, \quad y_{\pm} = \pm \sqrt{\frac{2m}{\lambda T^2}}, \quad f(y_{\pm}) = s_* := \frac{2m^2}{\lambda T^3}, \\ f''(0) = \frac{4m}{T}, \quad f''(y_{\pm}) = -\frac{8m}{T}. \end{aligned}$$

For $O = 1$ on the cutoff support, put $a = \sqrt{T/(4m)}$ and $b_1 = \sqrt{T/(8m)}$. Equation (3) reads

$$\begin{aligned} A_{\varepsilon} &= ae^{i\pi/4} + 2b_1 e^{is_*/\varepsilon - i\pi/4} + O(\varepsilon), \\ |A_{\varepsilon}|^2 &= \frac{3T}{4m} + \frac{T}{\sqrt{2}m} \sin(s_*/\varepsilon) + O(\varepsilon). \end{aligned} \tag{4}$$

Sequences with $s_*/\varepsilon = 2\pi k + \pi/2$ and $2\pi k + 3\pi/2$ give different limits. Meanwhile $\int \chi \delta(f') dy = T/(2m)$ by (2). Even averaging away the sine in (4) retains interference between y_+ and y_- , whose actions coincide. A separate diagonalization or branch label would be needed. $\square$

This is an obstruction to extending the halving identity to all critical paths by one overall normalization. It appears before a continuum limit or a choice of h . Retaining the pair of independent amplitudes also introduces off-diagonal terms; it cannot be identified with (2) without an additional prescription.

Theorem 2 (two existing limits for the quartic).

- (a) On every partition, select only a small neighborhood of the zero stationary path. Then $A_{\varepsilon, \pi}(O)/A_{\varepsilon, \pi}(1)$ converges to $O(0)$, and refining these branch measures gives the Dirac measure on the zero path.
- (b) For every $h > 0$, the symmetric time-sliced operators converge strongly on $L^2(\mathbb{R})$ to the unitary quartic evolution, independently of the sequence of partitions with mesh tending to zero:

$$A_h = -\frac{h^2}{2m} \frac{d^2}{dq^2}, \quad B = \lambda q^4, \quad H_h = A_h + B,$$

$$Q_{h,\tau} = e^{-i\tau B/(2h)} e^{-i\tau A_h/h} e^{-i\tau B/(2h)}, \quad Q_{h,\pi} \longrightarrow U_h(T) = e^{-iTH_h/h}. \quad (5)$$

Proof of (a). At zero the Hessian of (1) is the positive definite Dirichlet kinetic matrix: the quartic term has zero second derivative. The inverse function theorem isolates this branch at each partition. Equation (3) cancels its nonzero leading coefficient in the ratio. Its polygon is identically zero on every partition, proving the outer limit. This proves branch selection explicitly; its cutoff may depend on π .

Proof of (b). The positive polynomial Schrödinger operator is essentially self-adjoint on $C_c^\infty(\mathbb{R})$; see the continuous, bounded-below case recorded in Simon 1999, p. 2 (passage, Theorem 1.1 and following self-adjointness statement). Its domain is $D(A_h) \cap D(B)$. For completeness, the estimate ensuring this domain identification follows by integration by parts. With $a_h = h^2/(2m)$,

$$\|H_h f\|^2 = a_h^2 \|f''\|^2 + \lambda^2 \|q^4 f\|^2 + 2a_h \lambda \int q^4 |f'|^2 dq - 12a_h \lambda \int q^2 |f|^2 dq.$$

The pointwise inequality $12a_h \lambda q^2 \leq \lambda^2 q^8/2 + C_h$, with $C_h = 9a_h \lambda (6a_h/\lambda)^{1/3}$, absorbs the negative term. Closure and smooth cutoff approximation give the stated domain and its equivalent graph norm. Both factors in (5) are unitary, and on this domain

$$\tau^{-1}(Q_{h,\tau} - U_h(\tau))f \longrightarrow 0.$$

The difference quotients are uniformly bounded from the graph domain to L^2 . The orbit $\{U_h(s)f : 0 \leq s \leq T\}$ is compact in that domain, so convergence is uniform on the orbit. Telescoping an arbitrary partition therefore bounds the error by

$$T \sup_{\substack{0 < \tau \leq |\pi| \\ 0 \leq s \leq T}} \frac{\|(Q_{h,\tau} - U_h(\tau))U_h(s)f\|}{\tau} \longrightarrow 0.$$

Density extends convergence to L^2 . This is the product-formula argument of Teschl, Theorem 5.10, pp. 131–132 (passage), with the same telescoping proof for unequal cells. The kernel of $Q_{h,\tau}$ has phase (1) and normalization $(m/(2\pi i h \tau))^{1/2}$. Hence (5) is exactly the stated time slicing. $\square$

Composition and $\|U_h(T)\| = 1$ hold for every supplied $h > 0$. The proof gives convergence at fixed h , with graph estimates depending on h ; it makes no assertion of an unnormalized real-time amplitude limit as $h \downarrow 0$. Theorem 1 blocks the unrestricted classical-first halved prescription, while (a) gives a precise classical-first repair. An equivalence in Conjecture R must retain that distinction.

3. Quartic Euclidean time: a uniform two-limit theorem

A positive version gives a stronger existence test, with a direct uniform estimate. Keep the quartic potential and endpoints zero; change the action to

$$S_\pi^E = \sum_j \frac{m(q_{j+1} - q_j)^2}{2\tau_j} + W_\pi, \quad W_\pi = \frac{\lambda}{2} \sum_j \tau_j (q_j^4 + q_{j+1}^4).$$

Let $\nu_{h,\pi}$ be the normalized density proportional to $e^{-S_\pi^E/h}$, embedded as piecewise linear paths. Here h is the same action resolution called ε in the classical-first route.

Theorem 3. For every $h > 0$, arbitrary mesh refinement gives the quartic Feynman–Kac bridge ν_h on $C([0, T])$. Both iterated limits, and every joint limit $h \downarrow 0$, $|\pi| \downarrow 0$, give $\delta_{q \equiv 0}$. More precisely, for every $r > 0$ and every partition,

$$\nu_{h,\pi}(\|q\|_\infty > r) \leq 2 \exp \left\{ \frac{3\lambda h T^3}{16m^2} - \frac{2mr^2}{hT} \right\}. \quad (6)$$

Proof. Let B_h be the zero-endpoint Brownian bridge with covariance

$$\mathbb{E}[B_h(s)B_h(t)] = \frac{h}{m} \left(\min(s, t) - \frac{st}{T} \right).$$

Its values at the vertices have the normalized free kinetic density. Thus $\nu_{h,\pi}$ is their polygonal interpolation, tilted by $e^{-W_\pi(B_h)/h}/Z_{h,\pi}$. Continuity of bridge paths gives uniform convergence of the interpolation and convergence of the trapezoidal potential sum to $\lambda \int_0^T B_h^4 dt$. The weights lie between zero and one, and their expectation tends to a strictly positive value. Dominated convergence proves the fixed- h limit. Its identification with the heat kernel of H_h , including diffusion coefficient h/m , is the bounded-below Feynman–Kac formula (Simon, Theorem 1.1, passage). The unnormalized heat kernels obey exact convolution.

For uniformity in h , use $\mathbb{E}B_h(t)^4 = 3[ht(T-t)/(mT)]^2 \leq 3h^2T^2/(16m^2)$. The trapezoidal weights sum to T , so Jensen gives

$$Z_{h,\pi} \geq \exp[-\mathbb{E}W_\pi/h] \geq \exp[-3\lambda h T^3/(16m^2)].$$

The interpolated supremum is bounded by the bridge supremum. Reflection of Brownian motion at its first passage through r , followed by conditioning on the final value zero, gives $\mathbb{P}(\sup B_h > r) = e^{-2mr^2/(hT)}$; applying the same identity to $-B_h$ gives the two-sided bound with factor two. Multiplying by $Z_{h,\pi}^{-1}$ proves (6). It tends to zero uniformly in π as $h \downarrow 0$; the limiting bridge obeys the same bound. The unique Euclidean minimizing path is zero since every term of S^E is nonnegative and the endpoints vanish. This proves all the claimed orders of limits. $\square$

This theorem concerns the Euclidean positive measure, with its stated boundary data. Analytic continuation provides real-time operators through the self-adjoint H_h ; transferring probability estimates such as (6) to oscillatory real-time path measures would require another argument. Together, Theorems 1–3 give an explicit status for each limit: a failing unrestricted halved classical limit, an existing selected classical limit, an existing fixed- h real-time continuum limit, and commuting Euclidean limits with a bound. These consistency tests permit phase constants approaching zero.

4. Proposition I: the polygon and the angle circle

Now use the plane, a central potential $V(r)$, and scalar signed angular momentum $L = q \times p$. Work on a regular segment away from $r = 0$. A central kick changes p by $-aq$ for some scalar a , and a drift changes q by $\tau p/m$. Each separately preserves L . The oriented area of the drift triangle is exactly

$$\Delta\mathcal{A} = \frac{1}{2}q_j \times q_{j+1} = \frac{\tau_j L}{2m}. \quad (7)$$

This is the finite equal-area mechanism of Newton’s Proposition I. Symmetric kicks from (1) have the same property. Nonlinear forces generally change old endpoints when a finite cell is split; their consistent integrators converge on compact collision-free segments as the mesh vanishes. Equation (7) remains exact at each mesh. These swept areas serve the angular test here; the Galileo inertial–parabola defect in the Planck programme remains a separate observable.

In polar coordinates the kinetic numerator in (1) is

$$r_j^2 + r_{j+1}^2 - 2r_j r_{j+1} \cos(\theta_{j+1} - \theta_j).$$

The full discrete action and its exponential are single-valued under every change of angular representative $\theta_j \mapsto \theta_j + 2\pi k_j$, for every $\varepsilon > 0$. The angular discrete Euler–Lagrange equation conserves

$$L_j = \frac{mr_j r_{j+1}}{\tau_j} \sin(\theta_{j+1} - \theta_j).$$

Thus periodicity of the original path weight, together with the polygon law, places no arithmetic restriction on ε .

An extra requirement arises when a *stationary state* is assembled from a fixed angular-momentum branch. The separated Hamilton–Jacobi function on the angular cover is $S(r, \theta, t) = W_r(r; E, L) + L\theta - Et$.

Theorem 4 (angular descent). Assume the branch amplitude is a scalar, single-valued function of θ , and require its angular phase $e^{iL\theta/h}$ to descend from $\mathbb{R}$ to S^1 with trivial boundary holonomy. Then, necessarily and sufficiently,

$$\frac{L}{h} \in \mathbb{Z}, \quad \oint L d\theta = 2\pi n h. \quad (8)$$

Proof. Translation by 2π multiplies the phase by $e^{2\pi i L/h}$. Its equality to one is (8). For lifted polygon increments the product is $\prod_j e^{iL\Delta\theta_j/h} = e^{iL\sum_j \Delta\theta_j/h}$; inserting a vertex leaves it unchanged. Descent through a full turn then imposes exactly the same condition, independently of the mesh. $\square$

This is also the exact angular spectrum of $-i\hbar\partial_\theta$ on periodic scalar wavefunctions. Allowing a flat boundary phase $\psi(\theta + 2\pi) = e^{2\pi i\alpha}\psi(\theta)$ changes the relation to $L/h \in \mathbb{Z} + \alpha$. The trivial scalar choice is an explicit premise; neither central kicks nor angle periodicity chooses a state space. Distinct winding paths remain distinct integration histories, even when their endpoints agree.

For a fixed $\ell = |L| > 0$, (8) permits precisely $h = \ell/n$ with $n = 1, 2, \dots$, whose infimum is zero. Holding the winding number n fixed determines h from that orbit. A bound $n \leq N_*$ would produce the conditional floor $h \geq \ell/N_*$, with both orbit scale and bound supplied. A common h for several prescribed orbits requires their angular actions to be integer multiples of it; incommensurate actions cannot all satisfy that exact-state premise.

5. Hooke and Kepler: radial actions and Maslov phases

For regular noncircular bound tori with $\ell = |L| > 0$, define

$$I_\theta = \ell, \quad I_r = \frac{1}{\pi} \int_{r_-}^{r_+} \sqrt{2m(E - V(r)) - \ell^2/r^2} dr.$$

The angular cycle has no turning point. The radial cycle has two simple turning points. Supplying the usual WKB transport and its caustic phase gives the Keller–Maslov rule

$$\exp \left[\frac{i}{h} \oint p \cdot dq - \frac{i\pi\mu}{2} \right] = 1, \quad I = h(n + \mu/4). \quad (9)$$

Here $\mu_\theta = 0$, $\mu_r = 2$. Each simple turning point contributes $-\pi/2$ to the transported cycle phase, yielding $I_r = h(n_r + 1/2)$ and $I_\theta = h|n_\theta|$. This is semiclassical state consistency, with WKB transport supplied; the angular rule is exact on the stated scalar state space. Keller 1958 (metadata, *Annals of Physics* 4, 180–188) is the established source; Keller’s later account (abstract) describes the integer and half-integer conditions from WKB. We use (9) conditionally and derive the following mechanical actions.

Hooke, $V = m\omega^2 r^2/2$, $\omega > 0$ (**Proposition X**). The radial period is $T_r = \pi/\omega$, since r^2 oscillates at 2ω . Differentiating the action integral gives $\partial I_r / \partial E = T_r / (2\pi) = 1/(2\omega)$. The circular orbit has $E_c = \omega\ell$ and $I_r = 0$, hence

$$I_r = \frac{E}{2\omega} - \frac{\ell}{2}, \quad E = \omega(2I_r + \ell). \quad (10)$$

Equation (9) consequently gives the EBK values

$$E = h\omega(2n_r + |n_\theta| + 1).$$

For a circle $\ell = m\omega R^2$, taking $R \downarrow 0$ produces arbitrarily small positive classical angular actions at fixed m, ω . The radial turning-point argument applies to noncircular tori; circular orbits follow as the mechanical limit of (10), with a degenerate radial cycle requiring its own wave analysis.

Kepler, $V = -k/r$, $k > 0$. For $E < 0$ and $0 < \ell < k\sqrt{m/(-2E)}$, the ellipse has two distinct positive radial turning points. Its semimajor axis is $a = -k/(2E)$ and its period is $T_r = 2\pi\sqrt{ma^3/k}$. Thus $\partial I_r/\partial E = k\sqrt{m}/(-2E)^{3/2}$. At the circle $E_c = -mk^2/(2\ell^2)$, $I_r = 0$, giving

$$I_r = k\sqrt{\frac{m}{-2E}} - \ell, \quad E = -\frac{mk^2}{2(I_r + \ell)^2}. \quad (11)$$

The planar EBK rule yields

$$E = -\frac{mk^2}{2h^2(n_r + |n_\theta| + 1/2)^2}.$$

This is the planar action calculation; the three-dimensional Coulomb problem has additional angular variables. The angular law alone fixes ℓ/h and leaves I_r free. Classical circles satisfy $\ell = \sqrt{mkR}$, so $R \downarrow 0$ also gives angular infimum zero at fixed m, k , through regular circles of individually positive radius.

Theorem 5 (even both cycle conditions allow $h \downarrow 0$). Fix a Hooke or Kepler noncircular torus satisfying (9) at some $h_0 > 0$, with integers $n_r \geq 0$, $|n_\theta| \geq 1$. The *same* actions satisfy (9) at every

$$h_j = \frac{h_0}{2j+1}, \quad n_{r,j} = (2j+1)n_r + j, \quad n_{\theta,j} = (2j+1)n_\theta, \quad j = 0, 1, \dots \quad (12)$$

Proof. Substitution gives $h_j(n_{r,j} + 1/2) = h_0(n_r + 1/2)$ and $h_j|n_{\theta,j}| = h_0|n_\theta|$. Equations (10)–(11) then preserve the energy and angular momentum. $\square$

Some fixed tori admit no h satisfying both exact EBK relations, because their action ratio fails the required integer/half-integer relation. Whenever one value exists, (12) excludes a positive lower bound from these relations alone. Asymptotic WKB errors carry their usual separate status; (12) is an exact statement about the proposed cycle conditions.

Finally, the action in a closed spatial circuit must be identified correctly. Write $W = \oint p \cdot dq = 2\pi J_{\text{orb}}$, whereas $S_T = W - ET$. For Hooke, $J_{\text{orb}} = 2I_r + \ell = E/\omega$ and $T = 2\pi/\omega$, so $S_T = 0$ on every ellipse. Imposing $e^{iS_T/h} = 1$ there gives no restriction whatever. For Kepler, $J_{\text{orb}} = I_r + \ell$ and $ET = -\pi J_{\text{orb}}$, so $S_T = 3\pi J_{\text{orb}}$. Equal configuration after a dynamical period allows a stationary state to acquire its temporal phase $e^{-iET/h}$. The spatial cycle condition (9) therefore uses W and the caustic phase, with time held separate. Identifying it with the bare full-period halved action would change the condition.

6. Relation to the orbit bounds and Kepler threshold

The closed-orbit force-action note proves $J_{\text{orb}} \geq m^2 v_*^3 / F_{\text{max}}$ in Newtonian mechanics under a uniform speed floor and force ceiling. For eccentric orbits this is the total orbit action above; it differs from $I_\theta = \ell$. The excitation and force hypotheses supply that bound. Combining it with an integer rule would still require an upper bound on the integer to bound h from below.

On a circle $J_{\text{orb}} = \ell$. The bound-orbit observable note gives $2\sqrt{\det \Sigma} = \ell$ for its uniform-phase preparation. Equation (8) would turn this into $2\sqrt{\det \Sigma} = |n_\theta| h$ if that classical estimator is used to label an admitted angular action. Identifying the classical ensemble with a quantum state would need an additional preparation argument.

The relativistic Kepler note has a different, explicit mechanism: in the external singular potential $-k/r$, regular bound motion requires $\ell > k/c$. Combining this classical admissibility condition with angular descent gives

$$|n_\theta| h > k/c.$$

For fixed h it excludes low angular integers. For a fixed integer it requires $h > k/(c|n_\theta|)$. With unbounded integers it permits arbitrarily small h ; for a fixed allowed orbit use $h = \ell/n$. The threshold remains coupling dependent and closes in the Newtonian limit $c \rightarrow \infty$. A softened core has angular infimum zero already at fixed m, k, c , as proved in that note. Thus the singular-core threshold, the orbital action estimator, and the angular phase lattice each retain their distinct hypotheses.

7. Consequence for STATE

Item 2 gains a tested form of the 1998 conjecture: quartic interference requires a branch or diagonal prescription for critical-point recovery; fixed- h real-time refinement exists for all $h > 0$, and a Euclidean quartic test has commuting zero-resolution limits. Central-force angular descent gives an orbit-dependent integer relation whose allowed phase constants accumulate at zero, even with both EBK cycles imposed. The constructive next obligation is a physically justified control and observable prescription that addresses the interference in (4), together with a scale-selection premise surviving the sequences (8) and (12). These results settle the stated tests; the general controlled equivalence in Conjecture R retains those explicit open obligations.