

Planck Area from Half-Density Normalization (Draft)

Abstract

Half-densities are the natural “coordinate-free integrands” for composing kernels without choosing a background measure. But choosing a *universal* convention for turning half-density objects into dimensionless numerical amplitudes introduces a length ^{$d/2$} scale. In $d = 4$, this is an *area*. This note sharpens the hypothesis ladder needed for the claim “half-density normalization selects a universal area scale”, and isolates a simple dimension-matching condition under which the Planck area appears without fractional powers of couplings. A gravitational anchor based on a minimal-areal-speed principle is recorded as a separate heuristic thread [RiveroAreal] [RiveroSimple].

1. Purpose and Status

This is a dependent follow-up to `paper/main.md`. It is not yet a finished paper; its goal is to isolate one technical point that is only implicit in the main manuscript: the role of half-densities (and their scaling) in making composition laws coordinate-invariant *and* dimensionally well-defined.

Claims below are labeled as **Proposition** (math-precise under hypotheses) or **Heuristic** (programmatically bridge).

2. Half-Densities and Composition Kernels

Let M be a d -dimensional manifold. A (positive) density is a section of $|\Lambda^d T^* M|$, and a half-density is a section of $|\Lambda^{d/2} T^* M|^{1/2}$.

The key operational point is: when a kernel is a half-density in its integration variable, composition of kernels does not depend on an arbitrary choice of coordinate measure.

Heuristic H1.1 (Why half-densities). If $K_1(x, y)$ and $K_2(y, z)$ are chosen so that their product in the intermediate variable y is a density, then $\int_M K_1(x, y) K_2(y, z)$ is coordinate-invariant without fixing a preferred dy . This matches the structural role of kernel composition used in `paper/main.md` (Section 6).

Derivation D1.1 (Coordinate invariance of half-density pairing and composition). In a local chart $y = (y^1, \dots, y^d)$, write a half-density as $\psi(y) = \varphi(y) |dy|^{1/2}$. Under a change of variables $y = y(y')$, one has $|dy|^{1/2} = |\det(\partial y / \partial y')|^{1/2} |dy'|^{1/2}$, so the coefficient transforms as $\varphi'(y') = \varphi(y(y')) |\det(\partial y / \partial y')|^{1/2}$.

Hence the product of two half-densities is a density: $\psi_1 \psi_2 = (\varphi_1 \varphi_2) |dy|$, and its integral is chart-independent: $\int_M \psi_1 \psi_2$ is well-defined without choosing a

background measure beyond the density bundle itself.

Kernel composition is the same mechanism: if $K_1(x, y)$ and $K_2(y, z)$ are half-densities in y , then $K_1 K_2$ is a density in y and $\int_M K_1 K_2$ is coordinate invariant.

3. Dimensional Analysis: Normalizing a Half-Density Requires a Scale

A density on M carries the units of length^d once physical units are assigned to coordinates. A half-density therefore carries units $\text{length}^{d/2}$.

Proposition P1.1 (No canonical "half-density = function" identification). There is no canonical identification of a half-density $\psi \in |\Lambda^{dT^*M}|^{1/2}$ with an ordinary scalar function f on M . Choosing such an identification is equivalent to choosing a nowhere-vanishing reference half-density σ_* (equivalently a positive density $\rho_* = \sigma_*^2$) and writing $\psi = f \sigma_*$.

Derivation D1.2 (Dilation makes the $\text{length}^{d/2}$ weight explicit). On \mathbb{R}^d , consider a dilation $y \mapsto y' = ay$ with $a > 0$. Then $|dy'| = a^d |dy|$, so $|dy'|^{1/2} = a^{d/2} |dy|^{1/2}$. Thus even in flat space, half-densities carry an inherent $\text{length}^{d/2}$ scaling weight.

Proposition P1.2 (Universal *dimensionless* amplitudes force a $\text{length}^{d/2}$ constant). If one imposes the extra requirement that the scalar representative f in $\psi = f \sigma_*$ be dimensionless in physical units, then the reference half-density σ_* must carry all of the $\text{length}^{d/2}$ dimension. In particular, a *constant* (field-independent) choice of σ_* is equivalent to choosing a universal $\text{length}^{d/2}$ scale.

In $d = 4$, this universal $\text{length}^{d/2}$ scale is a universal *area* scale.

Heuristic H1.2 (Reciprocity claim). Half-densities alone do not force a particular scale: the forced fact is that converting half-density objects into scalar numerical amplitudes requires extra structure (a reference half-density). The “universal area scale” claim begins only after adding two further hypotheses: 1. the reference σ_* is taken to be *constant* (no dependence on background metric/fields), and 2. the constant is required to be fixed by universal constants/couplings of the theory.

Under these hypotheses, $d = 4$ is the unique dimension in which the needed $\text{length}^{d/2}$ constant can be supplied by the gravitational coupling without fractional powers (Derivation D1.3).

Derivation D1.3 (Dimension match: why $(d=4)$ is singled out by gravity). In d spacetime dimensions, the Einstein–Hilbert action $\frac{1}{16\pi G_d} \int d^d x \sqrt{|g|} R$ shows that (in $c = \hbar = 1$ units) Newton’s constant has dimension $[G_d] = \text{length}^{d-2}$. If the universal half-density normalization constant is required to be built from G_d without fractional powers, then its dimension must match $\text{length}^{d/2}$ with exponent 1, i.e. $\text{length}^{d/2} = \text{length}^{d-2}$, which holds

if and only if $d = 4$. In that case G_4 itself has dimension of area, and the corresponding area scale is the Planck area $L_P^2 \sim \hbar G_4 / c^3$.

4. Stationary Phase Produces Half-Density Prefactors (Short-Time Kernel)

The main manuscript uses stationary phase to explain why classical extremals dominate refinement limits. Here we add the complementary kernel-level fact: stationary phase does not only pick the extremal; it also produces a determinant prefactor that transforms as a half-density, i.e. the object needed for coordinate-free kernel composition.

Derivation D1.4 (Van Vleck prefactor is a bi-half-density). Let $S_{\text{cl}}(x, z; t)$ be the classical action as a function of endpoints and time, treated as a generating function. The standard short-time/stationary-phase approximation to the propagator has the form

$$K(x, z; t) \approx \frac{1}{(2\pi i \hbar)^{d/2}} \left| \det \left(-\frac{\partial^2 S_{\text{cl}}}{\partial x \partial z} \right) \right|^{1/2} \exp \left(\frac{i}{\hbar} S_{\text{cl}}(x, z; t) \right).$$

Under a change of coordinates $x = x(x')$, $z = z(z')$, the mixed Hessian transforms by the chain rule, and its determinant acquires Jacobian factors:

$$\det \left(-\frac{\partial^2 S_{\text{cl}}}{\partial x' \partial z'} \right) = \det \left(\frac{\partial x}{\partial x'} \right) \det \left(\frac{\partial z}{\partial z'} \right) \det \left(-\frac{\partial^2 S_{\text{cl}}}{\partial x \partial z} \right).$$

Taking square roots shows that the prefactor transforms with $|\det(\partial x / \partial x')|^{1/2} |\det(\partial z / \partial z')|^{1/2}$, i.e. exactly as a half-density factor at each endpoint. Thus the stationary-phase prefactor is naturally interpreted as making K a half-density in each variable, so that kernel composition does not depend on a background measure choice.

Heuristic H1.4 (Where Planck area can enter, minimally). Derivation D1.3 isolates one minimal route by which a Planck-scale quantity can enter: if the theory supplies a single universal coupling with dimension of length (Newton's constant) and one demands that the half-density normalization constant be built from that coupling *without fractional powers*, then $d = 4$ is singled out and the resulting constant has the dimension of an area, naturally identified with the Planck area $L_P^2 \sim \hbar G_4 / c^3$.

5. A Gravitational Anchor: Minimal Areal Speed and the $D = 4$ Cancellation

Rivero's "Planck areal speed" observation gives a concrete route by which Planck-scale discreteness reappears at Compton scales in inverse-square gravity [RiveroAreal] [RiveroSimple].

Heuristic H1.3 (Areal-speed selection). In $3 + 1$ Newtonian gravity (inverse-square), imposing a discrete areal-speed/area-time condition at a Planck scale can yield characteristic radii proportional to a reduced Compton length, with Newton’s constant canceling when expressed in Planck units. This is a non-trivial indication that “a universal area scale” can be operationally meaningful at low energies in $D = 4$.

Derivation D1.5 (Inverse-square circular orbit + Planck areal speed \rightarrow Compton radius). For a circular orbit under an inverse-square central force $F(r) = K/r^2$ (with coupling $K > 0$), the centripetal balance is $mv^2/r = K/r^2$. The areal speed is $\dot{A} = \frac{1}{2}rv$, so $v = 2\dot{A}/r$. Substituting into the force balance gives

$$m \left(\frac{2\dot{A}}{r} \right)^2 = \frac{K}{r} \implies r = \frac{4m \dot{A}^2}{K}.$$

For Newtonian gravity between a source mass M and test mass m , $K = GMm$, hence

$$r = \frac{4\dot{A}^2}{GM},$$

independent of the test mass m . If one now imposes $\dot{A} = k \dot{A}_P$, where Rivero’s Planck areal speed is $\dot{A}_P = cL_P$ [RiveroAreal], then using $L_P^2 = G\hbar/c^3$ yields

$$r = \frac{4k^2(cL_P)^2}{GM} = \frac{4k^2(G\hbar/c)}{GM} = 4k^2 \frac{\hbar}{cM}.$$

Thus r becomes a multiple of the reduced Compton length $L_M = \hbar/(cM)$, with Newton’s constant canceled out. In particular, $k = \frac{1}{2}$ gives $r = L_M$. This is the “Planck area per Planck time \Rightarrow Compton scale” cancellation highlighted in [RiveroAreal] and summarized in [RiveroSimple].

6. Interface with the Main Paper

The main manuscript argues that: 1. classical dynamics are recovered from quantum composition by stationary-phase concentration, and 2. refinement across scales forces RG-style consistency conditions when naive limits diverge.

This draft adds a complementary ingredient: the kernel side is most naturally formulated in half-density language, and stationary phase produces the bi-half-density prefactor directly. A universal convention for turning those half-densities into scalar amplitudes then requires a length ^{$d/2$} scale; in $d = 4$ this is an area scale.

7. Open Problems (Needed for a Real Paper)

1. Make the half-density normalization argument precise for a concrete groupoid or kernel model (tangent-groupoid or short-time propagator model).
2. Show how the area scale enters stationary-phase prefactors and how this interacts with RG scaling.
3. General-dimension analysis: clarify what replaces “area” in odd dimensions and whether a universal normalization is still defensible.
4. Identify minimal hypotheses under which “need of half-density scale \Rightarrow Planck area” is more than dimensional bookkeeping.