

A Connes-Style Translation of the $\mathbb{CP}^2$ /Aloff–Wallach Kaluza–Klein Programme

KK Orchestra working manuscript

June 1, 2026

Abstract

This manuscript translates the current $\mathbb{CP}^2$ /Aloff–Wallach Kaluza–Klein programme into Connes-style noncommutative geometry. The translation uses spectral triples, real structures, inner fluctuations, boundary triples, finite wall modules, and spectral-action language to separate five layers: compact geometry, endpoint diagnostics, the D6 branch interface, the orthodox $\mathfrak{m}_3$ custodial Higgs block, and the open retained-response source problem. The main conclusion is deliberately conservative. The Connes vocabulary clarifies the missing object: a source-derived quadratic form on the retained anti-graph mismatch. It also keeps the established electroweak result $\rho = 1$ in the single $\mathfrak{m}_3$ branch separate from any future De Vries-style spectral refinement.

Contents

1	Reading Guide	2
2	Local Corpus Map	3
3	Legacy Discussion Coverage	3
4	Source-Route Corpus Map	5
5	Source Library Map	6
6	Package Atlas	7
7	Mathematical Expansion Convention	8
7.1	Retained Mismatch Source	8
7.2	Schur, Boundary, and Determinant Expansions	9
7.3	Endpoint, Gravity, and Tower Expansions	9
8	Connes Anchors	10
8.1	Basic Spectral Data	10
8.2	Source Dimension and Probe Dimension	11
8.3	KK, QFT, and Connes Layer Parallel	11
9	The One-Line Source Problem	12

10	Geometry as Families of Spectral Triples	13
10.1	Compact Metrics	13
10.2	Spectral Action and Einstein-Hilbert Reduction	14
11	D1/D2/D3 and D5/D6/D7	14
11.1	Low-Dimensional Prototype	14
11.2	CP2-Suspended Carriers	14
12	Endpoint Filters	15
13	The D6 Branch Interface	16
13.1	Three Objects	16
13.2	Link Monodromy and Bosonic Boundary Data	16
14	The m_3 Higgs Module	17
14.1	Symmetric Branch	17
14.2	Broken Branch	17
14.3	Custodial Condition	17
15	Finite Wall Modules and Relative Sources	18
15.1	Finite Wall Module	18
15.2	Relative Finite Source	18
15.3	Relative Determinant Source	18
15.4	Auxiliary Wall Sources	19
16	Gravity, Boundary, and Yang–Mills Terms	19
16.1	Ordinary Gravity	19
16.2	Ordinary Gauge-Higgs Sector	19
17	Objections and Consistency Checks	19
17.1	Objections	19
17.2	Consistency Checks	20
18	Additional Translation Frontiers	20
18.1	Tower Criterion for an Emergent Dimension	20
18.2	S^5 , Spin(6), and Electromagnetism	20
18.3	Noncommutative and Division-Algebra Repairs	21
19	Possible Completions	23
20	Conclusion	24

1 Reading Guide

The document has two intended readers.

- For a physicist, the translation explains which parts of the KK model are ordinary gauge, Higgs, gravity, boundary, or defect physics. It also records where a signed retained-response Hessian would need a genuine source action.

- For a geometer, the translation identifies the algebras, Hilbert modules, Dirac operators, real structures, and boundary data attached to the D5, D6, and D7 pieces.

2 Local Corpus Map

The current produced corpus has the following NCG reading:

NCG_KK_QFT_CONNES_DICTIONARY.md	expandable KK/QFT/Connes reference dictionary,
M3_DOUBLET_CUSTODIAL_Lecture.tex	custodial $\mathfrak{m}_3$ module and $\rho = 1$,
CUSTODIAL_3D_TO_7D_EXTENSION.md	Hopf-to-Aloff-Wallach custodial dictionary,
KK_HIGGSING_CUSTODIAL_MODEL.md	gauge-Higgs inner-fluctuation model,
KK_HIGGS_POTENTIALS_1D3D_5D7D.md	spectral-action and reduced-potential ledger,
GRAVITY_ACTION_D765_D321.md	metric-family and heat-coefficient control,
TRANSITION_1D3D_5D7D.md	family-of-triples transition dictionary,
D6_FERMION_BRIDGE.md	Spin ^c and branch-interface constraint sketch,
D6_DIRAC_BRANCH_DEFECT_SPECTRAL_FLOW.md	singular, APS, eta, and spectral-flow data.

These files supply local evidence, filters, and source bookkeeping. None of them replaces the retained boundary-source calculation for A_B on h .

3 Legacy Discussion Coverage

The wider project discussion is assigned to current NCG homes in [20]. The table below records the manuscript-level reading.

Discussion family	NCG home	Status
Hilo and KK source synthesis	Source narrative and corpus control through source and produced-output crosswalks.	Source-status labels required before use.
Custodial and $\mathbf{m}_3$	Real or quaternionic doublet module, isotropic scalar metric, and ordinary Higgs mass matrix.	Orthodox $\rho = 1$ filter.
Gauge-Higgs and scalar potentials	Inner fluctuations, compact Dirac-family moduli, spectral-action potentials, and branch Hessians.	Ordinary Higgs and branch-selection layer.
Transition and probe dimension	Families of triples, compressed sector triples, and phase dictionaries.	Phase and filtration layer.
Witten-style tower	Family $(\mathcal{A}_\epsilon, \mathcal{H}_\epsilon, D_\epsilon)$ with labelled eigenfamily and source-computed gap.	Dimension-growth criterion.
D5 endpoint and electroweak charge	Product endpoint triple, lens quotient module, flat characters, Q_{EM} , and weak-inertia rows.	Endpoint and charge-period filters.
D6 branch corpus	Boundary/singular triples, Schur packages, determinant packages, auxiliary modules, finite wall modules, and twisted-line modules.	Source-package search layer.
Boundary, Calderon, and DtN work	Boundary triple, Weyl function, Calderon projector, and finite-part trace.	Domain and spectral data.
De Vries and golden-rule search	Neutral multiplier μ_r , named source operator, retained package $P_B, A_B, m_B, \gamma_B, \sigma_r$.	Target-for-derivation and obstruction ledger.
Normalization and response metric	Response trace on F_B , map scale γ_B , Schur weight w_B , and residual quotient.	Trace-normalization package.
Noncommutative, fuzzy, and CSDR repairs	Finite/equivariant triples, fuzzy flag modules, projective A_θ modules, Morita and trace-selection data.	Exotic repair candidates.
Spin ^c , fermion bridge, and division algebra	Spin ^c Hilbert modules, finite Hilbert-space constraints, real/quaternionic structures, and defect carriers.	Bosonic constraints and carrier language.
Relative gain, determinant, and finite trace	Relative source functional, finite trace module, determinant second variation, and counterterm rule.	Conditional source routes.

The residual mathematical target across the table is the same retained source package:

$$P_B : F_B \rightarrow E_B, \quad K_B = P_B^\dagger A_B P_B, \quad m_B, \quad \gamma_B, \quad \sigma_r.$$

A filename-level residual audit is recorded in [21]. Notes that were only implicit in the family tables receive a Connes package class, an expansion anchor, and an allowed-use label there.

Repository artifacts outside the current source-note-output corpus are covered in [22]. Top-level ledgers, archives, task prompts, calculation artifacts, settings, and build auxiliaries enter as governance, source narrative, historical evidence, prompt material, calculation artifact, verification artifact, or no mathematical role. Any mathematical reuse from those files still passes through a complete expansion block or a named current note.

The original claims C1–C7 and task prompts T1–T6/R4/R5 are translated in [23]. That ledger assigns them to the current package classes: branch/groupoid spectral data for C1, homogeneous compact triples for C2–C3 and T2, Spin^c modules for C4/T1, gauge-inertia traces for C5/T3/R4, the $\mathbf{m}_3$ custodial module for T4, source-library obstructions for T6, endpoint charge-factor diagnostics for R5, and the retained package

$$E_B, \quad F_B, \quad P_B, \quad A_B, \quad m_B, \quad \gamma_B, \quad \sigma_r$$

for C7/T5.

The wastebook correction history is translated in [24]. Its confirmed geometry, spin and Spin^c corrections, gauge-inertia caveats, corrected literature entries, weak-angle alternatives, flipped colour/electroweak identification, and rejected spectral shorthand are assigned to package homes before reuse. In particular, the old shorthand is replaced by the neutral retained-source notation

$$O_r^\dagger O_r = \mu_r \text{Id} + \Delta_r, \quad P_B^\dagger A_B P_B|_r = \gamma_B^2 \mu_r \text{Id} + \sigma_r.$$

The archived T/R verification reports are translated in [25]. They supply historical evidence for Spin^c and index filters, compact metric families, gauge inertias, custodial mass matrices, source-origin obstructions, no-go evidence, weak-angle comparisons, and flipped endpoint diagnostics. Current derivations still require the relevant operator, action, trace, and residuals in a live package note or in this manuscript.

Calculation artifacts under `calc/` are translated in [26]. Topology scripts are assigned to Spin^c and endpoint-period packages, metric scripts to compact spectral triples, weak-angle scripts and arrays to gauge-inertia packages, endpoint comparison scripts to representation diagnostics, custodial scripts to the $\mathfrak{m}_3$ ordinary Higgs Hessian, and cache files to no mathematical role. Any script-derived formula used here has to pass through a displayed expansion or a cited archive report before it enters an argument.

Produced-output premise remediation is tracked in [27]. Older boundary and vertical-sector formulas are read through named operator multipliers

$$P_{\text{bd}}^\dagger P_{\text{bd}}|_{E_j} = x_0 \mu_j^{\text{bd}} \text{Id}_{E_j} + \sigma_j^{\text{bd}}, \quad M_{\text{vert},r} = - \sum_a \rho_r(T_a)^2 = \mu_r^{\text{vert}} \text{Id}_{V_r}.$$

These are direct boundary or representation readouts. Retained-source use still requires one source package for

$$P_B, \quad A_B, \quad m_B, \quad \gamma_B, \quad \sigma_r.$$

The note-layer shorthand remediation index [29] applies the same rule to older research notes that still contain representation or boundary multiplier abbreviations. Role-level produced-output anchors in [28] give each current `out/` deliverable a Connes package form before reuse: manuscript, dictionary, custodial module, gauge-Higgs Hessian, potential, gravity, transition, Spin^c , or singular Dirac data.

4 Source-Route Corpus Map

The older D6 source-admission and deletion notes are read through the NCG package classes in [37]. The current convention is

$$P_B : F_B \rightarrow E_B, \quad h = T_B x - (q - P_B p), \quad K_B = P_B^\dagger A_B P_B.$$

Older notes that write $q - P_{\text{old}}^\dagger p$ translate by

$$P_B = P_{\text{old}}^\dagger.$$

Source-route family	NCG package	Use in the programme
Folded collar and compliance	Boundary Schur package with relaxed collar module Z_B , Weyl scale Λ_B , and Schur weight W_B .	Candidate retained source after trace and response data agree.
Boundary triple and Calderon	$D_{\min}, D_{\max}, \Gamma_0, \Gamma_1$, Weyl function, Calderon projector, finite-part trace.	Domain and spectral package; coefficient source only after K_B is derived.
Local B_4 routes	Local module with variables u , $R_{\text{def}}h$, and Hessian blocks $K_{\text{loc}}, J_{\text{loc}}, E_{\text{loc}}$.	Conditional Schur source after vacuum, Hessian, coupling, and trace are fixed.
Determinant and gluing	h -dependent operator family $L_B(h)$, determinant, counterterm, and finite trace.	Candidate after sign, subtraction, zero modes, and counterterms are source data.
Influence, constraint, and transmission	Auxiliary module Z_B , operator K_Z , coupling J_Z , contour or constraint sign.	Candidate after physical sign and response trace are derived.
Finite wall and smoothing	Finite or layered spectral module with finite trace and endpoint representation.	Candidate after blocks, traces, and limits are fixed.
Twisted angular capacity	Flat line L_δ on E_B , normal-disk capacity, and twisted response module.	Candidate after the bosonic line on h and normal-gradient coefficient are derived.
Gravity, cone, and ordinary Higgs routes	Compact metric, spectral-action, conical, Yang–Mills, or scalar Hessian data.	Stability, electroweak, and branch-selection filters.
Endpoint and representation routes	Product, quotient, Hopf/lens, finite-centre, and flat-character endpoint modules.	Charge, period, global-form, and roundness filters.

The common source-readout expansion is

$$K_B|_r = \gamma_B^2 \mu_r \text{Id} + \sigma_r, \quad \|p_r\|_{\text{kin}}^2 = m_B \|p_r\|^2,$$

so

$$k_r^{\text{can}} = m_B^{-1} K_B|_r.$$

When $A_B|_r = w_B \text{Id} + \Delta_r$, this becomes

$$k_r^{\text{can}} = \frac{w_B \gamma_B^2}{m_B} \mu_r + \delta_r^{\text{can}}.$$

The corpus map therefore has a positive role: it sorts local calculations into spectral packages. It also keeps endpoint, ordinary Higgs, gravity, and direct boundary-energy data out of the retained-source coefficient unless they enter one of the source expansions above.

5 Source Library Map

The local source library in `docs/` enters through six roles:

Connes and spectral-action sources	vocabulary and formulas,
KK and Hilo transcripts	project source narrative,
Standard Model and Higgs sources	electroweak checks,
global-form and line-operator sources	endpoint period filters,
boundary, Calderon, defect, and CSDR digests	candidate operators and modules,
Witten tower sources	infinite-family dimension criterion.

The role map controls citation use. Source material supplies anchors, filters, operators, or comparisons; the retained wall coefficient still requires a source expansion in one trace convention. The filename-level source audit in [39] assigns each current `docs/` source file one of these roles and records its use boundary. The role-level expansion anchors in [40] display the package forms behind those roles: almost-commutative triples, Higgs Hessians, endpoint line lattices, KK/CSDR reductions, boundary triples, defect determinants, and tower families.

The translation completion audit in [8] records the corpus enumeration, coverage evidence, verification commands, and remaining retained-source target for the active Connes-translation objective.

6 Package Atlas

The package atlas in [31] condenses the file coverage into reusable Connes data blocks. The manuscript-level version is:

Package	Algebra and Hilbert data	Operator or action	Source role
Almost-commutative Standard Model	$C^\infty(M) \otimes \mathcal{A}_F$, $L^2(M, S) \otimes \mathcal{H}_F$, J , γ .	$D_M \otimes 1 + \gamma_M \otimes D_F$, inner fluctuation D_A .	Source anchor for vocabulary and ordinary gauge-Higgs formulas.
Homogeneous KK carrier	$C^\infty(Y_u)$, equivariant sectors, compact spin or Spin^c modules.	Compact $D_K(u)$, metric action, Killing generators, heat coefficients.	Gravity, gauge-inertia, and stability data.
Custodial Higgs module	Finite or compact weak algebra, $\mathfrak{m}_3 \simeq \mathbb{C}^2$, scalar module.	Inner fluctuation, covariant derivative, scalar Hessian, vector mass matrix.	Orthodox $\rho = 1$ and triplet-hazard filter.
Endpoint and lens package	Product, quotient, or crossed-product endpoint algebra; flat characters and Q_{EM} .	Endpoint Dirac or quotient operator, period map, represented charges.	Charge, global-form, and weak-inertia filters.
Branch/interface package	Singular or groupoid algebra, sheet Hilbert spaces, B_4 , E_B , F_B .	Sheet exchange, graph/anti-graph projectors, branch-domain operator, APS/link data, boundary relation.	Carrier for h ; coefficient absent without a source package.
Boundary and Schur packages	Boundary algebra, extension domain, Z_B , retained module E_B .	Γ_0, Γ_1 , Weyl/Calderon data, Schur kernel on E_B .	Candidate source after operators and trace agree.
Retained mismatch source	Module algebra on $h \in E_B$, $p \in F_B$, map $P_B : F_B \rightarrow E_B$.	$S_{\text{mis}}^{(2)} =$ $\frac{1}{2} \langle h - P_B p, A_B(h - P_B p) \rangle$.	Active target: $A_B, m_B, \gamma_B, \sigma_r$.
Determinant, finite, or auxiliary source	h -dependent operator family, finite wall module, or auxiliary module.	Determinant Hessian, finite trace action, constraint or influence Schur kernel.	Candidate source after sign, counterterm, and trace are fixed.
Fuzzy, noncommutative, Spin^c , and tower packages	Finite matrices, projective A_θ modules, Spin^c modules, or $(\mathcal{A}_\epsilon, \mathcal{H}_\epsilon)$.	Finite Dirac block, noncommutative connection, index constraint, or labelled D_ϵ family.	Repair, constraint, or dimension-growth criterion.

The table has a negative function as well. Carrier packages and filters can reject a route or

restrict modules, but the retained-source coefficient appears only in the row whose data include $P_B, A_B, m_B, \gamma_B, \sigma_r$ in one convention.

Every equation below has one of five roles. A source anchor is imported from the Connes or Standard Model NCG vocabulary. A derived statement is a calculation recorded in the project notes. A conditional statement requires named hypotheses. A filter constrains a source package without computing it. An open target names data still absent from the project.

7 Mathematical Expansion Convention

The project uses compact dictionaries and crosswalks, but a row gets derivational use only after an expansion supplies variables, modules, operator or action, domain, first and second variation or spectral readout, normalization, residuals, and status label. Claim-only prose can orient, compare, reject a shortcut, or name a target; it cannot supply a coefficient, period, multiplier, or equality. The reusable rule is recorded in [4], and corpus coverage is recorded in [5]. The package atlas in [31] records the algebra, Hilbert data, operator or action, trace, residual status, and anchors for the major Connes homes. Each compact statement is read as expanded here, expanded in a named file, cited source, conditional target, filter, obstruction, or rejected shortcut before it can enter a derivation. In this manuscript, a source statement that lacks the variables, modules, maps, domain, variation or spectral readout, normalization, residuals, and use status is demoted to orientation or control text. It supplies no source coefficient, endpoint period, sector multiplier, equality, or prediction. The present manuscript uses the following templates.

7.1 Retained Mismatch Source

The retained boundary source uses the orientation

$$P_B : F_B \longrightarrow E_B, \quad h \in E_B, \quad p \in F_B,$$

where h is the anti-graph mismatch and p is the retained response. A positive mismatch source expands as

$$S_{\text{mis}}^{(2)}(h, p) = \frac{1}{2} \langle h - P_B p, A_B(h - P_B p) \rangle_{E_B}.$$

Equivalently,

$$\begin{aligned} S_{\text{mis}}^{(2)}(h, p) &= \frac{1}{2} \langle h, A_B h \rangle_{E_B} - \text{Re} \langle h, A_B P_B p \rangle_{E_B} \\ &\quad + \frac{1}{2} \langle p, P_B^\dagger A_B P_B p \rangle_{F_B}. \end{aligned}$$

The response-module kernel before canonical normalization is

$$K_B = P_B^\dagger A_B P_B.$$

If

$$K_B|_r = \gamma_B^2 \mu_r \text{Id} + \sigma_r, \quad \|p_r\|_{\text{kin}}^2 = m_B \|p_r\|_{F_B}^2,$$

then the canonical sector readout is

$$K_r^{\text{can}} = m_B^{-1} K_B|_r = \frac{\gamma_B^2}{m_B} \mu_r \text{Id} + m_B^{-1} \sigma_r.$$

For scalar softness $A_B|_r = w_B \text{Id} + \Delta_r^A$, the common quotient is

$$q_B = \frac{w_B \gamma_B^2}{m_B}.$$

The multiplier μ_r stays neutral until $P_B^\dagger A_B P_B$ has been computed from the source package.

7.2 Schur, Boundary, and Determinant Expansions

A local or finite auxiliary source uses variables z and h :

$$S^{(2)}(h, z) = \frac{1}{2} \langle z, Kz \rangle + \text{Re} \langle z, JRh \rangle + \frac{1}{2} \langle Rh, ERh \rangle.$$

Eliminating z gives

$$A_{\text{eff}} = R^\dagger (E - J^\dagger K^{-1} J) R.$$

The finite scale belongs to K, J, E, R and to the trace used on the local module.

A boundary or singular carrier records

$$D_{\min}, \quad D_{\max}, \quad \Gamma_0, \quad \Gamma_1,$$

and the Green form

$$\omega(\phi, \psi) = \langle \Gamma_1 \phi, \Gamma_0 \psi \rangle - \langle \Gamma_0 \phi, \Gamma_1 \psi \rangle.$$

The Weyl or Calderon operator, extension relation, and finite-part trace provide domain and spectral data. Source status begins when those data compute A_B or $K_B = P_B^\dagger A_B P_B$ on the retained modules.

A determinant route starts with an h -dependent family

$$L_B(h) = L_0 + V_1[h] + \frac{1}{2} V_2[h, h] + \cdots, \quad G_0 = L_0^{-1}.$$

The second variation has the schematic form

$$\delta_h^2 \log \det L_B = \text{Tr}(G_0 V_2[\cdot, \cdot]) - \text{Tr}(G_0 V_1[\cdot] G_0 V_1[\cdot]) + \delta_h^2 S_{\text{ct}}.$$

The sign, subtraction, zero-mode treatment, counterterm, and trace are part of the source data.

7.3 Endpoint, Gravity, and Tower Expansions

Endpoint rows record representation and period data:

$$\rho_{\text{end}}(Q_{\text{EM}}), \quad Q_{\text{EM}} H_0 = 0, \quad \ker(\Phi_{\text{period}}), \quad \chi : \mathbb{Z}_n \rightarrow \text{U}(1).$$

Gauge kinetic coefficients require normalized generators:

$$f_X = \text{Tr}_F(\rho(T_X)^2) \mathcal{I}_X.$$

The weak-angle ratio has prediction status only after the weak and hypercharge coefficients use the same normalization,

$$\sin^2 \theta_W = \frac{f_2}{f_2 + f_Y}.$$

The higher-dimensional metric expansion has the form

$$g_D = g_4 + G_{ab}(y, u)(dy^a + K_i^a A^i)(dy^b + K_j^b A^j),$$

and reduction of the Einstein–Hilbert or spectral action gives

$$S \mapsto M_4^2 R_4 + f_{ij}(u) F^i F^j + G_{AB}(u) \partial u^A \partial u^B - V(u) + S_{\partial \text{or defect}}.$$

Dimension growth requires an infinite labelled family,

$$\lambda_n(\epsilon)^2 = \lambda_0(\epsilon)^2 + \frac{n^2}{R_{\text{extra}}(\epsilon)^2} + \text{residual}_n(\epsilon),$$

with a source-computed gap scale. Finite multiplet restoration gives symmetry or phase restoration unless such a tower is present.

8 Connes Anchors

8.1 Basic Spectral Data

Source anchor. A spectral triple is written as

$$(\mathcal{A}, \mathcal{H}, D).$$

In even or real cases the additional data are

$$\gamma, \quad J,$$

with J carrying the right action and the KO-signs. In an almost-commutative product the source convention used here is

$$\mathcal{A} = C^\infty(M) \otimes \mathcal{A}_F, \quad \mathcal{H} = L^2(M, S) \otimes \mathcal{H}_F,$$

$$D = D_M \otimes 1 + \gamma_M \otimes D_F.$$

The inner fluctuation has the source form

$$D_A = D + A + \epsilon' J A J^{-1}, \quad A = \sum_i a_i [D, b_i].$$

The bosonic action is read through the spectral action

$$\text{Tr } f(D_A/\Lambda).$$

Source anchor. The Standard Model finite algebra used as the NCG reference is

$$\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}),$$

with finite metric dimension zero and finite KO-dimension six modulo eight. The distinction between metric dimension and KO-dimension is part of the translation: a finite algebra can carry KO-signs without becoming a smooth six-dimensional compact space.

8.2 Source Dimension and Probe Dimension

Conditional statement. A charged sector can resolve only the compressed algebra and operator available to that sector:

$$\mathcal{A}_{\text{eff}}(E, q) = \Pi_{E,q} \mathcal{A}_{\text{parent}} \Pi_{E,q}, \quad P_{\text{eff}}(E, q) = \Pi_{E,q} P_{\text{parent}} \Pi_{E,q}.$$

Here E denotes the relevant energy or radius window and q denotes a charge or representation label. This formula is a compatibility framework, not a source of the retained-response coefficient. The sector dictionary used in the notes is

neutral probe	trivial or suppressed internal action,
charged IR probe	endpoint, phase, lens, or boundary-character sector,
charged middle probe	branch, quotient, fuzzy, or groupoid sector,
charged UV probe	full carrier or infinite labelled tower sector.

The neutral line is a coupling or representation constraint. It cannot erase the charged-sector source data, and the observed sector dimension cannot replace the source expansion that computes A_B .

8.3 KK, QFT, and Connes Layer Parallel

Diagnostic translation. A proposed source relation should have three compatible readings: a KK geometric or boundary operator, a QFT quadratic source, and a Connes spectral representation. The comparison below records where the current programme has structure and where the source equation remains absent.

Layer	KK reading	QFT reading	Connes reading
Carrier metric	Compact space, fibration, radius, shape, branch locus.	Background fields and kinetic operator.	Spectral triple $(\mathcal{A}, \mathcal{H}, D)$, possibly singular or groupoid.
Gauge	Bundle connection or off-diagonal metric.	Yang–Mills connection and covariant derivative.	Inner fluctuation $D_A = D + A + \epsilon' JAJ^{-1}$.
Scalar/Higgs	Radius, shape, Wilson-line, boundary, or vertical deformation.	Scalar multiplet, potential, and mass matrix.	Finite/off-diagonal Dirac fluctuation or compact Dirac-family modulus.
Branch interface	Branched cover, fixed locus, collar, sheet exchange.	Defect, boundary condition, interface field, or constrained boundary value.	Singular triple, boundary triple, graph and anti-graph modules.
Retained source	Wall softness, boundary compliance, response map.	Two-field Hessian, Schur complement, determinant, or constraint source.	$E_B, F_B, P_B, A_B, m_B, \gamma_B, \sigma_r$.
Quantised labels	Harmonic sectors, monopole sectors, flat characters.	Multiplet, spin, isospin, and charge sectors.	Irreducible modules, K -theory classes, finite representations.
Endpoint	D5 lens, $S^5 \times S^2$, D7 Aloff–Wallach, EM period.	Charge lattice, line operators, global gauge group.	Product, quotient, or crossed-product endpoint triples.
Emergent dimension	Closing compact gap and infinite labelled tower.	KK tower with radius-dependent masses.	Family $(\mathcal{A}_\epsilon, \mathcal{H}_\epsilon, D_\epsilon)$ with dimension-spectrum change.

The missing relation can enter only on a row where the same operator crosses all three readings and returns a trace-normalized source package. A harmonic label without the QFT source action

gives representation bookkeeping. A QFT coefficient without the spectral module gives a fit. A finite Connes block without the KK endpoint and trace convention gives an algebraic toy model. Thus the present negative result is specific: the endpoint, Higgs, and module rows are populated, while the retained-source row still lacks a cross-column operator.

9 The One-Line Source Problem

Open target. The central NCG problem is to find a spectral, singular, boundary, determinant, or finite wall source that computes a boundary quadratic form on a retained mismatch h . The data should include

$$E_B, \quad F_B, \quad P_B, \quad A_B, \quad m_B, \quad \gamma_B, \quad \sigma_r.$$

The intended meanings are:

E_B	Hilbert module for the anti-graph mismatch h ,
F_B	Hilbert module for the retained response p ,
$P_B : F_B \rightarrow E_B$	source-derived response map,
A_B	wall kernel or softness operator on E_B ,
m_B	response trace coefficient,
γ_B	map-to-trace normalization,
σ_r	residual sector shift.

For a retained round sector the neutral source target is

$$P_B^\dagger A_B P_B|_r = q_B \mu_r \text{Id} + \sigma_r.$$

The multiplier μ_r is kept neutral until the named operator has been computed. If a source returns a common q_B different from one, the output is a scaled relation. If q_B or the residuals have unacceptable sector dependence, the branch leaves the unscaled comparison.

Conditional normalization check. For a compliance or soft-wall source with Schur weight $W|_r = w_B \text{Id} + \Delta_r^W$, response map $P_B = \gamma_B P_0$, and response trace $\|p_r\|_{\text{kin}}^2 = m_B |p_r|^2$, the canonical readout is

$$k_r^{\text{can}} = \left[\frac{w_B \gamma_B^2}{m_B} \right] \mu_r + \delta_r^{\text{can}}.$$

For a folded-collar scalar Robin source in one branch trace convention, the same test reduces to

$$q_B = \kappa_B \gamma_B^2.$$

The unscaled branch therefore requires a source-computed value $q_B = 1$ with controlled residuals. A common source-computed value different from one gives a scaled branch, and sector dependence rejects unscaled use.

The normalization split has two separate quotients:

$$c_B = \frac{\beta_B}{\alpha_B^2}, \quad q_B = \frac{w_B \gamma_B^2}{m_B}.$$

The first quotient controls the shape of the signed two-field Hessian. The second controls the map scale after the response trace on F_B is used. The two tests can pass or fail independently.

Conditional hierarchy check. Let Λ_B be the boundary Weyl or Dirichlet-to-Neumann operator of the folded branch collar, with retained sector compression

$$\Pi_r \Lambda_B \Pi_r = \lambda_r \text{Id} + \Xi_r.$$

For scalar wall softness $C_B|_r = \kappa_B \text{Id}$, set

$$\epsilon_r = \frac{\kappa_B}{\lambda_r}.$$

The soft-wall readout becomes

$$k_r^{\text{can}} = \frac{q_B}{1 + \epsilon_r} \mu_r + \sigma_r^{\text{can}}.$$

Thus q_B alone does not admit the branch: the same D6 source has to compute λ_r , the hierarchy ϵ_r , and the canonical sector shifts. Hard-wall order returns to direct boundary spectral data.

The operator search therefore has NCG status before it has numerical status. Candidate sources include a normalized group-action map, a boundary square-root operator, an interface polar modulus, a branch-link tangential operator, a finite interface block, or a Schur boundary source. Each candidate has to state its algebra, Hilbert module, domain, trace convention, and residual terms before μ_r can be evaluated.

The normalized group-action reference package is expanded in [19]. For a unitary module V_r and orthonormal generators X_i , it uses

$$P_{\text{act},r} \psi = \sum_i \rho_r(X_i) \psi \otimes e_i, \quad P_{\text{act},r}^\dagger P_{\text{act},r} = - \sum_i \rho_r(X_i)^2.$$

Irreducibility gives a neutral operator output

$$P_{\text{act},r}^\dagger P_{\text{act},r} = \mu_r^{\text{act}} \text{Id}, \quad \mu_r^{\text{act}} = - \frac{1}{\dim V_r} \sum_i \text{Tr}_{V_r}(\rho_r(X_i)^2).$$

If a D6 wall source produces

$$P_B = \gamma_B P_{\text{act}} + P_{\text{low}}, \quad A_B = \kappa_B \text{Id} + \Delta_B,$$

then

$$P_B^\dagger A_B P_B|_r = \kappa_B \gamma_B^2 \mu_r^{\text{act}} \text{Id} + \sigma_r.$$

The retained comparison still depends on

$$q_B = \kappa_B \gamma_B^2 / m_B$$

and on the residual norm. The group action therefore computes the neutral multiplier, not the wall coefficient.

10 Geometry as Families of Spectral Triples

10.1 Compact Metrics

Source anchor and conditional translation. For a smooth compact internal space K_d , the commutative internal triple is

$$\mathcal{A}_K = C^\infty(K_d), \quad \mathcal{H}_K = L^2(K_d, S_K), \quad D_K = D_{g_K(u)}.$$

The parameter u denotes radion-free shape data. Changing u changes the Dirac operator and hence the metric data of the triple.

For a fibration metric with connection, the higher-dimensional Dirac operator has the schematic conditional form

$$D_{4+d}(g, u, A) = D_4^{\text{hor}}(A, u) + \gamma_4 \otimes D_K(u) + \text{connection and curvature terms}.$$

Those extra terms encode the O'Neill and mean-curvature corrections of the KK metric.

10.2 Spectral Action and Einstein-Hilbert Reduction

Source anchor. The spectral action expansion has the form

$$\text{Tr } f(D_{4+d}^2/\Lambda^2) \sim \sum_k f_k \Lambda^k a_k(D_{4+d}^2).$$

The relevant coefficients contain the volume term, scalar curvature, connection curvature, scalar kinetic terms from metric variation, and boundary or defect contributions when such structures are present.

Derived bookkeeping. In the Einstein-frame reduction the corresponding zero-mode data are

$$g_{\mu\nu}^E, \quad A_\mu^I, \quad \phi, \quad u^A.$$

They translate respectively to the external Dirac operator, horizontal lifts or automorphism fluctuations, the overall spectral scale of D_K , and the family $D_K(u)$.

Derived scaling warning. For a common internal length L ,

$$\int_K \sqrt{g_K} R_K \sim L^{d-2}.$$

Thus the D5, D6, and D7 compact sectors have far larger raw gravitational inertia than the D1, D2, and D3 prototypes. The low-dimensional ladder is useful for topology and phase intuition, while the CP2-suspended ladder controls the quantitative compact action.

11 D1/D2/D3 and D5/D6/D7

11.1 Low-Dimensional Prototype

Derived comparison. The prototype triples are:

$$\begin{array}{lll} D1 & S^1 & \text{circle triple, zero compact scalar curvature,} \\ D2 & S^2 & \text{two-dimensional axis triple, topological curvature integral,} \\ D3 & S^3 & \text{Hopf carrier, nonzero compact curvature.} \end{array}$$

The prototype supports the familiar sequence from phase, to projective axis, to full Hopf orbit. It does not provide the CP2-suspended action coefficients.

11.2 CP2-Suspended Carriers

Derived and conditional translation. The D5 endpoint is the circle bundle

$$S^1 \longrightarrow M_5(n) = S(O(n)) \longrightarrow \mathbb{CP}^2.$$

NCG treatment uses a quotient, crossed-product, boundary, or endpoint triple. The connection curvature nF_{FS} enters the compact metric, and APS or Calderon data enter through boundary spectral terms.

Derived and conditional translation. The D6 mediator is

$$\mathbb{CP}^1 \longrightarrow Y_6 = \mathrm{SU}(3)/T^2 \longrightarrow \mathbb{CP}^2.$$

The spin^c adiabatic index argument can reduce to a CP^2 calculation, but the gravity action still sees finite vertical fibre curvature. The index limit and the metric stationarity equation are separate calculations. The $\mathrm{CP}^2/\mathrm{S}^4$ branch ingredient is

$$\pi_4 : \mathbb{CP}^2 \longrightarrow S^4, \quad \mathrm{Fix}(\mathrm{conj}) = \mathbb{RP}^2, \quad B_4 = p^{-1}(\mathbb{RP}^2) \subset Y_6.$$

The Connes package in [10] represents the two-sheet complement by a singular or groupoid algebra, with graph and anti-graph boundary modules. Its source expansion uses a finite-part trace R_B , a local branch field z , and the Schur kernel

$$A_{\mathrm{br}} = R_B^\dagger \left[E_h - E_h L_z (K_z + L_z^\dagger E_h L_z)^{-1} L_z^\dagger E_h \right] R_B.$$

Thus the branched cover supplies the D6 singular carrier and source template; the coefficient waits for $K_z, E_h, L_z, R_B, P_B, m_B, \gamma_B$ in one trace convention.

Derived and conditional translation. The D7 carrier is

$$\mathrm{SO}(3) \longrightarrow X_{1,1} = \mathrm{SU}(3)/\mathrm{U}(1)_{1,1} \longrightarrow \mathbb{CP}^2.$$

The homogeneous metric split is

$$\mathfrak{m} = \mathfrak{m}_0 \oplus \mathfrak{m}_3, \quad \dim_{\mathbb{R}} \mathfrak{m}_0 = 3, \quad \dim_{\mathbb{R}} \mathfrak{m}_3 = 4.$$

The ratio between the two metric scales remains a compactification shape variable. It is a gravity datum, not a substitute for a boundary wall kernel.

12 Endpoint Filters

Filter. The Standard-Model-shaped endpoint hierarchy used by this project is

$S^5 \times S^2$	primary product diagnostic,
$D5$ lens endpoint	flat-character and electromagnetic period test,
$D7$ Aloff–Wallach carrier	round response and custodial test,
S^7	secondary comparison object.

The product $S^5 \times S^2$ keeps the CP^2 Hopf/lens data and the weak axis visible. The seven-sphere is retained for restored-fibre intuition only when a source action connects it to the active carrier.

Conditional endpoint triple. The product diagnostic can be written as

$$\mathcal{A}_{\mathrm{prod}} = C^\infty(S^5 \times S^2) \otimes \mathcal{A}_F^{\mathrm{end}}, \quad \mathcal{H}_{\mathrm{prod}} = L^2(S^5 \times S^2, S) \otimes \mathcal{H}_F^{\mathrm{end}}.$$

The $D5$ lens side can instead be represented by

$$C^\infty(M_5(n)) \quad \text{or} \quad C^\infty(S^5) \rtimes \mathbb{Z}_n$$

with flat-character data. The endpoint calculation records the Hopf or lens period, the finite representation of Q_{EM} , and the chosen Standard-Model global form. A comparison with the Aloff–Wallach carrier still has to state the map from the product fibre $S_H^1 \times S_W^2$ to the $\text{SO}(3)$ projective Hopf fibre. These data test a D6 source package; they do not compute A_B or κ_B .

Conditional caveat. A D=11 parent can carry a larger abelian charge algebra when the electromagnetic line remains a distinct low-energy period. A $B - L$ -type generator is therefore admissible in a parent bookkeeping algebra under the condition

$$Q_{\text{EM}} \text{ survives as a named endpoint line.}$$

The caveat permits a D=11 $B - L$ line alongside electromagnetic charge while keeping their endpoint roles distinct. The parent generator cannot replace Q_{EM} , and it cannot serve as the D5/D6/D7 wall coefficient source.

Conditional charge-algebra form. Let $\mathfrak{z}_{11} \subset \mathfrak{u}(\mathcal{A}_F^{11})$ be the abelian parent charge sector, represented on $\mathcal{H}_F^{11}$ by ρ_{11} . For an endpoint projection or quotient Π_{end} , define

$$\rho_{\text{end}}(X) = \Pi_{\text{end}} \rho_{11}(X) \Pi_{\text{end}}.$$

The parent $B - L$ line is compatible with the endpoint only under the independence test

$$\rho_{\text{end}}(Q_{\text{EM}}^{11}) \neq 0, \quad \rho_{\text{end}}(Q_{\text{EM}}^{11}) \notin \mathbb{R} \rho_{\text{end}}(Q_{B-L}^{11}) + \rho_{\text{end}}(\mathfrak{k}_\Gamma),$$

where $\mathfrak{k}_\Gamma$ denotes the quotient-period kernel of the chosen Standard-Model global form. Global-form and line-operator sources can test Γ , charge periods, and fractionalization data. They still act as endpoint filters rather than a source for P_B, A_B, m_B, γ_B .

13 The D6 Branch Interface

13.1 Three Objects

Derived translation. The D6 branch material separates into:

link spinor spectral data	$A_{\text{link}}, H_{\text{link}}, D_{\text{link}}, J_{\text{link}}, G_B,$
branch graph data	$A_{Br}, H_{Br}, D_{Br}, \Pi_G, \Pi_A,$
bosonic boundary module	$E_B, F_B, P_B, A_B, m_B.$

The graph projection defines a domain condition. The anti-graph projection defines the boundary coordinate. A finite wall coefficient still needs a source.

13.2 Link Monodromy and Bosonic Boundary Data

Filter. The link spinor problem carries eta, holonomy, and zero-mode constraints. Current notes record a link angular shift, but the present bosonic mismatch module remains angular-neutral. A twisted boundary repair would require

$$E_B^{tw} = E_B \otimes L_\delta, \quad F_B^{tw} = F_B \otimes L_\delta, \quad P_B^{tw} = P_B \otimes \text{Id}_{L_\delta}.$$

The display above is a conditional module change. The line L_δ has to act on h , not only on link spinors, and the normal action coefficient still has to be derived.

Conditional angular-capacity source. If a D6 source derives the line L_δ on the bosonic mismatch, the normal-circle algebra $C^\infty(S^1)$ acts through a flat module with

$$D_\theta h = i(k + \delta_{\text{eff}})h, \quad \mu_\delta = \text{dist}(\delta_{\text{eff}}, \mathbb{Z}) > 0.$$

The regular normal-disk Dirichlet form then has the small-core coefficient

$$\kappa_{\text{ang}} = 2\pi a_B^\perp \mu_\delta.$$

This coefficient has source status only after the D6 action supplies both the bosonic line on h and the normal-gradient scale $a_B^\perp$. In the current angular-neutral convention the regular core gives vanishing softness.

14 The $\mathfrak{m}_3$ Higgs Module

14.1 Symmetric Branch

Derived electroweak control. The Aloff–Wallach block

$$\mathfrak{m}_3 \simeq \mathbb{C}^2 \simeq \mathbb{R}^4$$

is one real irreducible $U(2)$ -module. In the symmetric branch

$$\langle H \rangle = 0,$$

equivariance gives

$$\text{Hess}_H V|_0 = m_{3,\text{eff}}^2 \text{Id}_{\mathfrak{m}_3}.$$

The four real scalar components have one common mass. The branch has no electroweak vacuum radius.

14.2 Broken Branch

Derived electroweak control. When the source potential returns

$$m_{3,\text{eff}}^2 < 0, \quad \lambda_3 > 0,$$

the vacuum can be written

$$\langle H \rangle = (0, v/\sqrt{2}).$$

The covariant derivative term gives

$$(M_A^2)_{ab} = K_H(T_a \langle H \rangle, T_b \langle H \rangle).$$

The scalar Hessian splits into one radial direction and three Goldstone directions. After gauge fixing, the three Goldstone coordinates supply the longitudinal modes of W^+ , W^- , and Z .

14.3 Custodial Condition

Derived electroweak control. A single scalar metric on $\mathfrak{m}_3$ gives

$$\kappa_c = \kappa_n, \quad \rho = 1.$$

The vertical block $\mathfrak{m}_0$ supplies a hazard. The selected branch needs

$$\langle \Phi_0 \rangle = 0, \quad m_{0,\text{eff}}^2 + \lambda_{03} v^2 / 2 > 0.$$

Endpoint, defect, or D6 source terms touching the Higgs block are compatible only when their induced quadratic form is scalar on $\mathfrak{m}_3$ or when their precision correction has been computed.

15 Finite Wall Modules and Relative Sources

15.1 Finite Wall Module

Open target. A finite wall source would contain

$$\mathcal{A}_{\text{wall}}, \quad \mathcal{H}_{\text{wall}}, \quad D_{\text{wall}}, \quad J_{\text{wall}}, \quad \gamma_{\text{wall}},$$

with retained spaces

$$E_B^{\text{fin}}, \quad F_B^{\text{fin}}, \quad Z_B.$$

For a quadratic finite Schur parent, the source calculation would provide

$$K_F, \quad L_F, \quad E_F, \quad P_F, \quad m_F$$

and compute

$$A_F = E_F - E_F L_F (K_F + L_F^\dagger E_F L_F)^{-1} L_F^\dagger E_F.$$

Finite charges, anomaly data, and endpoint flat characters filter the representation. They do not replace this Schur calculation.

15.2 Relative Finite Source

Conditional source form. A relative finite source has the schematic form

$$S_{\text{rel}}(Q, R) = S_0(Q) - S_1(Q, R),$$

where both S_0 and S_1 are defined by a spectral-action difference, determinant ratio, boundary improvement, or supertrace pairing before comparison. A positive square alone gives stable mismatch energy. The signed retained-response readout needs the relative source to be physical, not selected after the target relation is known.

15.3 Relative Determinant Source

Conditional source form. A determinant or gluing route has to provide an h -dependent operator family

$$L_B(h; u) = L_0(u) + V_1[h; u] + \frac{1}{2} V_2[h, h; u] + \cdots, \quad G_0 = L_0(u)^{-1}.$$

For a renormalized relative determinant,

$$\Gamma_{\text{det}}(h; u) = \frac{s}{2} (\log \text{Det}_{\text{ren}} L_B(h; u) - \log \text{Det}_{\text{ren}} L_B(0; u)) + \Gamma_{\text{ct}}(h; u),$$

the quadratic retained kernel has the form

$$A_{\text{det}} = s [\text{Tr}_{\text{wall}}(G_0 V_2) - \text{Tr}_{\text{wall}}(G_0 V_1 G_0 V_1)] + A_{\text{ct}}.$$

The determinant sign, subtraction reference, counterterm rule, and trace convention are source data. Eta, spectral-flow, and flat-lift information remain filters unless they enter through this h -dependent family and produce a positive round kernel on the retained anti-graph module.

15.4 Auxiliary Wall Sources

Conditional source form. Influence fields, gauge constraints, finite transmission multipliers, smoothing layers, and dynamical wall fields can be written as auxiliary modules over the branch algebra:

$$Z_B, \quad K_Z, \quad J_Z, \quad E_Z.$$

They produce a candidate kernel A_{aux} on E_B by a Schur, constraint, influence, or layer calculation. Linear influence and gauge constraint routes additionally need a physical sign or contour reading; a real Euclidean linear field gives a reduced stiffness unless a direct source term is present. Positive wall-field parents have the Schur form

$$A_z = E_h - E_h L_z (K_z + L_z^\dagger E_h L_z)^{-1} L_z^\dagger E_h.$$

Finite transmission gives $A_y = B_y^\dagger K_y^{-1} B_y$ only after the multiplier has a physical source. Smoothing layers give a boundary operator only after the thickness, endpoint condition, and density or mass are derived. Every variant then returns to the retained-sector test

$$A_{\text{aux}}|_r = \kappa_B \text{Id} + \Sigma_r^A, \quad q_B = \kappa_B \gamma_B^2.$$

16 Gravity, Boundary, and Yang–Mills Terms

16.1 Ordinary Gravity

Derived separation. Compact gravity supplies radion, shape, graviphoton, and compact-potential data. It does not supply the retained two-field source by itself. Boundary spectral energy has the direct form

$$\langle q_r, P_\partial q_r \rangle.$$

This stabilizes or shifts boundary modes. It becomes a retained source only after a boundary, determinant, Schur, or extension construction introduces p , P_B , A_B , and m_B .

16.2 Ordinary Gauge-Higgs Sector

Derived separation. The gauge-Higgs reduction computes the ordinary scalar potential and vector masses. It can select the $\mathfrak{m}_3$ branch, suppress $\mathfrak{m}_0$, and protect $\rho = 1$. The ordinary Hessian is a positive scalar or vector mass matrix, not the retained response block. A gauge sector source for A_B would need a derived wall field, defect field, or finite Schur parent.

17 Objections and Consistency Checks

17.1 Objections

The following routes are rejected as source derivations in the current workspace:

positive mismatch square	stable norm, no relative readout,
direct DtN energy	boundary stiffness, no retained p ,
Green form alone	mixed pairing, no response stiffness,
ordinary Higgs Hessian	gauge-Higgs masses, no wall kernel,
image response norm	loss of the multiplier spectrum,
endpoint topology	filter only,
link eta monodromy	spinor constraint, no bosonic h line.

17.2 Consistency Checks

Any future source package should pass the following checks:

1. Name the algebra, Hilbert modules, and operator.
2. Represent h and p as retained fields.
3. Compute P_B, A_B, m_B, γ_B in one trace convention.
4. Bound residual sector shifts.
5. Run the $S^5 \times S^2$, D5 lens, electromagnetic period, D7 roundness, and $\rho = 1$ filters.
6. Classify the output as unscaled, scaled, rejected, or still open.

18 Additional Translation Frontiers

18.1 Tower Criterion for an Emergent Dimension

Conditional translation. A Witten-style growth of an extra dimension has a spectral test. One needs a family

$$(\mathcal{A}_\epsilon, \mathcal{H}_\epsilon, D_\epsilon)$$

and an infinite labelled eigenfamily

$$\lambda_n(\epsilon)^2 = \lambda_0(\epsilon)^2 + \frac{n^2}{R_{\text{extra}}(\epsilon)^2} + \text{res}_n(\epsilon).$$

The label n has to come from geometry, topology, a boundary spectrum, a flat-character lift, or a defect family, and the action has to compute the gap scale. A finite restored $\mathfrak{m}_3$ multiplet gives symmetry restoration rather than this tower criterion.

The Higgs-phase dictionary translates to a phase classification of the same spectral-family problem:

$D7/D3$	symmetric carrier with the full $\mathfrak{m}_3 \simeq \mathbb{C}^2$ fluctuation block,
$D6/D2$	selected branch, projective axis, or branch-cover mediator,
$D5/D1$	phase, boundary, lens, eta, or flat-character endpoint.

For probe statements one compresses the family by sector projectors,

$$D_{\text{eff}}(E, q) = \Pi_{E,q} D_\epsilon \Pi_{E,q}.$$

The observed dimension is the spectral behaviour of D_{eff} . It does not by itself supply the retained-response coefficient.

18.2 S^5 , Spin(6), and Electromagnetism

Filter. The D5 cover has the Hopf presentation

$$S^1 \longrightarrow S^5 \longrightarrow \mathbb{CP}^2.$$

In NCG language this can be treated as an equivariant endpoint triple, a quotient triple, or a crossed-product endpoint after the lens reduction. The spin cover relation

$$\text{Spin}(6) \simeq \text{SU}(4)$$

belongs to spinor, coefficient, or finite Hilbert modules. It can organize a $3 + 1$ representation check, while electromagnetic use still requires a derived map to Q_{EM} and its period. The D=11 $B - L$ caveat above leaves that endpoint line distinct.

18.3 Noncommutative and Division-Algebra Repairs

Conditional translation. Exotic middle layers can be represented by branch groupoids, finite or fuzzy triples, or by a noncommutative torus

$$A_\theta = \langle U, V \mid VU = \exp(2\pi i\theta)UV \rangle.$$

The parameter θ , the projective module, or the finite Dirac block has to be selected by Morita equivalence, K -theory, real multiplication, Spin^c compatibility, or a D6 source action before comparison.

For a fuzzy or CSDR repair one should name a finite matrix sequence

$$\mathcal{A}_N, \mathcal{H}_N, D_N, J_N, \gamma_N, \text{Tr}_N$$

together with the isotropy embedding, equivariant field constraints, continuous flag-coset limit, finite spectrum, and trace convention. For a higher noncommutative torus one uses

$$\mathcal{A}_\Theta = \langle U_i \mid U_i U_j = \exp(2\pi i \Theta_{ij}) U_j U_i \rangle$$

with a projective module, connection, trace, and Morita or K -theory class. These data become source candidates only when their action computes the retained package P_B, A_B, m_B, γ_B . The finite fuzzy branch package in [45] makes the matrix source explicit. It uses finite retained modules

$$E_B^N, \quad F_B^N, \quad Z_B^N$$

and a finite quadratic action with operators

$$P_N : F_B^N \rightarrow E_B^N, \quad K_N : Z_B^N \rightarrow Z_B^N, \quad L_N : Z_B^N \rightarrow E_B^N, \quad E_N : E_B^N \rightarrow E_B^N.$$

Eliminating z_N gives

$$A_N^{\text{wall}} = E_N - E_N L_N (K_N + L_N^\dagger E_N L_N)^{-1} L_N^\dagger E_N.$$

The retained finite response is then

$$K_N^{\text{resp}} = P_N^\dagger A_N^{\text{wall}} P_N, \quad K_N^{\text{resp}}|_r = \gamma_N^2 \mu_{N,r} \text{Id} + \sigma_{N,r}.$$

Continuum use requires the limit of

$$\mathcal{A}_N, D_N, \text{Tr}_N, E_B^N, F_B^N, P_N, A_N^{\text{wall}}$$

to match the branch or compact package before endpoint comparison.

The noncommutative-torus projective-module package in [46] gives the parallel expansion for the irrational-quotient repair. In the two-torus case,

$$A_\theta = \langle U, V \mid VU = e^{2\pi i\theta} UV \rangle_{\text{sm}}, \quad \tau_\theta \left(\sum a_{mn} U^m V^n \right) = a_{00}.$$

Choose a projective right A_θ -module $E_{p,q}$ with positive dimension

$$d_\theta = \tau_\theta(e_{p,q}) > 0$$

and connection ∇_i satisfying

$$\nabla_i(sa) = (\nabla_i s)a + s \delta_i(a), \quad F_{12} = [\nabla_1, \nabla_2].$$

The torus Dirac operator is

$$D_\theta = \gamma^1 \nabla_1 + \gamma^2 \nabla_2 + D_{\text{low}}, \quad \mathcal{H}_\theta = L^2(E_{p,q}) \otimes \mathbb{C}^2.$$

For the retained D6 variables one tensors the branch modules by $E_{p,q}$:

$$E_B^\theta = E_B \otimes_{A_0} E_{p,q}, \quad F_B^\theta = F_B \otimes_{A_0} E_{p,q}, \quad Z_B^\theta = Z_B \otimes_{A_0} E_{p,q},$$

with

$$h_\theta \in E_B^\theta, \quad p_\theta \in F_B^\theta, \quad z_\theta \in Z_B^\theta, \quad P_\theta : F_B^\theta \rightarrow E_B^\theta.$$

The quadratic source is

$$S_\theta^{(2)}(z; h, p) = \frac{1}{2} \tau_\theta [\langle z, K_\theta z \rangle + \langle h - P_\theta p - L_\theta z, E_\theta(h - P_\theta p - L_\theta z) \rangle].$$

Its first variation in z gives

$$(K_\theta + L_\theta^\dagger E_\theta L_\theta) z = L_\theta^\dagger E_\theta (h - P_\theta p),$$

and Schur elimination gives

$$A_\theta^{\text{wall}} = E_\theta - E_\theta L_\theta (K_\theta + L_\theta^\dagger E_\theta L_\theta)^{-1} L_\theta^\dagger E_\theta.$$

The retained response reads

$$K_\theta^{\text{resp}} = P_\theta^\dagger A_\theta^{\text{wall}} P_\theta, \quad K_\theta^{\text{resp}}|_r = \gamma_\theta^2 \mu_{\theta,r} \text{Id} + \sigma_{\theta,r}.$$

With a response trace norm

$$\|p_{\theta,r}\|_{\text{kin}}^2 = m_\theta \|p_{\theta,r}\|^2,$$

the canonical sector kernel is

$$k_{\theta,r}^{\text{can}} = m_\theta^{-1} K_\theta^{\text{resp}}|_r.$$

Thus a torus parameter has coefficient relevance only after the same projective module, trace, connection, Schur kernel, map normalization, and residuals have been selected before comparison.

The D6 division-algebra clue has a compatible NCG role as a quaternionic real structure, auxiliary-field identity, or source-action constraint. It can help select the $\mathfrak{m}_3$ branch and protect the isotropic doublet metric. It becomes a retained-response coefficient only after the D6 action supplies the quadratic form.

Spin^c , anomaly, and finite-Hilbert-space data enter the same constraint layer. On the CP^2 side one may use a twisted Hilbert module

$$\mathcal{H}_{\text{Spin}^c, n} = L^2(\mathbb{CP}^2, S_{\text{Spin}^c} \otimes L_n)$$

with a twisted Dirac operator $D_{\text{Spin}^c, n}$. The line twist can constrain endpoint periods, flat lifts, finite modules, and branch trace bundles. A fermion or auxiliary package affects the retained source only when the same spectral action, determinant, or D6 source expansion produces h, p, P_B, A_B, m_B in one convention.

19 Possible Completions

The most concrete NCG completions now available are:

- 1 D6-local B_4 boundary source deriving $R_{\text{def}}, K_z, E_h, L_z$,
- 2 boundary extension with source-derived $R_{\text{def}}, K_B, E_{Bd}, L_B$,
- 3 finite wall module computing A_F and m_F ,
- 4 localized D6 wall or influence field computing A_B ,
- 5 relative determinant family $L_B(h)$ with fixed sign,
- 6 twisted boundary module after h carries L_δ .

Each route remains an open construction target until its quadratic expansion has been computed before comparison.

For the first route, the boundary-extension package can be written with a positive boundary parent

$$S_B(z; h) = \frac{\tau}{2} (\langle z, K_B z \rangle_Z + \langle R_{\text{def}} h - L_B z, E_{Bd} (R_{\text{def}} h - L_B z) \rangle_{bd}).$$

Eliminating z gives the retained kernel

$$A_B = R_{\text{def}}^\dagger \left[E_{Bd} - E_{Bd} L_B (K_B + L_B^\dagger E_{Bd} L_B)^{-1} L_B^\dagger E_{Bd} \right] R_{\text{def}}.$$

The formula has algebraic status. It becomes a source derivation only after the D6 boundary physics supplies $R_{\text{def}}, K_B, E_{Bd}, L_B$ and the same convention supplies P_B, m_B, γ_B . The source-operator status ledger [30] assembles the direct defect, positive local field, boundary-Higgs, boundary-triple, and determinant variants into the same retained readout.

A first D6-local instantiation places a field z on the branch locus B_4 :

$$K_z = D_{B_4, z}^\dagger D_{B_4, z} + M_z^2 + V_z, \quad E_h = e_h \text{Id} + E_{\text{curv}} + E_{\text{branch}}, \quad L_z = \ell_0 \text{Id} + L_1.$$

In a commuting round sector,

$$A_{B_4}|_r = \frac{e_h(M_z^2 + \nu_r)}{M_z^2 + \nu_r + e_h \ell^2} \text{Id}.$$

The instantiation remains conditional until the D6 action derives the local operators, trace map, and response convention.

The fixed-locus, boundary-Higgs, intrinsic endpoint, and twisted-source repairs can be compressed into one local expansion. Let u_B denote the localized variables on B_4 , with source-selected vacuum u_0 . The second variation has the form

$$S_{\text{loc}}^{(2)} = \frac{\tau_B}{2} (\langle \delta u, K_{\text{loc}} \delta u \rangle + 2 \text{Re} \langle \delta u, J_{\text{loc}} R_{\text{def}} h \rangle + \langle R_{\text{def}} h, E_{\text{loc}} R_{\text{def}} h \rangle),$$

and Schur reduction gives

$$A_{\text{loc}} = R_{\text{def}}^\dagger \left(E_{\text{loc}} - J_{\text{loc}}^\dagger K_{\text{loc}}^{-1} J_{\text{loc}} \right) R_{\text{def}}.$$

The positive z -parent above is the case

$$K_{\text{loc}} = K_z + L_z^\dagger E_h L_z, \quad J_{\text{loc}} = -L_z^\dagger E_h, \quad E_{\text{loc}} = E_h.$$

A boundary-Higgs scalar, twisted fixed-locus field, or intrinsic B_4 endpoint operator therefore becomes useful only after the local D6 equations fix the vacuum, Hessian, coupling, and trace convention.

20 Conclusion

The Connes translation gives a sharper mathematical form of the programme. Compact metrics become families of spectral triples. Endpoint spaces become filters through product, quotient, and homogeneous triples. The D6 branch becomes a singular or groupoid interface with separate link, graph, and bosonic boundary modules. The $\mathfrak{m}_3$ Higgs block becomes a real or quaternionic doublet module whose scalar metric gives the orthodox $\rho = 1$ result in the selected branch.

The open object is still the same concrete source datum:

$$A_B \text{ or } \kappa_B \text{ on } h, \text{ with } P_B, m_B, \gamma_B, \sigma_r \text{ computed in one convention.}$$

Until such a source is written and expanded, De Vries-style arithmetic belongs to possible completions rather than to the main derivation.

References

- [1] A. Connes, *Noncommutative Geometry and the Standard Model with Neutrino Mixing*, local transcription docs/Connes_Noncommutative_Geometry_Standard_Model_Neutrino_Mixing_2006.md.
- [2] A. Chamseddine, A. Connes, and M. Marcolli, *Gravity and the Standard Model with Neutrino Mixing*, local transcription docs/Chamseddine_Connes_Marcolli_Gravity_Standard_Model_Neutrino_Mixing_2006.md.
- [3] Project source synthesis, *KaluzaKlein Hilo Higgsing KaluzaKlein Review*, local file docs/KaluzaKlein_Hilo_Higgsing_KaluzaKlein_Review.md.
- [4] Project note, *Mathematical Expansion Standard*, notes/Mathematical_Expansion_Standard.md.
- [5] Project note, *Mathematical Expansion Coverage Audit*, notes/Mathematical_Expansion_Coverage_Audit.md.
- [6] Project note, *NCG Translation Master Ledger*, notes/NCG_Translation_Master_Ledger.md.
- [7] Project note, *NCG Continuous Paper Outline*, notes/NCG_Continuous_Paper_Outline.md.
- [8] Project note, *NCG Translation Completion Audit*, notes/NCG_Translation_Completion_Audit.md.
- [9] Project note, *NCG Boundary Extension Retained Kernel Package*, notes/NCG_Boundary_Extension_Retained_Kernel_Package.md.
- [10] Project note, *NCG CP2 S4 Branched Cover Spectral Package*, notes/NCG_CP2_S4_Branched_Cover_Spectral_Package.md.
- [11] Project note, *NCG D6 B4 Local Boundary Source Candidate*, notes/NCG_D6_B4_Local_Boundary_Source_Candidate.md.
- [12] Project note, *NCG B4 Local Consistency Expansion*, notes/NCG_B4_Local_Consistency_Expansion.md.

- [13] Project note, *NCG Weighted Quotient Source Triage Translation*,
notes/NCG_Weighted_Quotient_Source_Triage_Translation.md.
- [14] Project note, *NCG Soft Wall DtN Hierarchy Translation*,
notes/NCG_Soft_Wall_DtN_Hierarchy_Translation.md.
- [15] Project note, *NCG Twisted Angular Capacity Translation*,
notes/NCG_Twisted_Angular_Capacity_Translation.md.
- [16] Project note, *NCG Relative Determinant Gluing Source Translation*,
notes/NCG_Relative_Determinant_Gluing_Source_Translation.md.
- [17] Project note, *NCG Influence Constraint Wall Source Translation*,
notes/NCG_Influence_Constraint_Wall_Source_Translation.md.
- [18] Project note, *NCG KK QFT Connes Layer Parallel*,
notes/NCG_KK_QFT_Connes_Layer_Parallel.md.
- [19] Project note, *NCG Equivariant Group Action Multiplier Package*,
notes/NCG_Equivariant_Group_Action_Multiplier_Package.md.
- [20] Project note, *NCG Legacy Discussion Coverage Index*,
notes/NCG_Legacy_Discussion_Coverage_Index.md.
- [21] Project note, *NCG File-Level Residual Coverage Audit*,
notes/NCG_File_Level_Residual_Coverage_Audit.md.
- [22] Project note, *NCG Repository Artifact Coverage Audit*,
notes/NCG_Repository_Artifact_Coverage_Audit.md.
- [23] Project note, *NCG Claim And Task Source Ledger*,
notes/NCG_Claim_Task_Source_Ledger.md.
- [24] Project note, *NCG Wastebook Correction Ledger*,
notes/NCG_Wastebook_Correction_Ledger.md.
- [25] Project note, *NCG Archive Verification Report Ledger*,
notes/NCG_Archive_Verification_Report_Ledger.md.
- [26] Project note, *NCG Calculation Artifact Ledger*,
notes/NCG_Calculation_Artifact_Ledger.md.
- [27] Project note, *NCG Produced Output Premise Remediation Audit*,
notes/NCG_Produced_Output_Premise_Remediation_Audit.md.
- [28] Project note, *NCG Produced Deliverable Expansion Anchors*,
notes/NCG_Produced_Deliverable_Expansion_Anchors.md.
- [29] Project note, *NCG Note Multiplier Shorthand Remediation Index*,
notes/NCG_Note_Multiplier_Shorthand_Remediation_Index.md.
- [30] Project note, *NCG B4 Source Operator Status Ledger*,
notes/NCG_B4_Source_Operator_Status_Ledger.md.

- [31] Project note, *NCG Package Atlas*, `notes/NCG_Package_Atlas.md`.
- [32] Project reference, *NCG KK QFT Connes Dictionary*,
`out/NCG_KK_QFT_CONNES_DICTIONARY.md`.
- [33] Project note, *NCG Scaled Unscaled Source Normalization*,
`notes/NCG_Scaled_Unscaled_Source_Normalization.md`.
- [34] Project note, *NCG Discussion Coverage Audit*, `notes/NCG_Discussion_Coverage_Audit.md`.
- [35] Project note, *NCG Probe Filtration Dimension Translation*,
`notes/NCG_Probe_Filtration_Dimension_Translation.md`.
- [36] Project note, *NCG Produced Output Crosswalk*, `notes/NCG_Produced_Output_Crosswalk.md`.
- [37] Project note, *NCG Source Audit Corpus Map*, `notes/NCG_Source_Audit_Corpus_Map.md`.
- [38] Project note, *NCG Source Library Crosswalk*, `notes/NCG_Source_Library_Crosswalk.md`.
- [39] Project note, *NCG Source Library File Coverage Audit*,
`notes/NCG_Source_Library_File_Coverage_Audit.md`.
- [40] Project note, *NCG Source Library Expansion Anchors*,
`notes/NCG_Source_Library_Expansion_Anchors.md`.
- [41] Project note, *NCG D11 Parent Charge EM Endpoint Translation*,
`notes/NCG_D11_Parent_Charge_EM_Endpoint_Translation.md`.
- [42] Project note, *NCG S5xS2 D5 Lens Endpoint Translation*,
`notes/NCG_S5xS2_D5_Lens_Endpoint_Translation.md`.
- [43] Project note, *NCG Tower Higgs Phase Dimension Translation*,
`notes/NCG_Tower_Higgs_Phase_Dimension_Translation.md`.
- [44] Project note, *NCG Noncommutative Fuzzy Repair Translation*,
`notes/NCG_Noncommutative_Fuzzy_Repair_Translation.md`.
- [45] Project note, *NCG Fuzzy Flag Finite Matrix Source Package*,
`notes/NCG_Fuzzy_Flag_Finite_Matrix_Source_Package.md`.
- [46] Project note, *NCG Noncommutative Torus Projective Module Source Package*,
`notes/NCG_Noncommutative_Torus_Projective_Module_Source_Package.md`.
- [47] Project note, *NCG SpinC Division Algebra Bosonic Constraints*,
`notes/NCG_SpinC_Division_Algebra_Bosonic_Constraints.md`.
- [48] Project note, *NCG Spectral Multiplier Operator Search*,
`notes/NCG_Spectral_Multiplier_Operator_Search.md`.