



The **Solar System**^[c] is the gravitationally bound system of the Sun and the objects that orbit it. The largest eight objects, which form a planetary system, are, in order of distance from the Sun: four terrestrial planets, Mercury, Venus, Earth and Mars; and four giant planets which include two gas giants, Jupiter and Saturn, and two ice giants, Uranus and Neptune. The terrestrial planets, which all have a definite surface, are mostly made of rock and metal. The gas giants are mostly made of hydrogen and helium, while the ice giants are mostly made of volatile substances such as water, ammonia, and methane. The terrestrial planets are also called the *inner Solar System* and the giant planets the *outer Solar System*.

The Solar System was formed 4.6 billion years ago from the gravitational collapse of a giant interstellar molecular cloud. Over time, the cloud formed the Sun and a protoplanetary disk that gradually coalesced to form planets and other objects. That is the reason why all eight planets have an orbit that lies near the same plane. In the present day, 99.86% of the Solar System's mass is in the Sun and most of the remaining mass is contained in the planet Jupiter. Six planets and many other bodies have natural satellites or moons orbiting around them. All of the giant planets and a few smaller bodies are encircled by planetary rings, composed of ice, dust and sometimes moonlets.

There are an unknown number of smaller dwarf planets and innumerable small bodies orbiting the Sun.^[d] These objects are distributed in the asteroid belt that lies between the orbits of Mars and Jupiter, the Kuiper belt, the scattered disc that lies beyond Neptune's orbit and at even further reaches of the Solar System (in which case they are classified as extreme trans-Neptunian objects). There is consensus among astronomers on the classification of the following nine objects as dwarf planets: the asteroid Ceres, the Kuiper-belt objects Pluto, Orcus, Haumea, Quaoar, and Makemake, and the scattered-disc objects Gonggong, Eris, and Sedna.^[d] Many small-body populations, including comets, centaurs and interplanetary dust clouds, freely travel between the regions of the Solar System.

Solar System



The Sun, planets, moons and dwarf planets^[a]
(true color, size to scale, distances not to scale)

Age	4.568 billion years
------------	---------------------

Location	<u>Local Interstellar Cloud</u> , <u>Local Bubble</u> , <u>Orion–Cygnus Arm</u> , <u>Milky Way</u>
-----------------	--

Nearest star	<u>Proxima Centauri</u> (4.2441 ly) <u>Alpha Centauri</u> (4.37 ly)
---------------------	--

Nearest known planetary system	<u>Proxima Centauri system</u> (4.2441 ly)
---------------------------------------	--

Planetary system	
Semi-major axis of outer known planet (Neptune)	30.11 AU (4.5 bill. km; 2.8 bill. mi)
Distance to Kuiper cliff	~30 AU from the Sun

Populations	
Stars	1 (<u>Sun</u>)

The solar wind, a stream of charged particles flowing outwards from the Sun, creates a bubble-like region known as the heliosphere. The heliopause is the point at which pressure from the solar wind is equal to the opposing pressure of the interstellar medium; it extends out to the edge of the scattered disc. The Oort cloud, which is thought to be the source for long-period comets, may also exist at a distance roughly a thousand times further than the heliosphere. Beyond these is the end of the Solar System. The Solar System is inside the Local Interstellar Cloud, which is part of the Local Bubble, a region of the wider Orion Arm of the Milky Way, and orbits the Galactic Center, at a distance of 27 000 light years (lys). The closest star to the Solar System is Proxima Centauri at a distance of 4.25 light-years.

Formation and evolution

The Solar System formed 4.568 billion years ago from the gravitational collapse of a region within a large molecular cloud.^[e] This initial cloud was likely several light-years across and probably birthed several stars.^[10] As is typical of molecular clouds, this one consisted mostly of hydrogen, with some helium, and small amounts of heavier elements fused by previous generations of stars. As the region that would become the Solar System, known as the pre-solar nebula,^[11] collapsed, conservation of angular momentum caused it to rotate faster. The center, where most of the mass collected, became increasingly hotter than the surrounding disc.^[10] As the contracting nebula rotated faster, it began to flatten into a protoplanetary disc with a diameter of roughly 200 AU (30 billion km; 19 billion mi)^[10] and a hot, dense protostar at the center.^{[12][13]} The planets formed by accretion from this disc,^[14] in which dust and gas gravitationally attracted each other, coalescing to form ever larger bodies. Hundreds of protoplanets may have existed in the early Solar System, but they either merged or were destroyed or ejected, leaving the planets, dwarf planets, and leftover minor bodies.^{[15][16]}

Known planets	<u>Mercury</u> <u>Venus</u> <u>Earth</u> <u>Mars</u> <u>Jupiter</u> <u>Saturn</u> <u>Uranus</u> <u>Neptune</u>
Known dwarf planets	<u>Ceres</u> <u>Orcus</u> <u>Pluto</u> <u>Haumea</u> <u>Quaoar</u> <u>Makemake</u> <u>Gonggong</u> <u>Eris</u> <u>Sedna</u>
Known natural satellites	758 (285 planetary · 473 minor planetary) ^[1]
Known minor planets	1,298,410 ^{[b][2]}
Known comets	4,586 ^{[b][2]}
Identified rounded satellites	19
Orbit about <u>Galactic Center</u>	
Invariable-to-galactic plane inclination	60.19° (ecliptic)
Distance to Galactic Center	27,000 ± 1,000 ly
Orbital speed	220 km/s; 136 mi/s
Orbital period	225–250 myr
Star-related properties	
Spectral type	G2V
Frost line	≈5 AU ^[3]
Distance to <u>heliopause</u>	≈120 AU
Hill sphere radius	≈1–3 ly





Diagram of the early Solar System's protoplanetary disk, out of which Earth and other Solar System bodies formed

Due to their higher boiling points, only metals and silicates could exist in solid form in the warm inner Solar System close to the Sun, and these would eventually form the rocky planets of Mercury, Venus, Earth, and Mars. Because metallic elements only comprised a very small fraction of the solar nebula, the terrestrial planets could not grow very large. The giant planets (Jupiter, Saturn, Uranus, and Neptune) formed further out, beyond the frost line, the point between the orbits of Mars and Jupiter where material is cool enough for volatile icy compounds to remain solid. The ices that formed these planets were more plentiful than the metals and silicates that formed the terrestrial inner planets, allowing them to grow massive enough to capture large atmospheres of hydrogen and helium, the lightest and most abundant elements. Leftover debris that never became planets congregated in regions such as the asteroid belt, Kuiper belt, and Oort cloud.^[15] The Nice model is an explanation for the creation of these regions and how the outer planets could have formed in different positions and migrated to their current orbits through various gravitational interactions.^[17]

Within 50 million years, the pressure and density of hydrogen in the center of the protostar became great enough for it to begin thermonuclear fusion.^[18] The temperature, reaction rate, pressure, and density increased until hydrostatic equilibrium was achieved: the thermal pressure counterbalancing the force of gravity. At this point, the Sun became a main-sequence star.^[19] The main-sequence phase, from beginning to end, will last about 10 billion years for the Sun compared to around two billion years for all other subsequent phases of the Sun's pre-remnant life combined.^[20] Solar wind from the Sun created the heliosphere and swept away the remaining gas and dust from the protoplanetary disc into interstellar space. As helium accumulates at its core the Sun is growing brighter;^[21] early in its main-sequence life its brightness was 70% that of what it is today.^[22]

The Solar System will remain roughly as it is known today until the hydrogen in the core of the Sun has been entirely converted to helium, which will occur roughly 5 billion years from now. This will mark the end of the Sun's main-sequence life. At that time, the core of the Sun will contract with hydrogen fusion occurring along a shell surrounding the inert helium, and the energy output will be greater than at present. The outer layers of the Sun will expand to roughly 260 times its current diameter, and the Sun will become a red giant. Because of its increased surface area, the surface of the Sun will be cooler (2,600 K (2,330 °C; 4,220 °F) at its coolest) than it is on the main sequence.^[20]



Overview of the evolution of the Sun, a G-type main-sequence star. 12 billion years after being formed by the Solar System's protoplanetary disk, the Sun will expand to become a red giant; Mercury, Venus and possibly the Earth will be swallowed.

The expanding Sun is expected to vaporize Mercury as well as Venus, and render Earth uninhabitable (possibly destroying it as well). Eventually, the core will be hot enough for helium fusion; the Sun will burn helium for a fraction of the time it burned hydrogen in the core. The Sun is not massive enough to commence the fusion of heavier elements, and nuclear reactions in the core will dwindle. Its outer layers will be ejected into space, leaving behind a dense white dwarf, half the original mass of the Sun but only the size of Earth.^[23] The ejected outer layers will form what is known as a planetary nebula, returning some of the material that formed the Sun—but now enriched with heavier elements like carbon—to the interstellar medium.^[24]


Structure and composition

The word *solar* means "pertaining to the Sun", which is derived from the Latin word *sol*, meaning Sun.^[25] The Sun is the dominant gravitational member of the Solar System, and its planetary system is maintained in a relatively stable, slowly evolving state by following isolated, gravitationally bound orbits around the Sun.^[26]

Orbits

The planets and other large objects in orbit around the Sun lie near the plane of Earth's orbit, known as the ecliptic. Smaller icy objects such as comets frequently orbit at significantly greater angles to this plane.^{[27][28]} Most of the planets in the Solar System have secondary systems of their own, being orbited by natural satellites called moons. Many of the largest natural satellites are in synchronous rotation, with one face permanently turned toward their parent. The four giant planets have planetary rings, thin bands of tiny particles that orbit them in unison.^[29]

As a result of the formation of the Solar System, planets and most other objects orbit the Sun in the same direction that the Sun is rotating. That is, counter-clockwise, as viewed from above Earth's north pole.^[30] There are exceptions, such as Halley's Comet.^[31] Most of the larger moons orbit their planets in prograde direction, matching the planetary rotation; Neptune's moon Triton is the largest to orbit in the opposite, retrograde manner.^[32] Most larger objects rotate around their own axes in the prograde direction relative to their orbit, though the rotation of Venus is retrograde.^[33]



Animations of the Solar System's inner planets and outer planets orbiting; the latter animation is 100 times faster than the former. Jupiter is three times as far from the Sun as Mars.

To a good first approximation, Kepler's laws of planetary motion describe the orbits of objects around the Sun.^{[34]:433–437} These laws stipulate that each object travels along an ellipse with the Sun at one focus, which causes the body's distance from the Sun to vary over the course of its year. A body's closest approach to the Sun is called its *perihelion*, whereas its most distant point from the Sun is called its *aphelion*.^{[35]:9-6} With the exception of Mercury, the orbits of the planets are nearly circular, but many comets, asteroids, and Kuiper belt objects follow highly elliptical orbits. Kepler's laws only account for the influence of the Sun's gravity upon an orbiting body, not the gravitational pulls of different bodies upon each other. On a human time scale, these additional perturbations can be accounted for using numerical models,^{[35]:9-6} but the planetary system can change chaotically over billions of years.^[36]

The angular momentum of the Solar System is a measure of the total amount of orbital and rotational momentum possessed by all its moving components.^[37] Although the Sun dominates the system by mass, it accounts for only about 2% of the angular momentum.^{[38][39]} The planets, dominated by Jupiter, account for most of the rest of the angular momentum due to the combination of their mass, orbit, and distance from the Sun, with a possibly significant contribution from comets.^[38]

Composition

The overall structure of the charted regions of the Solar System consists of the Sun, four smaller inner planets surrounded by a belt of mostly rocky asteroids, and four giant planets surrounded by the Kuiper belt of mostly icy objects. Astronomers sometimes informally divide this structure into separate regions. The inner Solar System includes the four terrestrial planets and the asteroid belt. The outer Solar System is


beyond the asteroids, including the four giant planets.^[40] Since the discovery of the Kuiper belt, the outermost parts of the Solar System are considered a distinct region consisting of the objects beyond Neptune.^[41]

The principal component of the Solar System is the Sun, a low-mass star that contains 99.86% of the system's known mass and dominates it gravitationally.^[42] The Sun's four largest orbiting bodies, the giant planets, account for 99% of the remaining mass, with Jupiter and Saturn together comprising more than 90%. The remaining objects of the Solar System (including the four terrestrial planets, the dwarf planets, moons, asteroids, and comets) together comprise less than 0.002% of the Solar System's total mass.^[f]


The Sun is composed of roughly 98% hydrogen and helium,^[46] as are Jupiter and Saturn.^{[47][48]} A composition gradient exists in the Solar System, created by heat and light pressure from the early Sun; those objects closer to the Sun, which are more affected by heat and light pressure, are composed of elements with high melting points. Objects farther from the Sun are composed largely of materials with lower melting points.^[49] The boundary in the Solar System beyond which those volatile substances could coalesce is known as the frost line, and it lies at roughly five times the Earth's distance from the Sun.^[3]

The objects of the inner Solar System are composed mostly of rocky materials,^[50] such as silicates, iron or nickel.^[51] Jupiter and Saturn are composed mainly of gases with extremely low melting points and high vapor pressure, such as hydrogen, helium, and neon.^[51] Ices, like water, methane, ammonia, hydrogen sulfide, and carbon dioxide,^[50] have a melting points of up to a few hundred kelvins.^[51] They can be found as ices, liquids, or gases in various places in the Solar System.^[51] Icy substances comprise the majority of the satellites of the giant planets, as well as most of Uranus and Neptune (the so-called "ice giants") and the numerous small objects that lie beyond Neptune's orbit.^{[50][52]} Together, gases and ices are referred to as volatiles.^[53]

Distances and scales



The Sun's, planets', dwarf planets' and moons' size to scale, labelled. Distance of objects is not to scale.



To-scale diagram of distance between planets, with the white bar showing orbital variations. The size of the planets is not to scale.

The astronomical unit [AU] (150,000,000 km; 93,000,000 mi) would be the distance from the Earth to the Sun if the planet's orbit were perfectly circular.^[54] For comparison, the radius of the Sun is 0.0047 AU (700,000 km; 400,000 mi).^[55] Thus, the Sun occupies 0.00001% (10^{-5} %) of the volume of a sphere with a radius the size of Earth's orbit, whereas Earth's volume is roughly one millionth (10^{-6}) that of the Sun. Jupiter, the largest planet, is 5.2 astronomical units (780,000,000 km; 480,000,000 mi) from the Sun and has a radius of 71,000 km (0.00047 AU; 44,000 mi), whereas the most distant planet, Neptune, is 30 AU (4.5×10^9 km; 2.8×10^9 mi) from the Sun.^{[48][56]}

With a few exceptions, the farther a planet or belt is from the Sun, the larger the distance between its orbit and the orbit of the next nearest object to the Sun. For example, Venus is approximately 0.33 AU farther out from the Sun than Mercury, whereas Saturn is 4.3 AU out from Jupiter, and Neptune lies 10.5 AU out from Uranus. Attempts have been made to determine a relationship between these orbital distances, like the Titius–Bode law^[57] and Johannes Kepler's model based on the Platonic solids,^[58] but ongoing discoveries have invalidated these hypotheses.^[59]

Some Solar System models attempt to convey the relative scales involved in the Solar System in human terms. Some are small in scale (and may be mechanical—called orreries)—whereas others extend across cities or regional areas.^[60] The largest such scale model, the Sweden Solar System, uses the 110-metre (361 ft) Avicii Arena in Stockholm as its substitute Sun, and, following the scale, Jupiter is a 7.5-metre (25-foot) sphere at Stockholm Arlanda Airport, 40 km (25 mi) away, whereas the farthest current object, Sedna, is a 10 cm (4 in) sphere in Luleå, 912 km (567 mi) away.^{[61][62]}

If the Sun–Neptune distance is scaled to 100 metres (330 ft), then the Sun would be about 3 cm (1.2 in) in diameter (roughly two-thirds the diameter of a golf ball), the giant planets would be all smaller than about 3 mm (0.12 in), and Earth's diameter along with that of the other terrestrial planets would be smaller than a flea (0.3 mm or 0.012 in) at this scale.^[63]

Interplanetary environment



The zodiacal light, caused by interplanetary dust

The outermost layer of the Solar atmosphere is the heliosphere, which permeates much of the Solar planetary system. Along with light, the Sun radiates a continuous stream of charged particles (a plasma) called the solar wind. This stream of particles spreads outwards at speeds from 900,000 kilometres per hour (560,000 mph) to 2,880,000 kilometres per hour (1,790,000 mph),^[64] filling the vacuum between the bodies of the Solar System. The result is a thin, dusty atmosphere, called the interplanetary medium, which extends to at least 100 AU (15 billion km; 9.3 billion mi). Beyond the heliosphere, large objects remain gravitationally bound to the sun, but the flow of matter in the interstellar medium homogenizes the distribution of

micro-scale objects.^[65]

The interplanetary medium is home to at least two disc-like regions of cosmic dust. The first, the zodiacal dust cloud, lies in the inner Solar System and causes the zodiacal light. It may have been formed by collisions within the asteroid belt brought on by gravitational interactions with the planets; a more recent proposed origin is the planet Mars.^[66] The second dust cloud extends from about 10 AU (1.5 billion km; 930 million mi) to about 40 AU (6.0 billion km; 3.7 billion mi), and was probably created by collisions within the Kuiper belt.^{[67][68]}

Activity on the Sun's surface, such as solar flares and coronal mass ejections, disturbs the heliosphere, creating space weather and causing geomagnetic storms.^[69] Coronal mass ejections and similar events blow a magnetic field and huge quantities of material from the surface of the Sun. The interaction of this magnetic field and material with Earth's magnetic field funnels charged particles into Earth's upper atmosphere, where its interactions create aurorae seen near the magnetic poles.^[70] The largest stable structure within the heliosphere is the heliospheric current sheet, a spiral form created by the actions of the Sun's rotating magnetic field on the interplanetary medium.^{[71][72]}

Life habitability

Besides solar energy, the primary characteristic of the Solar System enabling the presence of life is the heliosphere and planetary magnetic fields (for those planets that have them). These magnetic fields partially shield the Solar System from high-energy interstellar particles called cosmic rays. The density of cosmic rays in the interstellar medium and the strength of the Sun's magnetic field change on very long timescales, so the level of cosmic-ray penetration in the Solar System varies, though by how much is unknown.^[73]


Earth's magnetic field also stops its atmosphere from being stripped away by the solar wind.^[74] Venus and Mars do not have magnetic fields, and as a result the solar wind causes their atmospheres to gradually bleed away into space.^[75]

The zone of habitability of the Solar System is conventionally located in the inner Solar System, where planetary surface or atmospheric temperatures admit the possibility of liquid water.^[76] Habitability might also be possible in subsurface oceans of various outer Solar System moons.^[77]

Sun

The Sun is the Solar System's star and by far its most massive component. Its large mass (332,900 Earth masses),^[78] which comprises 99.86% of all the mass in the Solar System,^[79] produces temperatures and densities in its core high enough to sustain nuclear fusion of hydrogen into helium.^[80] This releases an enormous amount of energy, mostly radiated into space as electromagnetic radiation peaking in visible light.^{[81][82]}

Because the Sun fuses hydrogen into helium at its core, it is a main-sequence star. More specifically, it is a G2-type main-sequence star, where the type designation refers to its effective temperature. Hotter main-sequence stars are more luminous but shorter lived. The Sun's temperature is intermediate between that of the hottest stars and that of the coolest stars. Stars brighter and hotter than the Sun are rare, whereas substantially dimmer and cooler stars, known as red dwarfs, make up about 75% of the stars in the Milky Way.^{[83][84]}



The Sun in true white color

The Sun is a population I star; it has a higher abundance of elements heavier than hydrogen and helium ("metals" in astronomical parlance) than the older population II stars.^[85] Elements heavier than hydrogen and helium were formed in the cores of ancient and exploding stars, so the first generation of stars had to die before the universe could be enriched with these atoms. The oldest stars contain few metals, whereas stars born later have more. This higher metallicity is thought to have been crucial to the Sun's development of a planetary system because the planets form from the accretion of "metals".^[86]

Inner Solar System


The inner Solar System is the region comprising the terrestrial planets and the asteroid belt.^[87] Composed mainly of silicates and metals,^[88] the objects of the inner Solar System are relatively close to the Sun; the radius of this entire region is less than the distance between the orbits of Jupiter and Saturn. This region is also within the frost line, which is a little less than 5 AU (750 million km; 460 million mi) from the Sun.^[27]

Inner planets

The four terrestrial or inner planets have dense, rocky compositions, few or no moons, and no ring systems. They are in hydrostatic equilibrium, forming a rounded shape, and have undergone planetary differentiation, causing chemical elements to accumulate at different radii. They are composed largely of refractory minerals such as silicates—which form their crusts and mantles—and metals such as iron and nickel which form their cores. Three of the four inner planets (Venus, Earth and Mars) have atmospheres substantial enough to generate weather; all have impact craters and tectonic surface features, such as rift valleys and volcanoes. The term *inner planet* should not be confused with *inferior planet*, which designates those planets that are closer to the Sun than Earth (i.e. Mercury and Venus).^[89]

Mercury

Mercury (0.307–0.588 AU (45.9–88.0 million km; 28.5–54.7 million mi) from the Sun^[90]) is the closest planet to the Sun. The smallest planet in the Solar System (0.055 M_{\oplus}), Mercury has no natural satellites. The dominant geological features are impact craters or basins with ejecta blankets, the remains of early volcanic activity including magma flows, and lobed ridges or rupes that were probably produced by a period of contraction early in the planet's history.^[91] Mercury's very tenuous atmosphere consists of solar-wind particles trapped by Mercury's magnetic field, as well as atoms blasted off its surface by the solar wind.^{[92][93]} Its relatively large iron core and thin mantle have not yet been adequately explained.



Overview of the Inner Solar System up to the Jovian System



The four terrestrial planets Mercury, Venus, Earth and Mars

Hypotheses include that its outer layers were stripped off by a giant impact, or that it was prevented from fully accreting by the young Sun's energy.^{[94][95]} There have been searches for "Vulcanoids", asteroids in stable orbits between Mercury and the Sun, but none have been discovered.^{[96][97]}

Venus

Venus (0.718–0.728 AU (107.4–108.9 million km; 66.7–67.7 million mi) from the Sun^[90]) is close in size to Earth (0.815 M_{\oplus}) and, like Earth, has a thick silicate mantle around an iron core, a substantial atmosphere, and evidence of internal geological activity. It is much drier than Earth, and its atmosphere is ninety times as dense. Venus has no natural satellites. It is the hottest planet, with surface temperatures over 400 °C (752 °F), mainly due to the amount of greenhouse gases in the atmosphere.^[98] The planet has no magnetic field that would prevent the depletion of its substantial atmosphere, which suggests that its atmosphere is being replenished by volcanic eruptions.^[99] A relatively young planetary surface displays extensive evidence of volcanic activity, but is devoid of plate tectonics. It may undergo resurfacing episodes on a time scale of 700 million years.^[100]

Earth

Earth (0.983–1.017 AU (147.1–152.1 million km; 91.4–94.5 million mi) from the Sun) is the largest and densest of the inner planets, the only one known to have current geological activity, and the only place in the universe where life is known to exist.^[101] Its liquid hydrosphere is unique among the terrestrial planets, and it is the only planet where plate tectonics has been observed.^[102] Earth's atmosphere is radically different from those of the other planets, having been altered by the presence of life to contain 21% free oxygen.^{[103][104]} The planetary magnetosphere shields the surface from solar and cosmic radiation, limiting atmospheric stripping and maintaining habitability.^[105] It has one natural satellite, the Moon, the only large satellite of a terrestrial planet in the Solar System.

Mars

Mars (1.382–1.666 AU (206.7–249.2 million km; 128.5–154.9 million mi) from the Sun) is smaller than Earth and Venus (0.107 M_{\oplus}). It has an atmosphere of mostly carbon dioxide with a surface pressure of 6.1 millibars (0.088 psi; 0.18 inHg); roughly 0.6% of that of Earth but sufficient to support weather phenomena.^[106] Its surface, peppered with volcanoes, such as Olympus Mons, and rift valleys, such as Valles Marineris, shows geological activity that may have persisted until as recently as 2 million years ago.^[107] Its red color comes from iron oxide (rust) in its soil,^[108] while the polar regions show white ice caps consisting largely of water.^[109] Mars has two tiny natural satellites (Deimos and Phobos) thought to be either captured asteroids,^[110] or ejected debris from a massive impact early in Mars's history.^[111]

Asteroid belt

Asteroids except for the largest, Ceres, are classified as small Solar System bodies^[d] and are composed mainly of carbonaceous, refractory rocky and metallic minerals, with some ice.^{[112][113]} They range from a few metres to hundreds of kilometres in size. Asteroids smaller than one meter are usually called meteoroids and micrometeoroids (grain-sized), with the exact division between the two categories being debated over the years.^[114] As of 2017, the IAU designates asteroids having a diameter between about 30 micrometres and 1 metre as micrometeoroids, and terms smaller particles "dust".^[115]

The asteroid belt occupies the orbit between Mars and Jupiter, between 2.3 and 3.3 AU (340 and 490 million km; 210 and 310 million mi) from the Sun. It is thought to be remnants from the Solar System's formation that failed to coalesce because of the gravitational interference of Jupiter.^[116] The asteroid belt contains tens of thousands, possibly millions, of objects over one kilometre in diameter.^[117] Despite this, the total mass of the asteroid belt is unlikely to be more than a thousandth of that of Earth.^[45] The asteroid belt is very sparsely populated; spacecraft routinely pass through without incident.^[118]

Ceres


Ceres (2.77 AU (414 million km; 257 million mi) from the Sun) is the largest asteroid, a protoplanet, and a dwarf planet.^[d] It has a diameter of slightly under 1,000 km (620 mi) and a mass large enough for its own gravity to pull it into a spherical shape. Ceres was considered a planet when it was discovered in 1801, but as further observations revealed additional asteroids, it became common to consider it as one of the minor rather than major planets.^[119] It was then reclassified again as a dwarf planet in 2006 when the IAU definition of planet was established.^{[120]:218}

Pallas and Vesta

Pallas (2.77 AU from the Sun) and Vesta (2.36 AU from the Sun) are the largest asteroids in the asteroid belt, after Ceres. They are the other two protoplanets that survive more or less intact. At about 520 km (320 mi) in diameter, they were large enough to have developed planetary geology in the past, but both have suffered large impacts and been battered out of being round.^{[121][122][123]} Fragments from impacts upon these two bodies survive elsewhere in the asteroid belt, as the Pallas family and Vesta family. Both were considered planets upon their discoveries in 1802 and 1807 respectively, and like Ceres, eventually considered minor planets with the discovery of more asteroids. Some authors today have begun to consider Pallas and Vesta as planets again, along with Ceres, under geophysical definitions of the term.^[5]

Asteroid groups

Asteroids in the asteroid belt are divided into asteroid groups and families based on their orbital characteristics. Kirkwood gaps are sharp dips in the distribution of asteroid orbits that correspond to orbital resonances with Jupiter.^[124] Asteroid moons are asteroids that orbit larger asteroids. They are not as clearly distinguished as planetary moons, sometimes being almost as large as their partners (e.g. that of 90 Antiope). The asteroid belt includes main-belt comets, which may have been the source of Earth's water.^[125]



Linear map of the inner Solar System, showing many asteroid populations

Jupiter trojans are located in either of Jupiter's L₄ or L₅ points (gravitationally stable regions leading and trailing a planet in its orbit); the term *trojan* is also used for small bodies in any other planetary or satellite Lagrange point. Hilda asteroids are in a 2:3 resonance with Jupiter; that is, they go around the Sun three times for every two Jupiter orbits.^[126] The inner Solar System contains near-Earth asteroids, many of which cross the orbits of the inner planets.^[127] Some of them are potentially hazardous objects.^[128]

Outer Solar System

The outer region of the Solar System is home to the giant planets and their large moons. The centaurs and many short-period comets also orbit in this region. Due to their greater distance from the Sun, the solid objects in the outer Solar System contain a higher proportion of volatiles, such as water, ammonia, and methane than those of the inner Solar System because the lower temperatures allow these compounds to remain solid, without significant rates of sublimation.^[15]


Outer planets

The four outer planets, also called giant planets or Jovian planets, collectively make up 99% of the mass known to orbit the Sun.^[f] Jupiter and Saturn are together more than 400 times the mass of Earth and consist overwhelmingly of the gases hydrogen and helium, hence their designation as gas giants.^[129] Uranus and Neptune are far less massive—less than 20 Earth masses (M_{\oplus}) each—and are composed primarily of ice. For these reasons, some astronomers suggest they belong in their own category, ice giants.^[130] All four giant planets have rings, although only Saturn's ring system is easily observed from Earth. The term superior planet designates planets outside Earth's orbit and thus includes both the outer planets and Mars.^[89]


The ring-moon systems of Jupiter, Saturn, and Uranus are like miniature versions of the Solar System; that of Neptune is significantly different, having been disrupted by the capture of its largest moon Triton.^[131]

Jupiter

Jupiter (4.951–5.457 AU (740.7–816.4 million km; 460.2–507.3 million mi) from the Sun^[90]), at 318 M_{\oplus} , is 2.5 times the mass of all the other planets put together. It is composed largely of hydrogen and helium. Jupiter's strong internal heat creates semi-permanent features in its atmosphere, such as cloud bands and the Great Red Spot. The planet possesses a 4.2–14 Gauss strength magnetosphere that spans 22–29 million km, making it, in certain respects, the largest object in the Solar System.^[132] Jupiter has 95 known satellites. The four largest, Ganymede,



Plot of objects around the Kuiper belt and other asteroid populations, the J, S, U and N denotes Jupiter, Saturn, Uranus and Neptune



The outer planets Jupiter, Saturn, Uranus and Neptune, compared to the inner planets Earth, Venus, Mars, and Mercury at the bottom right

Callisto, Io, and Europa, are called the Galilean moons: they show similarities to the terrestrial planets, such as volcanism and internal heating.^[133] Ganymede, the largest satellite in the Solar System, is larger than Mercury; Callisto is almost as large.^[134]

Saturn

Saturn (9.075–10.07 AU (1.3576–1.5065 billion km; 843.6–936.1 million mi) from the Sun^[90]), distinguished by its extensive ring system, has several similarities to Jupiter, such as its atmospheric composition and magnetosphere. Although Saturn has 60% of Jupiter's volume, it is less than a third as massive, at 95 M_{\oplus} . Saturn is the only planet of the Solar System that is less dense than water. The rings of Saturn are made up of small ice and rock particles.^[135] Saturn has 145 confirmed satellites composed largely of ice. Two of these, Titan and Enceladus, show signs of geological activity;^[136] they, as well as five other Saturnian moons (Iapetus, Rhea, Dione, Tethys, and Mimas), are large enough to be round. Titan, the second-largest moon in the Solar System, is bigger than Mercury and the only satellite in the Solar System to have a substantial atmosphere.^{[137][138]}

Uranus

Uranus (18.27–20.06 AU (2.733–3.001 billion km; 1.698–1.865 billion mi) from the Sun^[90]), at 14 M_{\oplus} , has the lowest mass of the outer planets. Uniquely among the planets, it orbits the Sun on its side; its axial tilt is over ninety degrees to the ecliptic. This gives the planet extreme seasonal variation as each pole points toward and then away from the Sun.^[139] It has a much colder core than the other giant planets and radiates very little heat into space.^[140] As a consequence, it has the coldest planetary atmosphere in the Solar System.^[141] Uranus has 27 known satellites, the largest ones being Titania, Oberon, Umbriel, Ariel, and Miranda.^[142] Like the other giant planets, it possesses a ring system and magnetosphere.^[143]

Neptune

Neptune (29.89–30.47 AU (4.471–4.558 billion km; 2.778–2.832 billion mi) from the Sun^[90]), though slightly smaller than Uranus, is more massive (17 M_{\oplus}) and hence more dense. It radiates more internal heat than Uranus, but not as much as Jupiter or Saturn.^[144] Neptune has 14 known satellites. The largest, Triton, is geologically active, with geysers of liquid nitrogen.^[145] Triton is the only large satellite with a retrograde orbit, which indicates that it did not form with Neptune, but was probably captured from the Kuiper belt.^[146] Neptune is accompanied in its orbit by several minor planets, termed Neptune trojans, that either lead or trail the planet by about one-sixth of the way around the Sun, positions known as Lagrange points.^[147]

Centaurs

The centaurs are icy comet-like bodies whose orbits have semi-major axes greater than Jupiter's (5.5 AU (820 million km; 510 million mi)) and less than Neptune's (30 AU (4.5 billion km; 2.8 billion mi)). These are former Kuiper belt and scattered disc objects that were gravitationally perturbed closer to the Sun by the outer planets, and are expected to become comets or get ejected out of the Solar System.^[44] While most centaurs are inactive and asteroid-like, some exhibit clear cometary activity, such as the first centaur

discovered, 2060 Chiron, which has been classified as a comet (95P) because it develops a coma just as comets do when they approach the Sun.^[148] The largest known centaur, 10199 Chariklo, has a diameter of about 250 km (160 mi) and is one of the only few minor planets known to possess a ring system.^{[149][150]}

Comets

Comets are small Solar System bodies,^[d] typically only a few kilometres across, composed largely of volatile ices. They have highly eccentric orbits, generally a perihelion within the orbits of the inner planets and an aphelion far beyond Pluto. When a comet enters the inner Solar System, its proximity to the Sun causes its icy surface to sublimate and ionise, creating a coma: a long tail of gas and dust often visible to the naked eye.^[151]




Comet Hale-Bopp seen in 1997

Short-period comets have orbits lasting less than two hundred years. Long-period comets have orbits lasting thousands of years. Short-period comets are thought to originate in the Kuiper belt, whereas long-period comets, such as Hale-Bopp, are thought to originate in the Oort cloud. Many comet groups, such as the Kreutz sungrazers, formed from the breakup of a single parent.^[152]

Some comets with hyperbolic orbits may originate outside the Solar System, but determining their precise orbits is difficult.^[153] Old comets whose volatiles have mostly been driven out by solar warming are often categorised as asteroids.^[154]

Trans-Neptunian region

Beyond the orbit of Neptune lies the area of the "trans-Neptunian region", with the doughnut-shaped Kuiper belt, home of Pluto and several other dwarf planets, and an overlapping disc of scattered objects, which is tilted toward the plane of the Solar System and reaches much further out than the Kuiper belt. The entire region is still largely unexplored. It appears to consist overwhelmingly of many thousands of small worlds—the largest having a diameter only a fifth that of Earth and a mass far smaller than that of the Moon—composed mainly of rock and ice. This region is sometimes described as the "third zone of the Solar System", enclosing the inner and the outer Solar System.^[155]



Distribution and size of trans-Neptunian objects. The horizontal axis stand for the semi-major axis of the body, the vertical axis stands for the inclination of the orbit, and the size of the circle stands for the relative size of the object.

Kuiper belt

The Kuiper belt is a great ring of debris similar to the asteroid belt, but consisting mainly of objects composed primarily of ice.^[156] It extends between 30 and 50 AU (4.5 and 7.5 billion km; 2.8 and 4.6 billion mi) from the Sun. It is composed mainly of small Solar System bodies, although the largest few are probably large enough to be dwarf planets.^[157] There are estimated to be over 100,000 Kuiper belt

objects with a diameter greater than 50 km (30 mi), but the total mass of the Kuiper belt is thought to be only a tenth or even a hundredth the mass of Earth.^[44] Many Kuiper belt objects have satellites,^[158] and most have orbits that are substantially inclined ($\sim 10^\circ$) to the plane of the ecliptic.^[159]

The Kuiper belt can be roughly divided into the "classical" belt and the resonant trans-Neptunian objects.^[156] The latter have orbits whose periods are in a simple ratio to that of Neptune: for example, going around the Sun twice for every three times that Neptune does, or once for every two. The classical belt consists of objects having no resonance with Neptune, and extends from roughly 39.4 to 47.7 AU (5.89 to 7.14 billion km; 3.66 to 4.43 billion mi).^[160] Members of the classical Kuiper belt are sometimes called "cubewanos", after the first of their kind to be discovered, originally designated 1992 QB₁; they are still in near primordial, low-eccentricity orbits.^[161]



Size comparison of some large TNOs with Earth: Pluto and its moons, Eris, Makemake, Haumea, Sedna, Gonggong, Quaoar, Orcus, Salacia, and 2002 MS₄.

Pluto and Charon

The dwarf planet Pluto (with an average orbit of 39 AU (5.8 billion km; 3.6 billion mi) from the Sun) is the largest known object in the Kuiper belt. When discovered in 1930, it was considered to be the ninth planet; this changed in 2006 with the adoption of a formal definition of planet. Pluto has a relatively eccentric orbit inclined 17 degrees to the ecliptic plane and ranging from 29.7 AU (4.44 billion km; 2.76 billion mi) from the Sun at perihelion (within the orbit of Neptune) to 49.5 AU (7.41 billion km; 4.60 billion mi) at aphelion. Pluto has a 2:3 resonance with Neptune, meaning that Pluto orbits twice round the Sun for every three Neptunian orbits. Kuiper belt objects whose orbits share this resonance are called plutinos.^[162]

Charon, the largest of Pluto's moons, is sometimes described as part of a binary system with Pluto, as the two bodies orbit a barycenter of gravity above their surfaces (i.e. they appear to "orbit each other"). Beyond Charon, four much smaller moons, Styx, Nix, Kerberos, and Hydra, orbit Pluto.^[163]


Others

Besides Pluto, astronomers generally agree that at least four other Kuiper belt objects are dwarf planets,^[157] though there is some doubt for Orcus,^[164] and additional bodies have also been proposed:^[165]

- Makemake (45.79 AU average from the Sun), although smaller than Pluto, is the largest known object in the *classical* Kuiper belt (that is, a Kuiper belt object not in a confirmed resonance with Neptune). Makemake is the brightest object in the Kuiper belt after Pluto. Discovered in 2005, it was officially named in 2009.^[166] Its orbit is far more inclined than Pluto's, at 29° .^[167] It has one known moon.^[168]
- Haumea (43.13 AU average from the Sun) is in an orbit similar to Makemake, except that it is in a temporary 7:12 orbital resonance with Neptune.^[169] Like Makemake, it was discovered in 2005.^[170] Uniquely among the dwarf planets, Haumea possess a ring system, two known moons named Hi'iaka and Namaka, and rotates so quickly (once every 3.9 hours) that it is stretched into an ellipsoid. It is part of a collisional family of Kuiper belt objects that share similar orbits, which suggests a giant collision took place on Haumea and ejected its fragments into space billions of years ago.^[171]

- Quaoar (43.69 AU average from the Sun) is the second-largest known object in the classical Kuiper belt, after Makemake. Its orbit is significantly less eccentric and inclined than those of Makemake or Haumea.^[169] It possesses a ring system and one known moon, Weywot.^[172]
- Orcus (39.40 AU average from the Sun) is in the same 2:3 orbital resonance with Neptune as Pluto, and is the largest such object after Pluto itself.^[169] Its eccentricity and inclination are similar to Pluto's, but its perihelion lies about 120° from that of Pluto. Thus, the phase of Orcus's orbit is opposite to Pluto's: Orcus is at aphelion (most recently in 2019) around when Pluto is at perihelion (most recently in 1989) and vice versa.^[173] For this reason, it has been called the *anti-Pluto*.^{[174][175]} It has one known moon, Vanth.^[176]

Scattered disc



The orbital eccentricities and inclinations of the scattered disc population compared to the classical and resonant Kuiper belt objects

The scattered disc, which overlaps the Kuiper belt but extends out to near 500 AU, is thought to be the source of short-period comets. Scattered-disc objects are believed to have been perturbed into erratic orbits by the gravitational influence of Neptune's early outward migration. Most scattered disc objects (SDOs) have perihelia within the Kuiper belt but aphelia far beyond it (some more than 150 AU from the Sun). SDOs' orbits can also be inclined up to 46.8° from the ecliptic plane.^[177] Some astronomers consider the scattered disc to be merely another region of the Kuiper belt and describe scattered-disc objects as "scattered Kuiper belt objects".^[178] Some astronomers also classify centaurs as inward-scattered Kuiper belt objects along with the outward-scattered residents of the scattered disc.^[179]

Eris and Gonggong

Eris (67.78 AU average from the Sun) is the largest known scattered disc object, and caused a debate about what constitutes a planet, because it is 25% more massive than Pluto^[180] and about the same diameter. It is the most massive of the known dwarf planets. It has one known moon, Dysnomia. Like Pluto, its orbit is highly eccentric, with a perihelion of 38.2 AU (roughly Pluto's distance from the Sun) and an aphelion of 97.6 AU, and steeply inclined to the ecliptic plane at an angle of 44°. ^[181]


Gonggong (67.38 AU average from the Sun) is another dwarf planet in a comparable orbit to Eris, except that it is in a 3:10 resonance with Neptune.^[182] It has one known moon, Xiangliu.^[183]

Farthest regions

The point at which the Solar System ends and interstellar space begins is not precisely defined because its outer boundaries are shaped by two forces, the solar wind and the Sun's gravity. The limit of the solar wind's influence is roughly four times Pluto's distance from the Sun; this *heliopause*, the outer boundary of the heliosphere, is considered the beginning of the interstellar medium.^[65] The Sun's Hill sphere, the effective range of its gravitational dominance, is thought to extend up to a thousand times farther and encompasses the hypothetical Oort cloud.^[184]

Edge of the heliosphere

The Sun's stellar-wind bubble, the heliosphere, a region of space dominated by the Sun, has its boundary at the *termination shock*, which is roughly 80–100 AU from the Sun upwind of the interstellar medium and roughly 200 AU from the Sun downwind.^[185] Here the solar wind collides with the interstellar medium^[186] and dramatically slows, condenses and becomes more turbulent,^[185] forming a great oval structure known as the *heliosheath*. This structure has been theorized to look and behave very much like a comet's tail, extending outward for a further 40 AU on the upwind side but tailing many times that distance downwind.^[187] Evidence from the *Cassini* and *Interstellar Boundary Explorer* spacecraft has suggested that it is forced into a bubble shape by the constraining action of the interstellar magnetic field,^{[188][189]} but the actual shape remains unknown.^[190]




Artistic depiction of the Solar System's heliosphere

The outer boundary of the heliosphere, the *heliopause*, is the point at which the solar wind finally terminates and is the beginning of interstellar space.^[65] *Voyager 1* and *Voyager 2* passed the termination shock and entered the heliosheath at 94 and 84 AU from the Sun, respectively.^{[191][192]} *Voyager 1* was reported to have crossed the heliopause in August 2012, and *Voyager 2* in December 2018.^{[193][194]}

The shape and form of the outer edge of the heliosphere is likely affected by the fluid dynamics of interactions with the interstellar medium as well as solar magnetic fields prevailing to the south, e.g. it is bluntly shaped with the northern hemisphere extending 9 AU farther than the southern hemisphere.^[185] Beyond the heliopause, at around 230 AU, lies the bow shock, a plasma "wake" left by the Sun as it travels through the Milky Way.^[195]

Detached objects

Sedna (with an average orbit of 520 AU from the Sun) is a large, reddish object with a gigantic, highly elliptical orbit that takes it from about 76 AU at perihelion to 940 AU at aphelion and takes 11,400 years to complete. Mike Brown, who discovered the object in 2003, asserts that it cannot be part of the scattered disc or the Kuiper belt because its perihelion is too distant to have been affected by Neptune's migration. He and other astronomers consider it to be the first in an entirely new population, sometimes termed "distant detached objects" (DDOs), which also may include the object 2000 CR₁₀₅, which has a perihelion of 45 AU, an aphelion of 415 AU, and an orbital period of 3,420 years.^[196] Brown terms this population the "inner Oort cloud" because it may have formed through a similar process, although it is far closer to the Sun.^[197] Sedna is very likely a dwarf planet, though its shape has yet to be determined. The



The detached object Sedna and its orbit within the Solar System

second unequivocally detached object, with a perihelion farther than Sedna's at roughly 81 AU, is 2012 VP₁₁₃, discovered in 2012. Its aphelion is only about half that of Sedna's, at 458 AU.^{[198][199]}

Oort cloud


The Oort cloud is a hypothetical spherical cloud of up to a trillion icy objects that is thought to be the source for all long-period comets and to surround the Solar System at roughly 50,000 AU (around 1 light-year (ly)) from the Sun, and possibly to as far as 100,000 AU (1.87 ly). It is thought to be composed of comets that were ejected from the inner Solar System by gravitational interactions with the outer planets. Oort cloud objects move very slowly, and can be perturbed by infrequent events, such as collisions, the gravitational effects of a passing star, or the galactic tide, the tidal force exerted by the Milky Way.^{[200][201]}

Boundaries

Much of the Solar System is still unknown. The Sun's gravitational field is estimated to dominate the gravitational forces of surrounding stars out to about two light-years (125,000 AU). Lower estimates for the radius of the Oort cloud, by contrast, do not place it farther than 50,000 AU.^[202] Most of the mass is orbiting in the region between 3,000 and 100,000 AU.^[203] Despite discoveries such as Sedna, the region between the Kuiper belt and the Oort cloud, an area tens of thousands of AU in radius, is still virtually unmapped. Learning about this region of space is difficult, because it depends upon inferences from those few objects whose orbits happen to be perturbed such that they fall closer to the Sun, and even then, detecting these objects has often been possible only when they happened to become bright enough to register as comets.^[204] Objects may yet be discovered in the Solar System's uncharted regions.^[205] The furthest known objects, such as Comet West, have aphelia around 70,000 AU from the Sun.^[206]

Comparison with other star systems

Compared to many extrasolar systems, the Solar System stands out in lacking planets interior to the orbit of Mercury.^{[207][208]} The known Solar System also lacks super-Earths, planets between one and ten times as massive as the Earth,^[207] although the hypothetical Planet Nine, if it does exist, could be a super-Earth orbiting in the outer Solar System.^[209] Uncommonly, it has only small rocky planets and large gas giants; elsewhere planets of intermediate size are typical—both rocky and gas—so there is no "gap" as seen between the size of Earth and of Neptune (with a radius 3.8 times as large). As many of these super-Earths are closer to their respective stars than Mercury is to the Sun, a hypothesis has arisen that all planetary systems start with many close-in planets, and that typically a sequence of their collisions causes consolidation of mass into few larger planets, but in case of the Solar System the collisions caused their destruction and ejection.^{[207][210]}



Habitable zones of TRAPPIST-1 and the Solar System; here, the TRAPPIST-1 system is enlarged 25 times. The displayed planetary surfaces on TRAPPIST-1 are speculative.

The orbits of Solar System planets are nearly circular. Compared to other systems, they have smaller orbital eccentricity.^[207] Although there are attempts to explain it partly with a bias in the radial-velocity detection method and partly with long interactions of a quite high number of planets, the exact causes remain

Location

Celestial neighborhood




Diagram of the Local Interstellar Cloud, the G-Cloud and surrounding stars. As of 2022, the precise location of the Solar System in the clouds is an open question in astronomy.^[212]

The Solar System is surrounded by the Local Interstellar Cloud, although it is not clear if it is embedded in the Local Interstellar Cloud or if it lies just outside the cloud's edge.^{[213][214]} Multiple other interstellar clouds also exist in the region within 300 light-years of the Sun, known as the Local Bubble.^[214] The latter feature is an hourglass-shaped cavity or superbubble in the interstellar medium roughly 300 light-years across. The bubble is suffused with high-temperature plasma, suggesting that it may be the product of several recent supernovae.^[215]

The Local Bubble is a small superbubble compared to the neighboring wider Radcliffe Wave and *Split* linear structures (formerly Gould Belt), each of which are some thousands of light-years in length.^[216] All these structures are part of the Orion Arm, which contains most of the stars in the Milky Way that are visible to the unaided eye. The density of all matter in the local neighborhood is $0.097 \pm 0.013 \text{ } M_{\odot} \cdot \text{pc}^{-3}$.^[217]

Within ten light-years of the Sun there are relatively few stars, the closest being the triple star system Alpha Centauri, which is about 4.4 light-years away and may be in the Local Bubble's G-Cloud.^[218] Alpha Centauri A and B are a closely tied pair of Sun-like stars, whereas the closest star to Earth, the small red dwarf Proxima Centauri, orbits the pair at a distance of 0.2 light-year. In 2016, a potentially habitable exoplanet was found to be orbiting Proxima Centauri, called Proxima Centauri b, the closest confirmed exoplanet to the Sun.^[219]

The next closest known fusors to the Sun are the red dwarfs Barnard's Star (at 5.9 ly), Wolf 359 (7.8 ly), and Lalande 21185 (8.3 ly).^[220] The nearest brown dwarfs belong to the binary Luhman 16 system (6.6 ly), and the closest known rogue or free-floating planetary-mass object at less than 10 Jupiter masses is the sub-brown dwarf WISE 0855-0714 (7.4 ly).^[221]

Just beyond at 8.6 ly lies Sirius, the brightest star in Earth's night sky, with roughly twice the Sun's mass, orbited by the closest white dwarf to Earth, Sirius B. Other stars within ten light-years are the binary red-dwarf system Gliese 65 (8.7 ly) and the solitary red dwarf Ross 154 (9.7 ly).^{[222][223]} The closest solitary Sun-like star to the Solar System is Tau Ceti at 11.9 light-years. It has roughly 80% of the Sun's mass but only about half of its luminosity.^[224]

The nearest and unaided-visible group of stars beyond the immediate celestial neighborhood is the Ursa Major moving group at roughly 80 light-years, which is within the Local Bubble, like the nearest as well as unaided-visible star cluster the Hyades, which lie at its edge. The closest star-forming regions are the Corona Australis Molecular Cloud, the Rho Ophiuchi cloud complex and the Taurus molecular cloud; the latter lies just beyond the Local Bubble and is part of the Radcliffe wave.^[225]

Galactic context

The Solar System is located in the Milky Way, a barred spiral galaxy with a diameter of about 100,000 light-years containing more than 100 billion stars.^[226] The Sun resides in one of the Milky Way's outer spiral arms, known as the Orion–Cygnus Arm or Local Spur.^[227] The Sun lies about 26,660 light-years from the Galactic Center (where the supermassive black hole Sagittarius A* is located),^[228] and its speed around the center of the Milky Way is about 220 km/s, so that it completes one revolution every 240 million years.^[226] This revolution is known as the Solar System's galactic year.^[229] The solar apex, the direction of the Sun's path through interstellar space, is near the constellation Hercules in the direction of the current location of the bright star Vega.^[230] The plane of the ecliptic lies at an angle of about 60° to the galactic plane.^[g]




Diagram of the Milky Way on its side, with galactic features and the relative position of the Solar System labelled.

The Solar System's location in the Milky Way is a factor in the evolutionary history of life on Earth. Its orbit is close to circular, and orbits near the Sun are at roughly the same speed as that of the spiral arms.^{[232][233]} Therefore, the Sun passes through arms only rarely. Because spiral arms are home to a far larger concentration of supernovae, gravitational instabilities, and radiation that could disrupt the Solar System, this has given Earth long periods of stability for life to evolve.^[232] However, the changing position of the Solar System relative to other parts of the Milky Way could explain periodic extinction events on Earth, according to the Shiva hypothesis or related theories, but this remains controversial.^{[234][235]}

The Solar System lies well outside the star-crowded environs of the Galactic Center. Near the center, gravitational tugs from nearby stars could perturb bodies in the Oort cloud and send many comets into the inner Solar System, producing collisions with potentially catastrophic implications for life on Earth. The intense radiation of the Galactic Center could also interfere with the development of complex life.^[232] Stellar flybys that pass within 0.8 light-years of the Sun occur roughly once every 100,000 years. The closest well-measured approach was Scholz's Star, which approached to 52^{+23}_{-14} kAU of the Sun some 70^{+15}_{-10} kya, likely passing through the outer Oort cloud.^[236]

Humanity's perspective

Humanity's knowledge of the Solar System has grown incrementally over the centuries. Up to the Late Middle Ages–Renaissance, astronomers from Europe to India believed Earth to be stationary at the center of the universe^[237] and categorically different from the divine or ethereal objects that moved through the sky. Although the Greek philosopher Aristarchus of Samos had speculated on a heliocentric reordering of

23°00' 22°30' 22°00'

• ϕ Aqr

• λ Aqr

• θ Aqr

• ψ^3 Aqr

ECLIPTIC

• σ Aqr

• τ^7 Aqr

• ι Aqr

• δ Aqr

• 66 Aqr

• 88 Aqr

• 89 Aqr

2003-06-15

In the 17th century, Galileo publicized the use of the telescope in astronomy; he and Simon Marius independently discovered that Jupiter had four satellites in orbit around it.^[242] Christiaan Huygens followed on from these observations by discovering Saturn's moon Titan and the shape of the rings of Saturn.^[243] In 1677, Edmond Halley observed a transit of Mercury across the Sun, leading him to realize that observations of the solar parallax of a planet (more ideally using the transit of Venus) could be used to trigonometrically determine the distances between Earth, Venus, and the Sun.^[244] Halley's friend Isaac Newton, in his magisterial *Principia Mathematica* of 1687, demonstrated that celestial bodies are not quintessentially different from Earthly ones: the same laws of motion and of gravity apply on Earth and in the skies.^{[34]:142}

In the 20th century, humans began their space exploration around the Solar System, starting with placing telescopes in space.^[252] Since then, humans have landed on the Moon during the Apollo program; the Apollo 13 mission marked the furthest any human has been away from Earth at 400,171 kilometers (248,655 mi).^[253] All eight planets and two dwarf planets have been visited by space probes. This began with Mariner 2's fly-by of Venus in 1962, while Mariner 9 mission to Mars was the first to orbit another planet in 1971. The outer planets were first visited by Pioneer 10's encounter with Jupiter, and Pioneer 11's encounter with Saturn. The remaining gas giants were first visited by the Voyager spacecraft, one of which (Voyager 1) is the furthest object made by humankind and the first in interstellar space.^[254] In addition,

probes have also returned samples from comets^[255] and asteroids,^[256] as well as flown through the Sun's corona^[257] and made fly-bys of Kuiper belt objects.^[258] Six of the planets (all but Uranus and Neptune) have or had a dedicated orbiter.^[259]

See also



[solar system portal](#)



[outer space portal](#)



[astronomy portal](#)

- [List of gravitationally rounded objects of the Solar System](#)
- [List of Solar System objects by size](#)
- [Lists of geological features of the Solar System](#)
- [List of Solar System extremes](#)
- [Outline of the Solar System](#)

Notes

- The [asteroid belt](#) and [Kuiper belt](#) are not added because the individual asteroids are too small to be shown on the diagram.
- As of 14 August 2023.
- Capitalization** of the name varies. The [International Astronomical Union](#), the authoritative body regarding astronomical nomenclature, specifies capitalizing the names of all individual astronomical objects but uses mixed "Solar System" and "solar system" structures in their [naming guidelines document](http://www.iau.org/public/themes/naming/) (<http://www.iau.org/public/themes/naming/>) [Archived](https://web.archive.org/web/20210725053113/https://www.iau.org/public/themes/naming) (<https://web.archive.org/web/20210725053113/https://www.iau.org/public/themes/naming>) 25 July 2021 at the [Wayback Machine](#). The name is commonly rendered in lower case ('solar system'), as, for example, in the [Oxford English Dictionary](#) and [Merriam-Webster's 11th Collegiate Dictionary](#) (<http://www.m-w.com/dictionary/solar%20system>) [Archived](https://web.archive.org/web/20080127201148/http://www.m-w.com/dictionary/solar%20system) (<https://web.archive.org/web/20080127201148/http://www.m-w.com/dictionary/solar%20system>) 27 January 2008 at the [Wayback Machine](#).
- According to [IAU definitions](#), objects orbiting the Sun are classified dynamically and physically into three categories: *planets*, *dwarf planets*, and *small Solar System bodies*.
 - A [planet](#) is any body orbiting the Sun whose [mass](#) is sufficient for [gravity](#) to have pulled it into a (near-)[spherical](#) shape and that has [cleared its immediate neighborhood](#) of all smaller objects. By this definition, the Solar System has eight planets: Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune. Because it has not cleared its neighborhood of other [Kuiper belt](#) objects, Pluto does not fit this definition.^[4]
 - A [dwarf planet](#) is a body orbiting the Sun that is massive enough to be made near-spherical by its own gravity but that has not cleared [planetesimals](#) from its neighborhood and is also not a satellite.^[4] Dwarf planets are considered planets by some planetologists but not by the IAU.^[5] The IAU has recognized four other bodies in the Solar System as dwarf planets: Ceres, Haumea, Makemake, and Eris.^{[6][7]} Other objects [commonly accepted as dwarf planets](#) include [Gonggong](#), [Sedna](#), [Orcus](#), and [Quaoar](#). In a reference to Pluto, other dwarf planets orbiting in the [trans-Neptunian region](#) are sometimes called "plutoids",^[8] though this term is seldom used.
 - The remaining objects orbiting the Sun are known as [small Solar System bodies](#).^[4]