

RBE 500 Homework #3

Arjan Gupta

Problem 4.2

Verify Equation (4.7) by direct calculation.

$$S(a)p = a \times p \tag{4.7}$$

Solution

Suppose the vectors a and p are given as

$$a = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}, p = \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}$$

By the definition of the cross-product, we know

$$a \times p = \begin{bmatrix} a_2p_3 - a_3p_2 \\ a_3p_1 - a_1p_3 \\ a_1p_2 - a_2p_1 \end{bmatrix} \tag{1}$$

Also, by the definition of skew-symmetric matrices, we know the form of $S(a)$, where a is the vector we have already defined,

$$S(a) = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix}$$

Hence, by normal matrix multiplication,

$$S(a)p = \begin{bmatrix} 0 - a_3p_2 + a_2p_3 \\ a_3p_1 + 0 - a_1p_3 \\ -a_2p_1 + a_1p_2 + 0 \end{bmatrix} = \begin{bmatrix} a_2p_3 - a_3p_2 \\ a_3p_1 - a_1p_3 \\ a_1p_2 - a_2p_1 \end{bmatrix}$$

Which is the same as (1). Therefore, Equation (4.7) is proved.

Problem 4.3

Prove the assertion given in Equation (4.9) that $R(a \times b) = Ra \times Rb$ for $R \in SO(3)$.

Solution

Let $v = R(a \times b)$, and $u = Ra \times Rb$. If v and u are to be proved as the same vector, then it must be shown that they have the exact same magnitude and direction. Let us consider magnitude and direction of v and u separately.

Magnitude

Since $R \in SO(3)$, $\det R = 1$, which means that the linear transformation R does not change the length (norm) of any vector that it transforms (rotates). Hence we can say

$$\|R(a \times b)\| = \|a \times b\| \quad (1)$$

$$\|Ra\| = \|a\| \quad (2)$$

$$\|Rb\| = \|b\| \quad (3)$$

Using the definition of the cross product along with (1), we can state that

$$\|v\| = \|R(a \times b)\| = \|a \times b\| = \|a\|\|b\| \sin \theta \quad (4)$$

Again using the definition of the cross product along with (2) and (3), we can state that

$$\|u\| = \|Ra \times Rb\| = \|Ra\|\|Rb\| \sin \theta = \|a\|\|b\| \sin \theta \quad (5)$$

We can see that (4) and (5) are equal. Therefore,

$$\|v\| = \|u\|$$

Which means that the magnitude of v and u are the same.

Direction

Since $R \in SO(3)$ is a merely a rotational transformation of the 3 dimensional vector space, any directional relationships given by the right-hand curl rule before R has been applied must remain preserved and obtainable by the right-hand curl rule even after R has been applied.

Now, let us assume that the vectors a and b lie in the plane P . Also, assume $a \times b$ lies in the direction given by \hat{q} . We know that \hat{q} can be obtained by applying the right-hand curl rule from a to b .

After R is applied, P becomes plane P_R , and \hat{q} becomes \hat{q}_R . This means that $v = R(a \times b)$ lies along \hat{q}_R . Also, vector a and b become vectors Ra and Rb , which now lie in plane P_R . However, we can still obtain \hat{q}_R by applying the right-hand curl rule from Ra to Rb , since this relationship is preserved. Hence, $u = Ra \times Rb$ lies in the direction \hat{q}_R . Therefore, v and u lie in the same direction.

Since v and u lie in the same direction and have the same magnitude, they are the same vector. Hence, (4.9) is proved.

Problem 4.5

Suppose that $a = (1, -1, 2)$ and that $R = R_{x,90}$. Show by direct calculation that $RS(a)R^T = S(Ra)$.

Solution

We have,

$$R = R_{x,90} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(90) & -\sin(90) \\ 0 & \sin(90) & \cos(90) \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}$$
$$S(a) = \begin{bmatrix} 0 & -2 & -1 \\ 2 & 0 & -1 \\ 1 & 1 & 0 \end{bmatrix}$$

Now,

$$RS(a)R^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -2 & -1 \\ 2 & 0 & -1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

=?