

RBE 500 Homework #4

Arjan Gupta

Problem 4.6

Given $R = R_{x,\theta}R_{y,\phi}$, compute $\frac{\partial R}{\partial \phi}$. Evaluate $\frac{\partial R}{\partial \phi}$ at $\theta = \frac{\pi}{2}$, $\phi = \frac{\pi}{2}$. First parametrically compute, then evaluate by plugging the values in.

Solution

$$\frac{\partial}{\partial \phi} (R_{x,\theta}R_{y,\phi}) = R_{x,\theta} \frac{\partial}{\partial \phi} (R_{y,\phi})$$

Using the the fact that $\frac{d}{d\theta} (R_{y,\theta}) = S(j)R_{y,\theta}$,

$$\begin{aligned} R_{x,\theta} \frac{\partial}{\partial \phi} (R_{y,\phi}) &= R_{x,\theta} S(j) R_{y,\phi} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} \cos \phi & 0 & \sin \phi \\ 0 & 1 & 0 \\ -\sin \phi & 0 & \cos \phi \end{bmatrix} \\ &= \begin{bmatrix} -\sin(\phi) & 0 & \cos(\phi) \\ \cos(\phi) \sin(\theta) & 0 & \sin(\phi) \sin(\theta) \\ -\cos(\phi) \cos(\theta) & 0 & -\cos(\theta) \sin(\phi) \end{bmatrix} \end{aligned}$$

Now, plugging in the values $\theta = \frac{\pi}{2}$, $\phi = \frac{\pi}{2}$, we get

$$\begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

We performed the above computations using the following MATLAB code.

```

1  % Calculation code for problem 4.6 of the RBE500 textbook (HW 4)
2
3  clear; close all; clc;
4
5  syms theta phi
6
7  % Define the matrices in our problem
8  rotx_theta = [1 0 0; 0 cos(theta) -sin(theta); 0 sin(theta) cos(theta)];
9  Sj = [0 0 1; 0 0 0; -1 0 0];
10 roty_phi = [cos(phi) 0 sin(phi); 0 1 0; -sin(phi) 0 cos(phi)];
11
12 % Multiply the matrices
13 product = rotx_theta*Sj*roty_phi;
14
15 % Get latex output
16 latex(product)
17
18 phi_val = pi/2;
19 theta_val = pi/2;
20
21 % Now plug in values
22 product_val = [ -sin(phi_val), 0, cos(phi_val);
23 cos(phi_val)*sin(theta_val), 0, sin(phi_val)*sin(theta_val);
24 -cos(phi_val)*cos(theta_val), 0, -cos(theta_val)*sin(phi_val)]

```

Problem 4.10

Two frames $o_0x_0y_0z_0$ and $o_1x_1y_1z_1$ are related by the homogeneous transformation

$$H = \begin{bmatrix} 0 & -1 & 0 & 1 \\ 1 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

A particle has velocity $v_1(t) = (3, 1, 0)$ relative to frame $o_1x_1y_1z_1$. What is the velocity of the particle in frame $o_0x_0y_0z_0$?

Solution

The given H is the homogeneous transformation H_1^0 . Therefore, $R_1^0 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $o_1^0 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$.

At any given point in time, the position of the particle with respect to frame $o_1x_1y_1z_1$ is given as $p^1(t)$. We know that

$$p^0(t) = R_1^0 p^1(t) + o_1^0$$

Taking the derivative of both sides and using the product rule, we get

$$\dot{p}^0(t) = \dot{R}_1^0 p^1(t) + R_1^0 \dot{p}^1(t) + 0$$

But, $\dot{R}_1^0 = 0$ since R_1^0 is a constant in time. Therefore,

$$\begin{aligned} v^0(t) &= R_1^0 v^1(t) \\ v^0(t) &= \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix} \\ v^0(t) &= \begin{bmatrix} -1 \\ 3 \\ 0 \end{bmatrix} \end{aligned}$$

Which was calculated by the following MATLAB script


```
1 % Calculation code for problem 4.10 of the RBE500 textbook (HW 4)
2
3 clear; close all; clc;
4
5 R = [0 -1 0; 1 0 0; 0 0 1];
6 v1 = [3; 1; 0];
7
8 v0 = R*v1
```

Problem 4.15

Find the 6×3 Jacobian for the three links of the cylindrical manipulator of Figure 3.7. Find the singular configurations for this arm.

Solution

Figure 3.7 of our book is the following.



We can see that we have 3 joints, so $n = 3$. Let us also form the table given in the lecture videos:

| | Linear component | Angular component |
|-----------------|--|-----------------------|
| Revolute joint | $J_{v_i} = z_{i-1}^0 \times (o_n^0 - o_{i-1}^0)$ | $J_{v_i} = z_{i-1}^0$ |
| Prismatic joint | $J_{\omega_i} = z_{i-1}^0$ | $J_{\omega_i} = 0$ |

Using this table, and the fact that the upper half of the Jacobian contains linear components while the bottom half contains angular components, we have

$$J = \begin{bmatrix} z_0 \times (o_3 - o_0) & z_1 & z_2 \\ z_0 & 0 & 0 \end{bmatrix}$$

In this Jacobian matrix, we know that $z_0 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ and $o_0 = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$. To find z_1 , z_2 , and o_3 , we need to find

$T_n^0 = A_1 \dots A_n$ for $n = 1, 2, 3$. By following the DH convention (which Figure 3.7 already abides by), we have the following table for quantities $\alpha_i, a_i, \theta_i, d_i$.

| Link | α_i | a_i | θ_i | d_i |
|------|------------|-------|------------|-------|
| 1 | 0 | 0 | θ_1 | d_1 |
| 2 | 0 | 0 | 0 | d_2 |
| 3 | 0 | 0 | 0 | d_3 |

Which gives us the following A_i matrices.

$$A_1 = \begin{bmatrix} \cos(\theta_1) & -\sin(\theta_1) & 0 & 0 \\ \sin(\theta_1) & \cos(\theta_1) & 0 & 0 \\ 0 & 0 & 1 & d_1 \\ 0 & 0 & 0 & 1 \end{bmatrix}, A_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & d_2 \\ 0 & 0 & 0 & 1 \end{bmatrix}, A_3 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & d_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Now we can compute our T matrices using the following MATLAB code,

```

1  % Calculation code for problem 4.15 of the RBE500 textbook (HW 4)
2
3  clear; close all; clc;
4
5  syms theta1 d1 d2 d3;
6
7  % Form the A matrices
8  A1 = [cos(theta1) -sin(theta1) 0 0; sin(theta1) cos(theta1) 0 0; 0 0 1 d1; 0 0 0 1];
9  A2 = [1 0 0 0; 0 1 0 0; 0 0 1 d2; 0 0 0 1];
10 A3 = [1 0 0 0; 0 1 0 0; 0 0 1 d3; 0 0 0 1];
11
12 % Compute T matrices
13 T2 = A1*A2;
14 T3 = A1*A2*A3;
15
16 % Output to LaTeX
17 latex(T2)
18 latex(T3)

```

$$T_1^0 = A_1 = \begin{bmatrix} \cos(\theta_1) & -\sin(\theta_1) & 0 & 0 \\ \sin(\theta_1) & \cos(\theta_1) & 0 & 0 \\ 0 & 0 & 1 & d_1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T_2^0 = A_1 A_2 = \begin{bmatrix} \cos(\theta_1) & -\sin(\theta_1) & 0 & 0 \\ \sin(\theta_1) & \cos(\theta_1) & 0 & 0 \\ 0 & 0 & 1 & d_1 + d_2 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T_3^0 = A_1 A_2 A_3 = \begin{bmatrix} \cos(\theta_1) & -\sin(\theta_1) & 0 & 0 \\ \sin(\theta_1) & \cos(\theta_1) & 0 & 0 \\ 0 & 0 & 1 & d_1 + d_2 + d_3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

From these T matrices, we get

$$z_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, z_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, o_3 = \begin{bmatrix} 0 \\ 0 \\ d_1 + d_2 + d_3 \end{bmatrix}$$

Now we are ready to compute our Jacobian matrix.