ISOTOPIA-2.23

Simulation of medical isotope production

Arjan Koning

s

Copyright © 2026 Arjan Koning

nds.iaea.org/talys

ISOTOPIA is free software: you can redistribute it and/or modify it under the terms of the MIT License.

Permission is hereby granted, free of charge, to any person obtaining a copy of this software and associated documentation files (the "Software"), to deal in the Software without restriction, including without limitation the rights to use, copy, modify, merge, publish, distribute, sublicense, and/or sell copies of the Software, and to permit persons to whom the Software is furnished to do so, subject to the following conditions:

The above copyright notice and this permission notice shall be included in all copies or substantial portions of the Software.

THE SOFTWARE IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE SOFTWARE OR THE USE OR OTHER DEALINGS IN THE SOFTWARE.

Typesetting: The Legrand Orange Book, LaTeX Template, Version 2.1.1 (14/2/16), downloaded from: `www.LaTeXTemplates.com`. Original author: Mathias Legrand (legrand.mathias@gmail.com) with modifications by: Vel (vel@latextemplates.com). License: CC BY-NC-SA 3.0 `creativecommons.org/licenses/by-nc-sa/3.0/`.

March 2026

About the author

Arjan Koning is a nuclear physicist with a Masters Degree in Theoretical physics at the Univ. of Amsterdam, a PhD in the Natural Sciences on Multi-step direct reactions at the Univ. of Groningen, and a Professorship at the Univ. of Uppsala on theoretical nuclear reactions.
Arjan is currently Head of the Nuclear Data Section at the IAEA in Vienna. Before that, he has worked at ECN/NRG Petten, the Netherlands, on nuclear reaction data for science and technology, and as guest scientist at CEA/Bruyères-le-Châtel and Los Alamos National Laboratory on the development and computational implementation of nuclear reaction models. He has led several students to PhD degrees, has coordinated and chaired various international nuclear data projects such as the OECD/NEA JEFF and WPEC projects, and has advised governments and international organisations on nuclear research and development. Among his scientific accomplishments are innovations in nuclear reaction physics, especially for the optical model and pre-equilibrium reactions, and initiating the '3 T's': the TALYS nuclear model code, Total Monte Carlo uncertainty propagation and the TENDL nuclear data library. He has been awarded the Oppenheimer Coin from Los Alamos National Laboratory, and the title of Doctor Honoris Causa by Uppsala University. As of 2025, his h-index is 62 with more than 26 000 citations.
Although Arjan is currently in a managerial role, he aims to keep his scientific creativity alive by maintaining and extending TALYS plus all software that emerges from that. Pleas from his friends to also spend time on other things are sometimes honoured.

Preface

ISOTOPIA is a software package for the prediction of medical isotope production. In principle, the isotope yields for any production route with incident neutrons, photons, protons, deuterons, tritons, helions or alpha particles, for natural or enriched targets, can be calculated. While most computational ingredients for the simulation of isotope production are rather straightforward, the *crucial* ingredients for reliable simulation of the isotope yield are nuclear reaction cross sections radioactive decay data. For ISOTOPIA, we use the IAEA medical isotope data library for about 150 reaction channels, complemented by TENDL-2025 for all other reactions.

The idea to construct ISOTOPIA is rooted in the belief that radioisotopes produced with *both* reactors and accelerators are needed in the foreseeable future, to ensure that medical diagnosis and therapy remain successful *and* economically affordable. Since several years, this issue has entered the discussion with the need for the replacement of old reactors that produce isotopes. Accelerators, which are mostly used to produce neutron-poor PET isotopes, and also several therapeutic isotopes, are now also suggested as a possible alternative production device for "reactor nuclides" with the ^{99}Mo/^{99m}Tc generator as the most important case. While this issue has economical, logistic and political aspects, the current code will at least help to answer its scientific aspects. ISOTOPIA predicts the production yield of any diagnostic, therapeutic or theranostic isotope with reactors or accelerators (for both photons and charged-particles). Obviously, the *reliability* of that prediction depends strongly on the quality of the nuclear reaction cross sections.

At certain moments in time, a well-defined version of ISOTOPIA is frozen. You are now reading the tutorial of version 2.23. Future versions should become more and more reliable through improvements in both the ISOTOPIA code and the underlying cross section and radioactive decay data libraries.

ISOTOPIA is the engine behind the Medical Isotope Browser, nds.iaea.org/mib, which allows to analyze production routes in an efficient way via a GUI.

License, contact and reference

As mentioned on the first page and in the source code, ISOTOPIA falls in the category of MIT License software.

In addition to the MIT *terms* I have a *request*:

- When ISOTOPIA is used for your reports, publications, etc., please make a proper reference to the code. At the moment this is:
  A.J. Koning, ISOTOPIA: Simulation of medical isotope production with accelerators, unpublished tutorial (2025).

The webpage for ISOTOPIA is **nds.iaea.org/talys**.

Acknowledgements

I wish to thank a few persons who have contributed to the present state of ISOTOPIA:

- Marieke Duijvestijn, who set up the first calculation routes with the TALYS code for isotopic yields, which we needed for a project on the production of ^{68}Ga, when we were still working at NRG Petten,
- Marco Verpelli, for integrating ISOTOPIA in the Medical Isotope Browser and for suggesting the associated improvements to the code.
- Steven van der Marck for his initial help to construct a software tool like this and useful discussions,
- Will Webster and Bernard Erasmus for important feedback on the implementation of ISOTOPIA in the TALYS code,
- Roberto Capote Noy and the entire consortium responsible for the IAEA medical isotope data library, for helpful discussions and for providing the specific data library for about 150 isotopes,
- Natalie Gaughan, for testing many of the isotope yield calculations with the TALYS code,
- Ulli Koester, for taking a critical look at our Medical Isotope Browser and suggesting improvements.

Arjan Koning

Contents

1 Introduction 9
1.1 How to use this tutorial 11

2 Installation and getting started 13
2.1 The ISOTOPIA package 13
2.2 Installation 13
2.3 Verification 14
2.4 Getting started 14

3 Input description 15
3.1 Basic input rules 16

4 Radioisotope production 17
4.1 Equations for production and depletion of nuclides 17
4.2 Reaction constants and rates 19
4.2.1 Neutron irradiation 20
4.2.2 Photon irradiation 21
4.2.3 Charged-particle irradiation 21
4.3 Activities 23
4.4 Possible refinements 23

5 Nuclear data libraries 25
5.1 Radioactive decay data library 25

5.2 Cross section data library 25
5.2.1 Neutrons 25
5.2.2 Photonuclear 25
5.2.3 Charged particles 26
5.3 Creation of medical isotope data library 26

6 Reference Guide 31
6.1 ISOTOPIA keywords 31

7 Sample cases 63
7.1 Sample p-Mo100-Tc099m: Accelerator production of ^{99m}Tc 63
7.2 Sample a-Bi209-At211: alpha-induced production of therapeutic nuclide ^{211}At 67
7.3 Sample p-Ga069-Ge068: Production of the ^{68}Ge generator for ^{68}Ga 67
7.4 Sample p-Th232-Ac225: Production of ^{225}Ac from ^{232}Th 68

8 Outlook and conclusions 69

Bibliography 69

A Derivation of the nuclide production equations 73
A.1 Differential equations for production and depletion of nuclides 73
A.2 Solution of the differential equations for production and depletion 75
A.3 Solution of the differential equations for depletion only (cooling) 77

1. Introduction

The current worldwide market for medial isotopes is estimated to be of the order of 6-8 billion dollars per year, and is expected to increase to 15 billion dollar in the year 2035. This places high demands on the production capacity of these isotopes, in the form of facilities of different type, like accelerators, research reactors and other neutron sources.

The essential feature of medical isotopes is that they release radiation in the form of alpha-, beta- or gamma-particles which are then used for diagnosis, therapy or a combination of the two (theranostics). This means that the particular isotope has to be radioactive and must be formed by transmutation through a nuclear reaction. There are three main irradiation sources to produce medical isotopes

- charged-particle accelerators with particle types ranging from protons (the dominating route) to heavy ions
- photonuclear reactions
- neutron sources, usually from research reactors

The nuclear reactions induced by these sources fall in two categories

- activation reactions, whereby the produced isotope is relatively close to the target, formed by a relatively simple reaction,
- fission reactions leading to fission products, with the production of ^{99}Mo produced by research reactors as the best known example

For diagnosis, almost all radioisotopes fall into the categories of single photon emission computed tomography (SPECT) or positron emission tomography (PET). Although PET scans are qualitatively superior and therefore its market is enlarging, the ease of use and cost effectiveness of SPECT, with the ^{99}Mo/^{99m}Tc generator as the workhorse, entails that its dominance will probably not disappear in the coming decade. As a rule of thumb (exceptions exist!) neutron-rich nuclides, often retrieved as fission products or by activation via neutron capture in a nuclear reactor, are used for SPECT scans while neutron-poor nuclides, produced with accelerators, are used for PET scans. For therapy, also both reactor-produced and accelerator-produced isotopes are on the market.

The current version of ISOTOPIA is devoted to the production of isotopes with a charged-particle accelerator, photon or neutron source.

Often, on the basis of its biological, chemical and nuclear decay properties a radioisotope is identified as a promising candidate for medical diagnosis or therapy. The subsequent question "What is the best way to produce that medical isotope?" can not always be answered by a firm "yes" or "no" but fragments into several other questions, such as:

- What is the optimal production route for the isotope, i.e., what is the best target + projectile combination?
- For charged-particle irradiation, what is the optimal incident beam energy, the optimal target thickness, or equivalently, what is the projectile energy range inside the target?
- What is the optimal irradiation time and cooling period?
- What is the isotopic yield during and directly after irradiation and after a cooling period?
- What is the yield of elemental and isotopic contaminations after irradiation and after a cooling period?
- How much do the answers to the above questions depend on the use of an enriched or a natural target?

To answer such questions, we have implemented analytical models which allows to parameterize the various characteristics of a reactor/accelerator + target combination and which calculates the isotopic yields of various production routes as a function of irradiation time. This is possible since we now have modern cross section and decay data libraries.

Of course, the nuclear physics aspects to the above questions are not the end of the story. Several questions on economics and logistics (often correlated) need to be answered as well, e.g.

- How much irradiation is needed to meet the current demand?
- What are the logistical implications of the radioactive decay time of the desired product and its purity, which may depend on the production route?
- What are the costs to make a sufficiently enriched target?

Finally, a political decision to build a production facility is made which may discard the scientific and economical answers, but we have decided not to touch economical, logistical or political aspects here and concentrate ourselves on the purely scientific aspects.

As specific features of ISOTOPIA we mention

- Estimation of isotopic yield on the basis of a user-specified irradiation source and energy range, target thickness, irradiation time and cooling period. Covered are incident neutrons, photons, protons, deuterons, tritons, helions and alpha particles, on any type of target material.
- The activities of all residual products are presented as yields as a function of irradiation time. When appropriate, results are averaged over time to obtain the average yield in [MBq/mAh] of [MBq/(g.h)].
- Estimate of simultaneous production of contaminations.
- Robustness: thanks to the complete TENDL data library augmented by well-evaluated reaction channels of the IAEA medical isotope library, ISOTOPIA can be used for the prediction of any isotope that one is interested in: it is generally applicable and has been thoroughly tested on a basic quality level.
- A transparent source program.
- Input/output communication that is easy to use and understand.
- A user tutorial.
- A few sample cases.

The starting point of all calculations are cross sections for all reaction channels. They are obtained from two sources,

- the TENDL-2025 nuclear data library [1], produced by the nuclear reaction model code TALYS [2] and other sources,

- the IAEA medical isotope database, high-quality in-depth evaluations of important reaction channels, on the basis of available experimental data [3, 4, 5, 6].

The current tutorial only describes the ISOTOPIA code and its input and output. We stress however that a powerful Graphical User Interface has been built around it at IAEA, the Medical Isotope Browser, nds.iaea.org/mib, which drives the ISOTOPIA code and which is a very efficient way to study medical isotope production as a function of the irradiation characteristics.

1.1 How to use this tutorial

Although we would be honored if you would read this tutorial from the beginning to the end, we can imagine that not all parts are necessary, relevant or suitable to you. the contents are as follows:

Chapter 2: Installation guide for ISOTOPIA.
Chapter 3: The main input rules
Chapter 4: General formalism for isotope production.
Chapter 5: The nuclear reaction and structure information needed in the form of cross section and decay data.
Chapter 6: The reference guide with all the keywords
Chapter 7: Sample cases and verification.
Chapter 8: Outlook and conclusions.

2. Installation and getting started

2.1 The ISOTOPIA package

In what follows we assume ISOTOPIA will be installed on a Linux or MacOS operating system. The total ISOTOPIA package is in the *isotopia/* directory and contains the following directories and files:

- *LICENSE* is the license file,
- *README.md* outlines the contents of the package.
- *install_isotopia.bash*, *code_build.bash* and *path_change.bash* are scripts that take care of the installation,
- *source/* contains the Fortran source code of ISOTOPIA
- *files/* contains the natural abundances and the decay data library.
- *doc/* contains the documentation: this tutorial in pdf format.
- *samples/* contains the input and output files of the sample cases.

Outside *isotopia/*, you need to have the directory *isotopia.libs/* which contains all cross section data. In fact the place of this directory can be set in the *machine.f90* subroutine.

In total, you will need about 20 Gb of free disk space to install ISOTOPIA, which is almost entirely due to the cross section database in the *isotopia.libs/* directory. The code has so far *only* been tested on various Linux and MacOS systems, so we can not guarantee that it works on all operating systems. The only OS dependencies we can think of are the directory separators '/' we use in pathnames that are hardwired in the code. If there is any dependence on the operating system, the associated statements can be altered in the subroutine *machine.f90*.

ISOTOPIA has been successfully tested for the gfortran compiler.

2.2 Installation

The installation of ISOTOPIA is straightforward. You can download ISOTOPIA via either git

- **git clone https://github.com/arjankoning1/isotopia.git**

or by getting the tar file

- from **https://nds.iaea.org/talys/isotopia.tar**
- **tar zxf isotopia.tar**

We here provide the necessary steps to do the installation, For a Unix/Linux system, the installation is expected to be done by the simple command

- *install_isotopia.bash*

which calls the scripts

- *code_build.bash* where you may set, if needed, the name of your compiler and the place where you want to store the ISOTOPIA executable.
- **path_change.bash** to set the pathname in the source code.

An alternative installation option is

- **cd isotopia/source**
- **make**

If this does not work for some reason, we here provide the necessary steps to do the installation manually. For a Unix/Linux system, the following steps should be taken:

- **cd isotopia/source** If *code_build.bash* has not already replaced the path name in *machine.f90*, do it yourself. We think this is the only Unix/Linux machine dependence of ISOTOPIA. We expect no complaints from the compiler.
- **gfortran -c *.f90**
- **gfortran *.o -o isotopia**
- **mv isotopia ../bin**

The above commands represent the standard compilation options. Consult the tutorial of your compiler to get an enhanced performance with optimization flags enabled. We note that an ISOTOPIA run generally takes only less than a second so extreme optimization is not necessary.

2.3 Verification

If ISOTOPIA is installed, testing the sample cases is the logical next step. The *samples/* directory contains the script *verify* that runs all the test cases. Each sample case has its own subdirectory, which contains a subdirectory *org/*, where we stored the input files and *our* calculated results. It also contains a subdirectory *new*, where we have stored the input files only and where the *verify* script will produce *your* output files. A full description of the keywords used in the input files is given in Chapter 3. Chapter 7 describes all sample cases in full detail. Note that in each subdirectory a file with differences with our original output is created.

2.4 Getting started

If you have created your own working directory with an input file named e.g. *isotopia.inp*, then a ISOTOPIA calculation can easily be started with:

isotopia < **isotopia.inp** > **isotopia.out**

where the names *isotopia.inp(out)* are not obligatory: you can use any name for these files. Besides the general output file *isotopia.out* various other output files, with activities per isotope,are produced.

3. Input description

For the communication between ISOTOPIA and its users, we have constructed an input/output method which shields beginners from all the possible options for parameters that can be specified in ISOTOPIA, while enabling at the same time maximal flexibility for experienced users.

An input file of ISOTOPIA consists of keywords and their associated values. Before we list all the input possibilities, let us illustrate the use of the input by the following example. It represents a minimum input file for ISOTOPIA:

```
projectile p
element    Mo
mass       100
ebeam     16.
```

This input file represents the simplest question that can be asked to ISOTOPIA: if a 100 % enriched ^{100}Mo target is bombarded by 16 MeV protons, what isotopes are produced as a function of time, and what is the radioactive yield? Behind this simple input file, however, there are several default values for the various assumptions, parameters, output flags, etc., that you may or may not be interested in. When you use a minimal input file like the one above, you leave it to the author of ISOTOPIA to choose all the parameters for you, including accelerator and target specification, as well as the level of detail of the output file. If you want to use specific parameters other than the default, or want to have more specific information in the output file(s), more keywords are required. Obviously, more keywords means more flexibility and, in the case of adequate use, better results. For example, one probably would like to enter the accelerator beam current, target size and material density in the input. In this Chapter, we will first give the basic rules that must be obeyed when constructing an input file for ISOTOPIA.

3.1 Basic input rules

Theoretically, it would be possible to make the use of ISOTOPIA completely idiot-proof, i.e. to prevent the user from any input mistakes that possibly can be made and to continue a calculation with "assumed" values. Although we have invested a relatively large effort in the user-friendliness of ISOTOPIA, we have not taken such safety measures to the extreme limit and ask at least some minimal responsibility from the user. Once you have accepted that, only very little effort is required to work with the code. Successful execution of ISOTOPIA can be expected if you stick to the following simple rules and possibilities of the input file:

1. One input line contains one keyword. Usually it is accompanied by only one value, but some keywords (e.g. **Tcool**) need to be accompanied by more than one value on the same line.
2. A keyword and its value(s) *must* be separated by at least 1 blank character.
3. The keywords can be given in arbitrary order. If you use the same keyword more than once, the value of the last one will be adopted.
4. All characters can be given in either lowercase or uppercase.
5. A keyword *must* be accompanied by a value. To use default values, the keywords should simply be left out of the input file.
6. An input line starting with a # in column 1 is neglected. This is helpful for including comments in the input file or to temporarily deactivate keywords.
7. A minimal input file always consists of 4 lines and contains the keywords **projectile**, **element**, **mass** and **ebeam**. These 4 keywords *must* be given in any input file.
8. An input line may not exceed 80 characters.

As an example of rules 3, 4, and 6, it can be seen that the following input file is completely equivalent to the one given in the beginning of this chapter:

```
Ebeam     16.
Element    Mo
MASS       100
projectile p
#rho 5.
```

In the following erroneous input file, only the second and third lines are correct, while rules 2, and 5 are violated in the other lines.

```
Ebeam22.
projectile p
MASS 103
Element
```

In cases like this, ISOTOPIA will give a specific error message for the first encountered problem and the execution will be stopped. We like to believe that we have covered all such cases and that it is impossible to let ISOTOPIA crash (at least with our compilers) without giving an appropriate error message, but you are of course invited to prove and let us know about the contrary (Sorry, no cash rewards). Typing errors in the input file should be spotted by ISOTOPIA, e.g. if you write **projjectile d**, it will tell you the keyword is not in our list, and the code will gracefully stop.

4. Radioisotope production

This section describes the equations to calculate the production yield of a radioactive nucleus by a nuclear reaction. Starting from a general formalism detailed in the Appendix one can introduce realistic approximations and derive simpler equations that hold in basically all cases of interest. In ISOTOPIA we have implemented only analytical solutions to the production and depletion equations, which are expected to be sufficient for nuclides formed by activation.

4.1 Equations for production and depletion of nuclides

The production and depletion of radionuclides can be described by a system of Bateman equations for the number of nuclides N of type i at any time t.

$$\frac{dN_i(t)}{dt} = \sum_k (\lambda_{ki} + R_{ki}) N_k(t) - (\lambda_i + \sum_j R_{ij}) N_i(t), \tag{4.1}$$

where N_i can be produced by nuclear reactions via all feeding nuclides k with reaction constants R, while they are simultaneously depleted through reactions to all nuclides j and decay constants λ. The most general solution of this Equation is obtained with matrix exponential solvers. However, in almost all cases of interest for radioisotope production by means of activation, as detailed in the Appendix, one can make well-justified assumptions to come to the following temporal development of a system of nuclides during irradiation,

$$\begin{aligned}
N_T(t) &= N_0\, e^{-R_T t}, \\
N_p(t) &= N_0 \frac{R_{Tp}}{\lambda_p - R_T} \left[e^{-R_T t} - e^{-\lambda_p t}\right], \\
N_i(t) &= N_0 \frac{R_{Ti}}{\lambda_i - R_T} \left[e^{-R_T t} - e^{-\lambda_i t}\right] + N_0 \frac{R_{Tp}\lambda_{pi}}{\lambda_p - R_T} \left[\frac{e^{-R_T t} - e^{-\lambda_i t}}{\lambda_i - R_T} - \frac{e^{-\lambda_p t} - e^{-\lambda_i t}}{\lambda_i - \lambda_p}\right],
\end{aligned} \tag{4.2}$$

where,

- N_T: the number of target nuclides,
- N_i: the number of produced nuclides of interest,
- N_p: the number of parent nuclides which decay to nuclide i,
- N_0: $= N_T(t=0)$, the initial number of target nuclides,
- λ_i: total decay constant per atom for nuclide i, where $\lambda_i = \ln 2/T_{i,1/2}$, with $T_{i,1/2}$ the half life. λ can be interpreted as the probability per unit time for a single atom to decay and has a dimension of s^{-1}.
- λ_{pi}: decay constant per atom for nuclide p decaying to nuclide i,
- R_T: total reaction constant per atom for target nuclide T, which can be interpreted as the probability per unit time that a single target atom undergoes a reaction, and has a dimension of s^{-1},
- R_{Ti}: reaction constant per atom for the reaction of target nuclide T to nuclide i.

In the above equations, the total decay constant for nuclide i is the sum over the partial decay constants of all K daughter nuclides

$$\lambda_i = \sum_{k=1,k\neq i}^{K} \lambda_{ik}. \tag{4.3}$$

The λ_i in the loss term usually feeds only one or two different channels, namely by beta decay to the ground state or isomer of the daughter nuclide d,

$$\lambda_i = \lambda_{id} = \lambda_{id}^{g.s.} + \lambda_{id}^{iso}. \tag{4.4}$$

The total nuclear reaction constant per atom for nuclide T is

$$R_T = \sum_{k=1,k\neq T}^{K} R_{Tk} \tag{4.5}$$

Eq. (4.2), which is derived in the Appendix, is more general than what is used and required in most activation analyses of radioisotope production. For many cases of interest the only term that is used is that of a radioactive nuclide which is directly produced from the target and which decays itself,

$$N_i(t) = N_0 \frac{R_{Ti}}{\lambda_i - R_T} [e^{-R_T t} - e^{-\lambda_i t}] \tag{4.6}$$

The reaction constant R_T is often very small compared to the irradiation time, of the order of 10^{-10} s^{-1}, i.e. $N_T(t) \sim N_0$ and therefore one often discards the first line of Eq. (4.2). For very high burn-up it is better to include this though. With a negligible R_T, the entire Eq. (4.6) would then be further reduced to

$$N_i(t) = N_0 \frac{R_{Ti}}{\lambda_i}[1 - e^{-\lambda_i t}], \tag{4.7}$$

which is the one used in many analyses.

We have however implemented the full set of Eq. (4.2), which considers, with the first line, the burn-up of the target, and also, with the second and third lines, the possible production of the nuclide of interest via an indirect decay route ('parent').

If we neglect the formation via a parent isotope, the irradiation time for which a maximal yield is obtained can be derived by setting the derivative of $N_i(t)$ in Eq. (4.6) to zero, leading to

$$t_{max} = \frac{ln(\lambda_i/R_T)}{\lambda_i - R_T}. \tag{4.8}$$

The number of target atoms available for activation at t=0, N_0, for $1 \leq m \leq M$ (with M the number of natural isotopes constituting the target element) equals:

$$N_0 = \frac{N_A}{A} B_m M_{tar}, \tag{4.9}$$

where $N_A = 6.022\ 10^{23}$ is Avogadro's number, A is the mass number, B_m is the abundance of isotope m with $\sum_m^M B_m = 1$, and M_{tar} is the mass of the target in [g], obtained by the product of the mass density ρ in [g/cm^3] and the volume of the target V_{tar} in [cm^3],

$$M_{tar} = \rho\ V_{tar}, \tag{4.10}$$

The formalism given above is still general with respect to the irradiation source and applies to neutron, charged-particle or photon irradiation. It is the reaction constant R which differs for the three types of irradiation. It depends on two important ingredients: the particular geometry-dependent aspects of the irradiation, and the cross sections. It is clear that nuclear data, both for nuclear decay that determines λ, and the reaction cross sections that determine R, are the critical ingredient in the formalism, and we will discuss that later.

4.2 Reaction constants and rates

For an external spectrum, like that of a photon or neutron source, the reaction constant per atom for the production of an isotope i is the product of the number of source particles, also known as the total particle flux, ϕ^{tot} given in $[cm^{-2}s^{-1}]$ and an average of the residual production cross section σ_i over the normalized energy-dependent particle source spectrum f,

$$R_{Ti} = \phi^{\text{tot}} \int f(E)\sigma_i(E)dE \tag{4.11}$$

The integral is also known as spectrum-averaged cross section or sometimes the effective cross section,

$$< \sigma_i >= \int f(E)\sigma_i(E)dE, \tag{4.12}$$

where,

$$\int f(E)dE = 1. \tag{4.13}$$

This leads to the compact formula

$$R_{Ti} = \phi^{\text{tot}} < \sigma_i > . \tag{4.14}$$

Analogously, for the target the reaction constant in [s^{-1}] of all reaction channels, is given by:

$$R_T = \phi^{\text{tot}}(< \sigma_{non} > - < \sigma_{in} >) \tag{4.15}$$

where $< \sigma_{non} >$ is the spectra-averaged non-elastic cross section and $< \sigma_{in} >$ is the spectrum-averaged inelastic cross section. The difference $< \sigma_{non} > - < \sigma_{in} >$ provides the probability to create an nuclide different from the original target nuclide in the nuclear reaction.

As we see below, for charged-particle irradiation the expression is different as the spectrum is generated inside the target.

For typical irradiation sources like a research reactor or an accelerator the total flux is of the order of 10^{12-14} particles per $[cm^{-2}s^{-1}]$. Note that σ needs to be expressed in 10^{-24} cm for 1 barn, leading to the small absolute values for R. For cross sections in the range of 1 mb to 1 barn, one often obtains reaction constants of the order of R $\sim 10^{-10} - 10^{-13}s^{-1}$.

The reaction constant only depends on the intensity of the incident particle source and not on the target volume of irradiation. This means it can be consistently compared among different types of particles sources and also with the decay constant: the higher the particle flux, the larger the production of the residual isotope relative to its decay. In the production and decay equations (4.2) the reaction constant compete with radioactive decay constants with typical values of $\lambda \sim 10^{-2} - 10^{-4}s^{-1}$ for short lived nuclides $\lambda \sim 10^{-8} - 10^{-12}s^{-1}$ for long lived nuclides.

4.2.1 Neutron irradiation

In general, medical isotope production via neutron reactions takes place in research reactors (with isotope production as an important side application) or dedicated isotope production reactors. Promising alternatives exist, such as using 14 MeV peaked spectra, but those are used at a smaller scale or still in development. Here we give access to all possible estimated neutron spectra that are to our knowledge available. Over the years, many different neutron spectra have been compiled in the CONDERC database, for research reactors, power reactions, fusion designs etc. Hence, inserting such a normalized spectrum in Eq. (4.12) and providing the total flux in Eq. (4.14) will lead directly to the production rate.

There are usually 2 refinements required, which will reduce the calculated activity relative to Eq. (4.11),

- The flux may not be homogeneous due to neutron absorbing materials in neigboring positions in the reactor core, this means that irradiation-specific estimates of the neutron spectrum need to be made, e.g. with a code like MCNP.
- Self-shielding

Regarding the first issue, it is difficult to give a general prescription for a specific irradiation condition, so we assume here that the flux is the same in the entire reactor core.

Self shielding

For self-shielding a reasonable estimate can be given. Eq. (4.11) is then replaced by

$$R_{Ti} = \phi^{\text{tot}} \int G(E) f(E) \sigma_i(E) dE \tag{4.16}$$

where the shelf-shielding factor is given by

$$G(E) = \frac{1 - e^{-\Sigma(E).x}}{\Sigma(E).x} \tag{4.17}$$

where x is the thickness of the material and Σ macroscopic cross section

$$\Sigma(E) = n_T \sigma_{att} \tag{4.18}$$

where for the attenuation cross section σ_{att} we take the total cross section σ_{tot}. To see the overall impact of self shielding it is helpful to define the self-shielding factor averaged over the reactor spectrum,

$$G_{av} = \int G(E) f(E) dE \tag{4.19}$$

4.2.2 Photon irradiation

Photonuclear reactions are often induced by Bremsstrahlung spectra. Various models of increasing sophistication for such a spectrum exist, from simple and more refined analytical models to spectra created by a full Monte Carlo simulation by codes like GEANT4. Here we use a thick-target Kramer spectrum for $f(E)$ in Eq. (4.11).

4.2.3 Charged-particle irradiation

For charged-particle reactions, the expression for the reaction constant is somewhat different from that of Eq. (4.11) since the spectrum is generated inside the target,

$$R_{Ti} = n_p \int_{E_{\text{back}}}^{E_{\text{beam}}} \frac{\sigma_i(E)}{S(E)} dE \tag{4.20}$$

where E_{beam} denotes the incident beam energy and E_{back} is the average particle energy leaving the backside of the target. If the projectiles travels through the target, the average projectile beam energy will decrease and the energy loss inside the target is determined by the stopping power in S [MeV/cm], which will be described later. n_p is the projectile rate density in [particles/(cm^{3}s)],

$$n_p = N_{\text{proj}}/V_{\text{tar}} \tag{4.21}$$

where V_{tar} is the active target volume in [cm^3], and N_{proj} is the number of projectiles impinging on the target per [s], given by

$$N_{\text{proj}} = I_{\text{beam}}/(z_p q_e) \tag{4.22}$$

with I_{beam} the beam current in [A], z_p the charge number of the projectile and q_e the electron charge. The effective path length in [cm] is

$$L_{\text{eff}} = \int_{E_{\text{back}}}^{E_{\text{beam}}} \frac{1}{S(E)} dE, \tag{4.23}$$

The integration limits E_{beam} and E_{back} determine the effective path length and vice versa. V_{tar} is then given by the product of the beam surface S_{beam} in [cm^2] and the effective target length,

$$V_{tar} = S_{beam} L_{\text{eff}} \tag{4.24}$$

Eq. (4.20) can be rewritten in terms of the general formula for R, Eq. (4.11), by defining the spectrum function

$$g(E) = \frac{1/S(E)}{\int_{E_{\text{back}}}^{E_{\text{beam}}} \frac{1}{S(E)} dE}, \tag{4.25}$$

where,

$$\int g(E) dE = 1. \tag{4.26}$$

Then, we obtain

$$R_{Ti} = \frac{N_{proj}}{V_{tar}} L_{\text{eff}} \int g(E)\sigma_i(E)dE \tag{4.27}$$

or

$$R_{Ti} = \frac{N_{proj}}{V_{tar}} L_{\text{eff}} < \sigma_i >_S \tag{4.28}$$

where we use the subscript S to denote that the cross section is averaged over the spectrum associated with the stopping power.

Stopping power

The stopping power describes the average energy loss of projectiles in the target by atomic collisions as a function of their energy in [MeV/cm]. We use the Bethe-Bloch formula [7]:

$$S(E) = \frac{dE}{dx} = 0.1535\, \rho\, \frac{Z}{A}\, \frac{z_p^2}{\beta^2}\, [ln\left(\frac{W_{max}^2}{I^2}\right) - 2\beta^2], \ \text{MeV/cm} \tag{4.29}$$

with Z the target charge number, and z_p the projectile charge number, while β represents a beam particle traveling at a relative velocity

$$\beta = \frac{v}{c} = \sqrt{\frac{E_{beam}(E_{beam} + 2m_0c^2)}{(E_{beam} + m_0c^2)^2}}, \tag{4.30}$$

with rest mass m_0. Further, m_e is the electron mass, and $\gamma = 1/\sqrt{1-\beta^2}$. The maximum energy transfer in a single collision, W_{max} is given by

$$W_{max} = 2m_ec^2(\beta\gamma)^2, \tag{4.31}$$

if the incident particle is much heavier than the electron mass. For the mean excitation potential I a semi-empirical formula is adopted:

$$\frac{I}{Z} = 9.76 + 58.8Z^{-1.19} \ \text{eV}. \tag{4.32}$$

This expression is claimed to be tested only if $Z \geq 13$, but we use it for all values.

Nuclear reaction and decay rates

With this, all ingredients to calculate the nuclide inventory $N_i(t)$ at any time during or after the irradiation process are defined, with exception of the residual production cross sections and decay data which we handle in a separate Section.

When including the target volume, we can introduce the reaction rate The total production rate of nuclide j in [atoms/s] is obtained by multiplying the reaction constant with the number of target atoms:

$$P_{Tj}(t) = R_{Tj}N_T(t). \tag{4.33}$$

and similarly the decay rate

4.3 Activities

With $N_i(t)$ known, the expression for the activity A_i of the produced nuclide i as a function of the irradiation time can now be given by

$$A_i(t) = \lambda_i \, N_i(t). \tag{4.34}$$

Usually the activity is given in GBq or MBq. If the half-life of the produced isotope is large compared to the irradiation time, the expression for $N_i(t)$ simplifies to,

$$N_i(t) = N_0 \, R_{Ti} \, t, \tag{4.35}$$

and hence the activity of i as

$$A_i(t) = \lambda_i \, N_0 \, R_{Ti} \, t, \tag{4.36}$$

In that case, the yield scales linearly with the irradiation time t and the reaction constant R_{Ti}, and the production yield expressed in e.g. [GBq/mAh] for accelerators or the average production rate in [GBq/day] for reactors, is a meaningful quantity.

4.4 Possible refinements

The formalism and ingredients specified above give a reasonable to good estimate of isotope production, which is the first aim of this project. Obviously, several improvements are possible which may lead to better estimates. For all irradiation sources these are:

- A better assessment of the cross sections. The unertainty of these directly proapagate to the estimated yield.
- More accurate decay data. Here, we use the JEFF radioactive decay data library. A combination of the best available decay data libraries may improve the estimated yield and impurities.

In particular for neutron sources,

- A reactor-geometry dependent flux, in which more or less absorbing positions in the reactor are taken into account. Related to this a user-specific reactor spectrum would be used.
- The particular geometry of the target to be irradiated, iand the associated consequence for self shielding

In particular for photon sources,

- A better model for the Bremsstrahlungs spectrum
- A better estimate for the fraction of photons, after their conversion from the electrom beam, which are available for target irradiation

In particular for charged particle sources,

- A better model for the stopping power

5. Nuclear data libraries

In the previous chapter it was outlined that the entire simulation of isotope production basically depends on a few ingredients,

- the solution of the production and decay equations, Eq. (4.2),
- Depending on the activation method:
  - for charged-particle induced activation: a model for the stopping power, in our case Eq. (4.29),
  - for photonuclear induced activation: a model for the Bremsstrahlung spectrum,
  - for neutron-induced induced activation: a model the neutron spectrum,
- a radioactive decay data library,
- a cross section data library,

5.1 Radioactive decay data library

For the radioactive decay data library, we have taken the JEFF-3.3-RD. For the current version of ISOTOPIA, all we use is a basic decay mode, i.e. whether the nuclide decays via electron or positron emission along the decay chain, and of course the half life.

5.2 Cross section data library

5.2.1 Neutrons

We use the TENDL-2025 nuclear data library in pointwise format.

5.2.2 Photonuclear

We use the TENDL-2023 nuclear data library.

5.2.3 Charged particles

The cross section database of ISOTOPIA is a merger of two sources,

- TENDL-2025 [1], is a complete nuclear data library for all projectile, target, product, energy combinations. For projectiles on any target, cross sections are included for all residual products which can be produced for incident energies up to 200 MeV. For incident charged-particles, TENDL has been automatically produced, using adjusted nuclear model parameters for those reactions where experimental data are available. That means that compared to experimental data the TENDL curves are usually good. Statistical analyses of the overall quality, also in comparison with other high-energy libraries, exist.
- The IAEA Medical Isotope database [3, 4, 5, 6], has been produced by a series of IAEA collaborations and has resulted in a recommended database for about 150 cross section sets for important medical isotopes produced by charged-particle induced reactions. For these reactions, the experimental data were considered to be so abundant and/or of so high quality that empirical Pade fits were made to produce the final evaluation, and no nuclear model code was required. This has led to excitation functions for restricted, though often the most important, nuclides and energy ranges.

For charged-particles the medical isotope data library for ISOTOPIA is the result of a smooth merger between these two data libraries. Obviously, TENDL is used for projectiles, targets, products and reaction channels which do not exist in the IAEA database. For the 150 reaction channels of the IAEA database, we have complemented the IAEA database outside its energy range by simply normalizing TENDL to the first and last energy point of the IAEA excitation functions. Often, but not always(!), the most relevant energy range is considered by the IAEA tables. Of course, for totally new production routes only the TENDL cross sections are available. Also, though for the about 150 well-known reaction channels a nuclear model estimate may be redundant, the better a nuclear model code can reproduce the experimental data or its Pade fit, the more confidence we may have for the *other* produced channels for the same reaction, which could represent important contaminations to consider.

The combination of IAEA database and TENDL currently may bring the most complete and best quality library for medical isotope production, but may be improved by adjusting TALYS nuclear model parameters to (even) better fit experimental data for many nuclides.

5.3 Creation of medical isotope data library

The nuclear data library that is used by ISOTOPIA is called *isotopia.libs*. For future reference, we describe here how it is created.

- First, the "raw" data file from the IAEA medical isotope database are renamed to filenames consistent with the entire system around TALYS. For example, the ^{100}Mo(p,2n)^{99m}Tc cross section is stored in p-Mo100-rp043099m.iaea.med. This is a manual or semi-automated step.
- Next, we rename the headers of the files to bring them in sync with the rest of the *libraries/* database. For this we use the script /isotopia/misc/medrename.bash
- To unify with the rest of the database even more, we extend every IAEA file up to 200 MeV using a smooth transition with TENDL, for which we use /isotopia/misc/medmerge.bash.
- Note that there are various collections of historic IAEA files, for therapeutic, diagnostic, monitor reactions etc. They all get the treatment mentioned above, and older files are overwritten with newer files.
- In directory *isotopia.libs* there is a script *create.isotopia.libs*. This script basically merges

the latest version of TENDL with the collection of specific IAEA medical isotope files. In practice this goes as follows: every single reaction file from TENDL is copied, and only those files which exist in the IAEA database are overwritten by the latter. For completeness, also all data from the EXFOR database are included.

This process leads to a data library *isotopia.libs* which has at the highest level the incident particles,

```
p/
d/
t/
h/
a/
```

The next level contains the nuclides

```
....
p/Mo097/
p/Mo098/
p/Mo100/
etc.
```

The next level contains the nuclear data library and EXFOR,

```
p/Mo100/exfor/
p/Mo100/iaea.2024/
```

All relevant EXFOR data that is available in computational XC4 format is stored in the exfor/ directory. For the particular case of medical isotope production we need only 2 subdirectories

```
p/Mo100/exfor/xs      : the channel cross sections, in particular the non-elastic
p/Mo100/exfor/residual: all residual production cross sections
```

Next, the *iaea* directory contains the evaluated nuclear data tabulated per reaction channel in simple tables. For consistency with the *libraries/* database used for other projects we have kept the same directory structure. For *isotopia.libs*, we have

```
p/Mo100/iaea.2024/tables/xs      : the non-elastic cross section p-Mo100-MT005.iaea.2024
p/Mo100/iaea.2024/tables/residual: all residual prod. c.s. e.g. p-Mo100-rp043099m.iaea.2024
```

Finally, as an example of a file with residual production cross sections, *p-Mo100-rp043099m.iaea.2024* looks as follows

```
# header:
#   title: Mo100(p,x)Tc99m cross section
#   source: IAEA medical isotope consortium
#   user: Arjan Koning
#   date: 2024-10-06
#   format: YANDF-0.2
# endf:
#   library: iaea.2024 + TENDL-2023
#   author: IAEA medical isotope consortium
#   year: 2024
```

```
# target:
#   Z: 42
#   A: 100
#   nuclide: Mo100
# reaction:
#   type: (p,x)
#   ENDF_MF: 6
#   ENDF_MT: 5
# residual:
#   Z: 43
#   A: 99
#   nuclide: Tc99m
#   level:
#     isomer: 1
# datablock:
#   quantity: cross section
#   columns: 3
#   entries:     457
##       E             xs            dxs
##     [MeV]          [mb]           [mb]
   6.000000E+00   1.100000E+00   1.100000E+00
   6.100000E+00   9.000000E-01   9.000000E-01
   6.200000E+00   8.000000E-01   8.000000E-01
   6.300000E+00   8.000000E-01   8.000000E-01
..................
```

where this particular example contains data from the IAEA evaluation, extended with TENDL. By far, the majority of files come straight from TENDL, as e.g. *p-Mo100-rp043098.iaea.2024*

```
# header:
#   title: Mo100(p,x)Tc98 cross section
#   source: ENDF
#   user: Arjan Koning
#   date: 2024-10-07
#   format: YANDF-0.2
# endf:
#   library: tendl.2023
#   author: A.J. Koning
#   year: 2023
# target:
#   Z: 42
#   A: 100
#   nuclide: Mo100
# reaction:
#   type: (p,x)
#   ENDF_MF: 6
#   ENDF_MT: 5
```

```
# residual:
#   Z: 43
#   A: 98
#   nuclide: Tc98
# datablock:
#   quantity: cross section
#   columns: 2
#   entries: 31
##       E             xs
##     [MeV]          [mb]
   1.685404E+01   0.000000E+00
   1.700000E+01   6.829083E-02
   1.800000E+01   3.388542E+01
   1.900000E+01   1.525024E+02
..................
```

In sum, TENDL is used for the energy ranges outside the IAEA evaluations and for all other isotopes.

6. Reference Guide

In this part, all ISOTOPIA keywords will be described, one per page. The description of each keyword is as follows:

- Name of the keyword
- Explanation
- Examples
- Range of allowed values
- Default value
- Comments (optional), when we feel that some extra warnings or explanation for proper use is appropriate.

In principle, all keywords may be referred to in the other parts of this tutorial.

6.1 ISOTOPIA keywords

As explained above, the minimum input file has 3 keywords, and leaves all choices for the other parameters to the author. In general, you probably want to be more specific. Below, we will explain all the possible keywords. For each keyword, we give an explanation, a few examples, the default value, and the theoretically allowed numerical range. Remember that you can always find all adopted default values for all parameters at the top of the standard *output* file of ISOTOPIA.

abundance

File with tabulated abundances. The **abundance** keyword is only active for the case of a natural target, i.e. if **mass 0**. By default, the isotopic abundances are read from the *isotopia/files/abundance/* database. It can however be imagined that one wants to analyze production from targets of a certain isotopic enrichment. On the input line, we read **abundance** and the filename. From each line of the file, TALYS reads Z, A and the isotopic abundance with the format (2i4,f11.6). An example of an abundance file, e.g. *abnew*, different from that of the database, is

```
82 206  24.100000
82 207  22.100000
82 208  52.400000
```

where we have left out the "unimportant" ^{204}Pb (1.4%). ISOTOPIA automatically normalizes the abundances to a sum of 1, leading in the above case to 24.44 % of ^{206}Pb, 22.41 % of ^{207}Pb and 53.14 % of ^{208}Pb in the actual calculation.

Examples

abundance abnew

Range

abundance can be equal to any filename, provided it is present in the working directory.

Default

If **abundance** is not given in the input file, abundances are taken from *files/abundance* and calculations for all isotopes are performed.

Adepth

The depth, relative to the target mass number, to which isotopes are considered for cross sections that contribute to the production path. For example if **Adepth 5** for the p + ^{100}Mo reaction, only yields for elements defined by **Zdepth** and isotopes with mass numbers between 95 and 101 will be calculated.

Examples

Adepth 2
Adepth 10

Range

0 $\leq$ **Adepth** $\leq$ **Atarget** where Atarget is the mass number of the target.

Default

Adepth 20.

Area

Area of the target in cm^2.

Examples

Area 1o.

Range

0 $<=$ **Area** $<=$ **10000**.

Default

Area 1. cm^2.

crosspath

Path for the cross section database. You should have this path hardwired in subroutine *machine.f*, but it may be helpful to easily change between different versions of the cross section database.

Examples

crosspath /home/koning/libraries
crosspath /home/qaim/tendl.2019

Range

crosspath should exist.

Default

Default: **crosspath ∼/isotopia.libs/**

currentunit

Unit for accelerator power.

Examples

currentunit mua: micro-Ampere
currentunit ma: milli-Ampere
currentunit A: Ampere

Range

currentunit should be equal to one of the units above.

Default

currentunit ma.

decay

Flag to include decay from parent nuclide. Basically used for code diagnostics.

Examples

decay y
decay n

Range

y or **n**

Default

decay y

Eback

For charged particle irradiation only: The lower end of the energy range at the back end of the target in MeV. This energy degradation is directly related to the effective thickness of the target.

Examples

Eback 130.
Eback 12.

Range

0. $\leq$ **Eback** $<$ **Ebeam** and 0. $\leq$ **Eback** $<$ 250. MeV

Default

Eback = Ebeam - 5 MeV.

Ebeam

For charged particle irradiation only: The incident energy of the particle beam in MeV.

Examples

Ebeam 140.
Ebeam 16.

Range

$0 <$ **Ebeam** $< 250.$ MeV.

Default

None.

element

Either the nuclear symbol or the charge number Z of the target nucleus can be given. Possible values for **element** range from Li (3) to C4 (124).

Examples

element pu
element 41
element V
element B9

Range

3 $\leq$ **element** $\leq$ **124** or **Li** $\leq$ **element** $\leq$ **C4**.

Default

None.

Comments

- To accommodate target nuclides with $Z > 110$ the element names are defined as follows: Rg(111), Cn(112), Nh(113), Fl(114), Mc(115), Lv(116), Ts(117), Og(118), B9(119), C0-4(120-124), which takes as a basis for $Z = 100$ the symbol $A0$, etc. unless an official name has been assigned to it, which is currently the case for $Z \leq 118$. Clearly, if we ever need or wish to go beyond $Z = 124$, there are enough symbols left. Obviously the symbols for Z above 118 will be changed as soon as official names are assigned to them.

fluxtotal

For neutrons and photons: Total particle flux in [$cm^{-}2s^{-}1$].

Examples

fluxtotal 2.e14
fluxtotal 5.e13

Range

0 $<=$ **fluxtotal** $<=$ **1e38**.

Default

fluxtotal 1.e14.

format

The format of the metadata in the output files. It may be helpful to specify the version. You may edit the 'format' variable variable in input.f90 and put your own version there. However, you can also use the input keyword.

Examples

format YANDF-0.5beta2

Range

Any string

Default

YANDF-0.4

Ibeam

For charged particles: Particle beam current in the unit of **currentunit** (which has default mA).

Examples

Ibeam 0.1
Ibeam 5.

Range

0 $<=$ **Ibeam** $<=$ **10000**.

Default

Ibeam 1.

mass

The target mass number A. The case of a natural element can be specified by **mass 0**. Then, an ISOTOPIA calculation for each naturally occurring isotope will be performed (see also the **abundance** keyword, p. 32), after which the results will be properly weighted and summed.

Examples

mass 112
mass 0

Range

mass 0 or **5** $<$ **mass** $\leq$ **339**. The extra condition is that the target nucleus, i.e. the combination of the target **mass** and **element** must be present in the cross section database, which corresponds to all nuclei with a half life longer than 1 day.

Default

None.

massunit

Unit for isotope yield (to give relative production rate) or mass unit, in weight.

Examples

massunit mug: microgram
massunit mg: milligram
massunit g: gram
massunit kg: kilogram

Range

massunit should be equal to one of the units above.

Default

massunit g.

outcross

Flag to output cross sections to separate files.

Examples

outcross y
outcross n

Range

y or **n**

Default

outcross n

projectile

Seven different symbols can be given as **projectile**, namely **n, g, p, d, t, h, a** representing neutron, gamma (photon), proton, deuteron, triton, ^{3}He, and alpha, respectively.

Examples

projectile n
projectile g
projectile p
projectile d

Range

projectile must be equal to **n, g, p, d, t, h, a**.

Default

None.

radiounit

Unit for radioactivity, to be used in the output files for isotope production.

Examples

radiounit Bq: Becquerel
radiounit kBq: kiloBecquerel
radiounit MBq: MegaBecquerel
radiounit GBq: GigaBecquerel
radiounit Ci: Curie
radiounit mCi: milliCurie
radiounit kCi: kiloCurie

Range

radiounit should be equal to one of the units above.

Default

radiounit Bq.

rho

Material density of the target in g/cm^3.

Examples

rho 10.

Range

0 $<=$ **rho** $<=$ **100**.

Default

rho is read from a hardwired material density table.

selfshield

For neutron sources: flag to take into account self shielding.

Examples

selfshield y
selfshield n

Range

y or **n**

Default

selfshield n

source

The source of the calculations. obviously this is ISOTOPIA, but it may be helpful to specify the version such that it appears in the metadata of the many output files. You may edit the 'source' variable variable in input.f90 and put your own version there. However, you can also use the input keyword.

Examples

source ISOTOPIAbeta3v20260223

Range

Any string

Default

ISOTOPIA

targetmass

For neutrons or photons: Mass of the target in the **massunit** unit (default: g).

Examples

targetmass 0.1
targetmass 5.

Range

0 $<=$ **targetmass** $<=$ **1e9**.

Default

targetmass is not used but calculated from **rho** and volume of target.

thickness

For neutrons or photons: thickness of the target in cm. For protons this is calculated from the stopping power.

Examples

thickness 0.1
thickness 5.

Range

0 <= **thickness** <= **10000** cm.

Default

thickness 0.05 cm.

timeunit

Unit for time for irradiation, cooling or relative production rate.

Examples

timeunit s: second
timeunit h: hour
timeunit d: day

Range

timeunit should be equal to one of the units above.

Default

timeunit h.

Tirrad

Irradiation time. A general input for the irradiation time has been enabled. On the input line we read integer values and time units, which can be y (years), d (days), h (hours), m (minutes) or s (seconds). These all need to be separated by blanks.

Examples

Tirrad	**2 d 5 h**
Tirrad	**32 h 30 m**
Tirrad	**1 d 6 h 24 m 12 s**

Range

0 <= **Tirrad** <= **1e6** for every time unit.

Default

Tirrad 1 d.

Tcool

Target cooling time. A general input for the cooling time has been enabled. On the input line we read integer values and time units, which can be y (years), d (days), h (hours), m (minutes) or s (seconds). These all need to be separated by blanks.

Examples

Tcool	**2 d 5 h**
Tcool	**32 h 30 m**
Tcool	**1 d 6 m 24 m 12 s**

Range

0 <= **Tcool** <= **1e6** for every time unit.

Default

Tcool 1 d.

user

The current user who produces the result. You may edit the 'user' variable in input.f90 and put your own name there. It will appear in the metadata of the various output files. However, you can also use the input keyword.

Examples

user Vlad Avrigeanu

Range

Any string

Default

Arjan Koning

xsfile

Local file with cross sections. With this keyword, residual production cross sections from TENDL can be overruled with values given in your working directory. On the input line, we read **xsfile**, Z and A of the residual product, the filename, and optionally the number of the produced isomer. This file should consist of Energy [MeV] - cross section [mb] values in x-y form, with a similar structure as those of the cross section database.

Examples

xsfile 25 56 Mn56.exp
xsfile 42 100 Mo100.loc 1

Range

xsfile should exist.

Default

None.

yieldunit

Unit for isotope yield (to give relative production rate) or mass unit, in weight.

Examples

yieldunit mug: microgram
yieldunit mg: milligram
yieldunit g: gram
yieldunit kg: kilogram

Range

yieldunit should be equal to one of the units above.

Default

yieldunit g.

ZAoutput

Flag to use numerical Z and A values in the filenames instead of nuclide names.

Examples

ZAoutput y
ZAoutput n

Range

y or **n**

Default

ZAoutput n

Zdepth

The depth, relative to the target charge number, to which isotopes are considered for cross sections that contribute to the production path. For example if **Zdepth 2** for the p + Mo reaction, only yields for Tc, Mo, Nb and Zr isotopes will be calculated.

Examples

Zdepth 2
Zdepth 10

Range

0 $\leq$ **Zdepth** $\leq$ **Ztarget** where Ztarget is the mass number of the target.

Default

Zdepth 10.

Table 6.1: The keywords of ISOTOPIA.

Keyword	Range	Default	Page
abundance	filename	no default	32
Adepth	0-Atarget	20	33
Area	0. - 10000.	1.	34
crosspath	directory name	~/isotopia.libs/	35
currentunit	muA, mA, or A	mA	36
decay	y,n	y	37
Eback	1.e-11 - 250.	Ebeam - 5	38
Ebeam	1.e-11 - 250.	None	39
element	3 - 124 or Li - C4	None	40
fluxtotal	0. - 1.e38	1.e14	41
format	format name	YANDF-0.4	42
Ibeam	0. - 10000.	1.	43
mass	0, 5 - 339	None	44
massunit	mug, mg, g or kg	g	45
outcross	y,n	n	46
projectile	n,g,p,d,t,h,a,	None	47
radiounit	(k,M,G)Bq, (m,k)Ci	Bq	48
rho	0. - 100.	None	49
selfshield	y,n	n	50
source	name of source	ISOTOPIA	51
targetmass	0. - 1.e9	None	52
thickness	0. - 10000.	0.05	53
timeunit	s, h or d	h	54
Tirrad	0 - 1.e6	1 d	55
Tcool	0 - 1.e6	1 d	56
user	your name	Arjan Koning	57
xsfile	file name	none	58
yieldunit	mug, mg, g or kg	g	59
ZAoutput	y,n	n	60
Zdepth	0-Ztarget	10	61

7. Sample cases

The purpose of ISOTOPIA is that every imaginable production route for a medical isotope can be simulated. To strengthen this statement, we will discuss a few sample cases. In each case, the ISOTOPIA input file, the relevant output and if appropriate, a figure will be presented. The description of the first sample case is the longest, since the output of ISOTOPIA will be discussed in complete detail. Obviously, that output description is also applicable to the other sample cases. The entire collection of sample cases serves as (a) verification of ISOTOPIA: the sample output files should coincide, apart from numerical differences of insignificant order, with the output files obtained by you, and (b) validation of ISOTOPIA: the results should be comparable to those obtained in other publications. The computation time of ISOTOPIA is very small. In general, we will explain the keywords again as they appear in the input files below. If not, consult Table 6.1, which will tell you where to find the explanation of a keyword.

7.1 Sample p-Mo100-Tc099m: Accelerator production of ^{99m}Tc

This sample case considers one of the most discussed alternatives for reactor-produced ^{99}Mo: the direct production of ^{99m}Tc from ^{100}Mo with a proton accelerator via the (p,2n) reaction. The input file for this case is as follows,

```
projectile p
element Mo
mass 100
Ebeam 16.
Eback 12.
Ibeam 0.15
```

Running this input file with

isotopia < isotopia.inp > isotopia.out

gives an output file with the general characteristics of the reaction. The output file starts with a display of the version of ISOTOPIA you are using, the name of the author, the Copyright statement and then an exact copy of the input file as given by the user is returned.

```
   ISOTOPIA-2.1    (Version: December 29, 2024)

           Prediction of medical isotope production with accelerators

Copyright (C) 2024  A.J. Koning

User: Arjan Koning
Date: 2024-12-22

########## USER INPUT ##########

USER INPUT FILE

projectile p
element mo
mass 100
ebeam 16.
eback 12.
ibeam 0.15

USER INPUT FILE + DEFAULTS

Keyword           Value   Variable     Explanation

user      Arjan Koning    user         user for this calculation
source    ISOTOPIA        source       source for this calculation
format    YANDF-0.2       oformat      format for output
#
# Nuclear reaction
#
projectile          p     ptype0       type of incident particle
element            Mo     Starget      symbol of target nucleus
mass              100     Atarget      mass number of target nucleus
#
# Accelerator
#
Ebeam              16.000 Ebeam        incident energy in MeV
Eback              12.000 Eback        lower end of energy range in MeV
Ibeam               0.150 Ibeam        beam current in mA
Tirrad              1     Tirrad       d of irradiation time
radiounit             gbq radiounit    unit for radioactivity
yieldunit             num yieldunit    unit for isotope yield
#
# Target
#
```

```
 Area                 1.000 Area         target area in cm^2
 Tcool                1     Tcool        d of cooling time
 rho                 -1.000 rhotarget    target density [g/cm^3]
 #
 # Cross section and decay data
 #
 Zdepth              10     Zdepth       depth to which Z numbers are scanned for cross sections
 Adepth              20     Adepth       depth to which A numbers are scanned for cross sections
```

Not only the keywords that you have specified in the input file are listed, but also all the defaults that are set automatically. The corresponding Fortran variables are also printed, together with a short explanation of their meaning. This table can be helpful as a guide to change further input parameters for a next run. You may also copy and paste the block directly into your next input file.

In the next output block we print the main parameters that characterize the production process. Note that some of these have been given in the input, while others are automatically calculated.

```
 Summary of isotope production for p + Mo100

 Maximal irradiation time:       0 years   1 days  0 hours  0 minutes  0 seconds
 Cooling time:       0 years   1 days  0 hours  0 minutes  0 seconds
 E-beam [MeV]:   1.600000E+01
 E-back [MeV]:   1.200000E+01
 Beam current [mA]:        0.150 mA
 Target material density [g/cm^3]:   1.022000E+01
 Target area [cm^2]:   1.000000E+00
 Effective target thickness [cm]:   2.097696E-02
 Effective target volume [cm^3]:   2.097696E-02
 Effective target mass [g]:   2.143845E-01
 Number of target atoms:    1.291053E+21
 Number of incident particles [s^-1]:   9.362260E+14
 Produced heat in target [kW]:   6.000000E-01
```

The block with final results is

```
 (Maximum) production and decay rates per isotope

 Total production rate [s^-1]:   9.624934E-10

#  Nuc     Production rate Decay rate     Activity       #isotopes      Yield          Isotopic fr
#             [s^-1]         [s^-1]         [GBq]          [   ]        [GBq/mAh]
 Tc 101     9.115965E-13   8.135531E-04   1.176901E+00   1.446618E+12   3.427384E-03   0.00004
 Tc 100     6.088757E-11   4.387007E-02   7.860895E+01   1.791858E+12   3.032753E-01   0.00051
 Tc  99     7.742050E-10   1.026401E-13   2.616365E+02   7.370476E+16   6.840031E-10   0.54037
 Tc  99 g   5.875616E-10   1.026401E-13   6.726828E-06   6.553800E+16   5.190665E-10   0.40869
 Tc  99 m   2.162431E-10   3.203675E-05   2.616365E+02   8.166760E+15   5.800662E-02   0.05093
 Mo 100     1.174375E-10   0.000000E+00   0.000000E+00   1.290946E+21   0.000000E+00   1.00000
..................
  End of ISOTOPIA calculation for p + Mo100
```

The time-dependent production of each isotope is found the files *ZZAAA.act* or *ZZAAAm.act* if is concerns isomeric production, with ZZ the charge number in (i2.2) format and AA the mass number in (i3.3) format.

For this case, obviously most important is the production of ^{99m}Tc, given in file *Tc099m.act*, which looks as follows,

```
# header:
#   title: Mo100(p,x)Tc99m Isotope production
#   source: ISOTOPIA
#   user: Arjan Koning
#   date: 2024-12-22
#   format: YANDF-0.2
# target:
#   Z: 42
#   A: 100
#   nuclide: Mo100
# reaction:
#   type: (p,x)
#   ENDF_MF: 6
#   ENDF_MT: 5
# residual:
#   Z: 43
#   A: 99
#   nuclide: Tc99m
# parameters:
#   Beam current [mA]:  1.500000E-01
#   E-Beam [MeV]:  1.600000E+01
#   E-Back [MeV]:  1.200000E+01
#   Initial production rate [s^-1]:  2.162431E-10
#   Decay rate [s^-1]:  3.203675E-05
#   Initial production yield [GBq/mAh]:  5.800662E-02
#   Total activity at EOI [GBq]:  2.616365E+02
#   Irradiation time:       0 years   1 days  0 hours  0 minutes  0 seconds
#   Cooling time:       0 years   1 days  0 hours  0 minutes  0 seconds
#   Half-life:       0 years   0 days  6 hours  0 minutes 36 seconds
#   Maximum production:       0 years   3 days 18 hours 17 minutes 18 seconds
# datablock:
#   quantity: Isotope production
#   columns: 6
#   entries: 100
##     Time         Activity       Isotopes         Yield     Isotopic_frac.      Time
##     [sec]          [GBq]           []          [GBq/mAh]         []              [h]
   1.728000E+03   1.503532E+01   4.693146E+14   5.800662E-02   1.336947E-01   4.800000E-01
   3.456000E+03   2.926088E+01   9.133534E+14   5.488257E-02   1.306088E-01   9.600000E-01
   5.184000E+03   4.272030E+01   1.333478E+15   5.192677E-02   1.275825E-01   1.440000E+00
   6.912000E+03   5.545484E+01   1.730975E+15   4.913017E-02   1.246355E-01   1.920000E+00
   8.640000E+03   6.750352E+01   2.107065E+15   4.648413E-02   1.217707E-01   2.400000E+00
.....................
```

In the file above, the general irradiation characteristics are repeated, as well as the total production rates. The activity as a function of time is presented in Fig. 7.1. Note that the production yield decreases as a function of time since ^{99m}Tc has a half life of 6 hours. Also the last column in this file can be related to the purity or specific activity of ^{99m}Tc relative to the other Tc isotopes and ^{99c}Tc.

Note that the directory also lists the following files



Figure 7.1: Production of Tc-99m

```
Mo099.act
Mo100.act
Tc099.act
Tc099g.act
Tc099m.act
Tc100.act
Tc101.act
```

which represent the isotopes, of which some can be regarded as 'impurities', which are produced simultaneously.

7.2 Sample a-Bi209-At211: alpha-induced production of therapeutic nuclide ^{211}At

This production route is listed in Table 1 of Ref. [8]. The corresponding input file for this case is

```
projectile a
element Bi
mass 209
Ebeam 28.
Eback 10.
Ibeam 0.100
```

Ref. [8] lists an initial production yield of 17.5 Mbq/(μA.h), while ISOTOPIA produces a value of 22.5, see the output file. Note that there is still some uncertainty with regards to the production cross section.

7.3 Sample p-Ga069-Ge068: Production of the ^{68}Ge generator for ^{68}Ga

There are various production routes for this isotope. The input file for this case is

```
projectile p
element Ga
mass 69
Ebeam 25.
Eback 15.
Tirrad 7 d
radiounit Ci
outcross y
```

Note that in this input file we have include some other features of ISOTOPIA, which of course can also be used in all other input files. The irradiation time is 7 days, while we request the output to be in Ci instead of GBq. In this case, the output file of most interest is probably *Ge068.act*. Also, all residual production cross sections for this reaction are printed in separate files.

7.4 Sample p-Th232-Ac225: Production of ^{225}Ac from ^{232}Th

As an example of the high-energy end of the domain of ISOTOPIA, we include the following sample case,

```
projectile p
element Th
mass 232
Ebeam 192
Eback 170.
Ibeam 0.10
```

Note that activation files for many isotopes are produced. The higher the incident energy, the more side-products are made, and many of them could be unwanted.

8. Outlook and conclusions

This tutorial describes ISOTOPIA, a software package for the prediction of medical isotope production using an accelerator.

The formalism employed to calculate all isotopic activity yields, is based on the direct production and β-decay of the isotopes. Feeding of the isotope through β-decay of other nuclei produced in the nuclear reactions is taken into account for one parent nuclide only, which for activation is generally sufficient. In general, one would like to include the processes of all β-decay feeding as well as nuclear reactions of already produced nuclei (which occurs at a high target burn-up) in an exact calculation. For most cases of interest, this is however not necessary. Nevertheless, we may upgrade ISOTOPIA with a general differential equation solver in the future.

ISOTOPIA is complete in the sense that it produces results for all possible isotope production paths. The quality of the prediction is directly related to the current quality of the cross section database. The backbone of the cross section database is TENDL-2025. TENDL has gradually transformed from a calculated data library into an evaluated data library. This means that in addition to global theoretical improvements in the TALYS code, an increasing number of experimental cross section sets will be taken into account in future versions of TENDL. Until that time, and probably also after, the addition of the IAEA medical isotope database, for about 150 important reaction channels, is indispensable.

In terms of functionality, we can think of at least one obvious extension, which is the ability to answer the "inverse question": *Given a desired isotope*, how do the various production routes relate to each other, in terms of projectile type, energy ranges, impurities, etc. and what is the optimal production route? In the current version of ISOTOPIA, we can only perform the "forward route": on the basis of an intuitively promising production route, we calculate all results of interest. A loop over the entire ISOTOPIA code needs to be built for this extension, after which a clever optimization (search) routine would provide the answer. Given the short calculation time of a typical ISOTOPIA run, this looks feasible. The best is proably to build a full database by running ISOTOPIA over all nuclides and projectiles, and then analyze that database.

Bibliography

[1] A.J. Koning et al. “TENDL: Complete Nuclear Data Library for innovative Nuclear Science and Technology”. In: *Nuclear Data Sheets* 155 (2019), p. 1.

[2] A.J. Koning, S. Hilaire, and S. Goriely. “TALYS: modeling of nuclear reactions”. In: *European Physical Journal A* 59 (2023), p. 131.

[3] E. Betak et al. “Nuclear data for the production of therapeutic radionuclides”. In: *IAEA Technical Reports Series, IAEA, Vienna, Austria (2011) ISBN 978-92-0-115010-3. editors S.M. Qaim, F. Tarkanyi, R. Capote, technical report STI/DOC/010/473* (2011).

[4] A. Hermanne et al. “Reference cross sections for charged-particle monitor reactions”. In: *Nucl. Data Sheets* 148 (2018), pp. 338–382.

[5] F. T. Tarkanyi et al. “Recommended nuclear data for medical radioisotope production: diagnostic positron emitters”. In: *J. Rad.Nucl.Chem.* 319 (2019), pp. 487–531.

[6] J. W. Engle et al. “Recommended Nuclear Data for the Production of Selected Therapeutic Radionuclides”. In: *Nucl. Data Sheets* 155 (2019), pp. 56–74.

[7] W. R. Leo. “Techniques for Nuclear and Particle Physics Experiments”. In: *Springer Verlag, Berlin* (1994).

[8] B. Scholten S.M. Qaim I. Spahn and B. Neumaier. “Uses of alpha particles, especially in nuclear reaction studies and medical radionuclide production”. In: *Radiochimica Acta* 104 (2016), p. 601.

[9] H. Bateman. In: *Proc. Cambridge Phil. Soc.* 15 (1910), p. 423.

[10] W. Rubinson. In: *J. Chem. Phys.* 17 (1949), p. 542.

A. Derivation of the nuclide production equations

A.1 Differential equations for production and depletion of nuclides

The most general situation is that of the irradiation of a piece of material, consisting of many different nuclides, which were either present at the start of the irradiation, have been formed, or will be formed during the irradiation, by either the primary flux of particles, by secondary particles, or by radioactive decay. If we have a total of K different nuclides and the number of each nuclide k is N_k, then the temporal development of such a system, during irradiation and decay, is described by K differential equations:

$$\begin{aligned}
\frac{dN_1(t)}{dt} &= \sum_{k=1,k\neq 1}^{K} \Lambda_{k\to 1} N_k(t) - \Lambda_1 N_1(t) \\
&\dots \\
\frac{dN_i(t)}{dt} &= \sum_{k=1,k\neq i}^{K} \Lambda_{k\to i} N_k(t) - \Lambda_i N_i(t) \\
&\dots \\
\frac{dN_K(t)}{dt} &= \sum_{k=1,k\neq K}^{K} \Lambda_{k\to K} N_k(t) - \Lambda_K N_K(t)
\end{aligned} \tag{A.1}$$

In each equation, the first term is a feeding term. In the most general case, various parent nuclides k may contribute to the formation of nuclide i, hence the summation over k, with $\Lambda_{k\to i}$ the partial formation rate for any possible parent nuclide k to nuclide i. Each partial formation rate can be expressed as

$$\Lambda_{k\to i} = \lambda_{k\to i} + R_{k\to i} \tag{A.2}$$

with $\lambda_{k\to i}$ the (partial) radioactive decay rate and $R_{k\to i}$ the (partial) nuclear reaction rate for any possible parent nuclide k to nuclide i. The second terms of Eq. (A.1) are loss terms due to

radioactive decay and nuclear reactions from nuclide i to any other nuclide. Here the total depletion rate (we interchange "depletion" with "formation" whenever appropriate) for nuclide i is

$$\Lambda_i = \lambda_i + R_i \tag{A.3}$$

where the total decay rate for nuclide i is

$$\lambda_i = \sum_{k=1,k\neq i}^{K} \lambda_{i\to k}$$

and the total nuclear reaction rate for nuclide i is

$$R_i = \sum_{k=1,k\neq i}^{K} R_{i\to k} \tag{A.4}$$

The entire loop over k may run over nuclides in their ground or isomeric states. Theoretically, the sum over reaction rates could include secondary particles (neutrons, photons, alpha particles etc.) formed after the first interaction of the incident beam with the material, over the entire outgoing energy spectrum. Since the number of nuclides i, $N_i(t)$ may appear simultaneously in many equations, due to its possible formation, or depletion, by many different nuclear reactions, it is clear that such a coupled system can only be solved by complicated mathematical and computational techniques. In fact, the most exact simulation would involve a Monte Carlo 3D transport calculation in which all primary and secondary particles are taken into account, including complete cross section libraries for all possible particles, coupled with an activation code that keeps track of the nuclide inventory. If we neglect such thick target transport issues, a system of equations like (A.1) is often solved by methods developed by Bateman [9] for radioactive decay and later generalized for source terms by Rubinson [10].

Fortunately, the situation is often not as complex as sketched above, since very reasonable approximations can be introduced into Eq. (A.1), which represent cases which are used in practice. First, let us start with the common case of the irradiation of a target which contains only one natural element at the start of the irradiation. This set of equations can be then be separated into a linear combination of contributions by each target *isotope*. Hence, we solve Eq. (A.1) one by one for each target isotope and add these contributions at the end to get the activation for the target material. Then, for such a mono-isotopic target T we have at the start of the irradiation

$$\begin{aligned} t=0: N_T &= N_T(0) = N_0 \\ N_i &= 0 \\ &\dots \\ N_K &= 0 \end{aligned} \tag{A.5}$$

Since in practice our target nuclide is not radioactive, the loss terms reduce to $\Lambda_T = R_T$ in Eq. (A.1) for N_T. Next, if a substantial part of the target is converted into other nuclides, beam particles may interact with atoms other than the original target atoms. However, this is only a concern for very long irradiation times. For most practical applications, like medical isotope production, we can often assume that the burn-up of the target is small and that the target composition does not change much during the irradiation (this will be confirmed by some of our sample cases). Hence, in addition we will assume there are no nuclear reaction or radioactive decay feeding terms for the *target* nuclide T.

This means that the temporal evolution of the target nuclide is described by the following simple equation

$$\frac{dN_T(t)}{dt} = -R_T N_T(t) \tag{A.6}$$

which basically states that the only thing that happens to the target material is burn-up through nuclear reactions.

For the nuclides that are produced during the irradiation, which are of course more interesting, we can make similar reasonable assumptions. We assume there is no loss of *produced* nuclides of interest through nuclear reactions with beam particles, i.e. we assume nuclides which are produced are not hit twice (again, this assumption becomes less accurate at very long irradiation times). This is equivalent to stating that the nuclides of interest are only produced from nuclear reactions on the target nuclides or from decay of other products formed during the irradiation. Also, consistent with the assumption for the target nuclide that nuclides are not hit twice, we have $\Lambda_i = \lambda_i$ for the depletion term, and assume that no other nuclear reactions lead to the nuclide of interest. Thus for the produced nuclides i we obtain

$$\frac{dN_i(t)}{dt} = \sum_{k=1,k\neq i}^{K} \lambda_{k\to i} N_k(t) + R_{T\to i} N_T(t) - \lambda_i N_i(t) \tag{A.7}$$

Nuclides often have only one or a few radioactive decay modes. Usually, only beta decay to the ground state or an isomer needs to be considered, although in some cases alpha decay may occur as well. If we neglect alpha decay, the summation in the first term above reduces to only one term. Thus, Eq. (A.7) reduces to

$$\frac{dN_i(t)}{dt} = \lambda_{p\to i} N_p(t) + R_{T\to i} N_T(t) - \lambda_i N_i(t) \tag{A.8}$$

where we now use the subscript p for the parent nuclide which decays to nuclide i. The parent nuclide itself is produced from nuclear reactions on the target and is described by the equation,

$$\frac{dN_p(t)}{dt} = R_{T\to p} N_T(t) - \lambda_p N_p(t) \tag{A.9}$$

where for simplicity we have left out a possible feeding term $\lambda_{g\to p} N_g(t)$ from its own parent (i.e. the grandparent g of nuclide i). It neglects radioactive decay to channels produced by multiple proton emission, e.g. ^{120}Te(p,2p)^{119}Sb and the contribution from its feeding channel ^{120}Te(p,2n)^{119}I + $2\beta^+$. Although the (p,2p) channel is generally smaller and therefore often not relevant for most practical medical isotope production routes, it would be safer that at higher energies feeding from radioactive decay is taken into account. For that a more general solution of Eq. (A.1) would have to be implemented, similar to solutions for neutron-induced problems where processes like e.g. multiple decay of fission fragments needs to be taken into account. For most charged-particle induced reactions of interest, the current approximation is justified.

A.2 Solution of the differential equations for production and depletion

Here we derive the solutions (4.2) of the differential equations (A.6)-(A.9). Eq. (4.2) has been implemented in ISOTOPIA.

For the target nuclide we have

$$\frac{dN_T(t)}{dt} = -R_T N_T(t). \tag{A.10}$$

This is easily integrated to give

$$N_T(t) = N_0\, e^{-R_T t}. \tag{A.11}$$

For every parent nuclide we have

$$\frac{dN_p(t)}{dt} = R_{T\to p} N_T(t) - \lambda_p N_p(t). \tag{A.12}$$

Multiplying both sides with $e^{\lambda_p t}$ gives

$$e^{\lambda_p t}\frac{dN_p(t)}{dt} = e^{\lambda_p t}\left[R_{T\to p} N_T(t) - \lambda_p N_p(t)\right], \tag{A.13}$$

or,

$$e^{\lambda_p t}\frac{dN_p(t)}{dt} + e^{\lambda_p t}\lambda_p N_p(t) = e^{\lambda_p t} R_{T\to p} N_T(t), \tag{A.14}$$

or,

$$\frac{d[e^{\lambda_p t} N_p(t)]}{dt} = e^{\lambda_p t} R_{T\to p} N_T(t). \tag{A.15}$$

Integrating this, after insertion of Eq. (A.11), gives

$$\begin{aligned} e^{\lambda_p t} N_p(t) &= N_0 R_{T\to p} \int dt e^{(\lambda_p - R_T)t} \\ &= \frac{N_0 R_{T\to p}}{\lambda_p - R_T} e^{(\lambda_p - R_T)t} + C, \end{aligned} \tag{A.16}$$

or,

$$N_p(t) = \frac{N_0 R_{T\to p}}{\lambda_p - R_T} e^{-R_T t} + C e^{-\lambda_p t}. \tag{A.17}$$

Inserting the initial condition $N_p(t=0) = 0$ gives

$$0 = \frac{N_0 R_{T\to p}}{\lambda_p - R_T} + C, \tag{A.18}$$

giving the solution for the parent nuclide,

$$N_p(t) = \frac{N_0 R_{T\to p}}{\lambda_p - R_T}\left[e^{-R_T t} - e^{-\lambda_p t}\right]. \tag{A.19}$$

For every nuclide i, directly produced by a nuclear reaction on the target nuclide or from decay of the parent nuclide, we have

$$\frac{dN_i(t)}{dt} = \lambda_{p\to i} N_p(t) + R_{T\to i} N_T(t) - \lambda_i N_i(t). \tag{A.20}$$

Multiplying both sides with $e^{\lambda_i t}$ gives analogous to Eqs. (A.13)-(A.15),

$$\frac{d[e^{\lambda_i t} N_i(t)]}{dt} = e^{\lambda_i t}\left[\lambda_{p\to i} N_p(t) + R_{T\to i} N_T(t)\right], \tag{A.21}$$

or, after insertion of the solutions for N_T, Eq. (A.11), and N_p, Eq. (A.19),

$$\frac{d[e^{\lambda_i t}N_i(t)]}{dt} = e^{\lambda_i t}\left[\lambda_{p\to i}\frac{N_0 R_{T\to p}}{\lambda_p - R_T}(e^{-R_T t} - e^{-\lambda_p t}) + N_0 R_{T\to i}e^{-R_T t}.\right] \tag{A.22}$$

Integrating this gives

$$\begin{aligned} e^{\lambda_i t}N_i(t) &= \frac{N_0\lambda_{p\to i}R_{T\to p}}{\lambda_p - R_T}\left[\int dt e^{(\lambda_i - R_T)t} + \int dt e^{(\lambda_i-\lambda_p)t}\right] \\ &+ N_0 R_{T\to i}\int dt e^{(\lambda_i - R_T)t} + C, \end{aligned} \tag{A.23}$$

or,

$$\begin{aligned} N_i(t) &= \frac{N_0\lambda_{p\to i}R_{T\to p}}{\lambda_p - R_T}\left[\frac{e^{-R_T t}}{\lambda_i - R_T} + \frac{e^{-\lambda_p t}}{\lambda_i - \lambda_p}\right] \\ &+ \frac{N_0 R_{T\to i}}{\lambda_i - R_T}e^{-R_T t} + Ce^{-\lambda_i t}. \end{aligned} \tag{A.24}$$

Inserting the initial condition $N_i(t=0) = 0$ gives

$$0 = \frac{N_0\lambda_{p\to i}R_{T\to p}}{\lambda_p - R_T}\left[\frac{1}{\lambda_p - R_T} + \frac{1}{\lambda_i - \lambda_p}\right] + \frac{N_0 R_{T\to i}}{\lambda_i - R_T} + C, \tag{A.25}$$

so we finally obtain

$$N_i(t) = N_0\,\frac{R_{T\to i}}{\lambda_i - R_T}\left[e^{-R_T t} - e^{-\lambda_i t}\right] + N_0\frac{\lambda_{p\to i}R_{T\to p}}{\lambda_p - R_T}\left[\frac{e^{-R_T t} - e^{-\lambda_i t}}{\lambda_i - R_T} - \frac{e^{-\lambda_p t} - e^{-\lambda_i t}}{\lambda_i - \lambda_p}\right]. \tag{A.26}$$

The three solutions of interest are thus given by Eqs. (A.11), (A.19), and (A.26), and these are summarized in Eq. (4.2).

A.3 Solution of the differential equations for depletion only (cooling)

After the irradiation time, T_{ir}, only decay of the various produced nuclides takes place, and the corresponding differential equations become simpler. For $t > T_{ir}$ we have for the target nuclide

$$N_T(t) = N_T(T_{ir}) \tag{A.27}$$

The parent nuclides are no longer produced from a nuclear reaction via the target (i.e. the 'R' term is no longer there) which reduces Eq. (A.9) to

$$\frac{dN_p(t)}{dt} = -\lambda_p N_p(t). \tag{A.28}$$

This is easily integrated to give

$$N_p(t) = N_p(T_{ir})\,e^{-\lambda_p(t-T_{ir})}. \tag{A.29}$$

For the nuclide of interest, Eq. (A.8) reduces to

$$\frac{dN_i(t)}{dt} = \lambda_{p\to i}N_p(t) - \lambda_i N_i(t) \tag{A.30}$$

Analogous to Eqs. (A.12) - (A.19) we can solve this to give

$$N_i(t) = \frac{N_p(T_{ir})\lambda_p}{\lambda_i - \lambda_p}\left[e^{-\lambda_p(t-T_{ir})} - e^{-\lambda_i(t-T_{ir})}\right]. \tag{A.31}$$