

Memo

To
memo-to

Date
june 2017

Reference
0.1

Number of pages
1

From
memo-from

Direct line
+31 (0)88 335 8568

E-mail
arjen.markus@deltares.nl

Subject
Discontinuities and differential equations

Copy to
—

1 Introduction

Simple model for dissolution of solid matter

2 Differential equation

Basic:

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2} \quad (1)$$

with initial condition:

$$C(t = 0) = \begin{cases} 0 & \text{if } x < 0 \\ C_0 & \text{if } x \geq 0 \end{cases} \quad (2)$$

Applying the Laplace transformation (transforming C to Γ) gives:

$$\begin{cases} s\Gamma - C_0 = D \frac{d^2 \Gamma}{dx^2} & \text{if } x < 0 \\ s\Gamma = D \frac{d^2 \Gamma}{dx^2} & \text{if } x \geq 0 \end{cases} \quad (3)$$

This leads – together with the boundary and continuity conditions – to the solution:

$$\begin{cases} C = \frac{1}{2} C_0 \operatorname{erfc}\left(\frac{x}{2\sqrt{Dt}}\right) & \text{if } x < 0 \\ C = C_0 \left(1 - \frac{1}{2} \operatorname{erfc}\left(\frac{x}{2\sqrt{Dt}}\right)\right) & \text{if } x \geq 0 \end{cases} \quad (4)$$