

## 1. MODEL ANSWER

**Question.** Use Plancherel's identity to find a relation between the solution and the initial data in terms of mean square norm called  $L^2$  norm of the Cauchy problem:

$$u_t + u_{xxxx} + 2u = 0, \quad -\infty < x < \infty, \quad t > 0$$

with initial data  $u(x, 0) = u_0(x)$ ,  $-\infty < x < \infty$ ; and also find  $\lim_{t \rightarrow \infty} \int_{-\infty}^{\infty} |u(x, t)|^2 dx$ .

*Model Answer:* We have given

$$(1.1) \quad u_t + u_{xxxx} + 2u = 0, \quad -\infty < x < \infty, \quad t > 0,$$

with initial data  $u(x, 0) = u_0(x)$ ,  $-\infty < x < \infty$ . Taking Fourier transform with respect to  $x$ , from equation (1.1), we get

$$\begin{aligned} \hat{u}_t(w, t) + (iw)^4 \hat{u} + 2\hat{u} &= 0 \\ \implies \hat{u}_t(w, t) + (w^4 + 2)\hat{u} &= 0 \\ (1.2) \quad \implies \hat{u}(w, t) &= \hat{u}_0(w)e^{-t(w^4+2)} \end{aligned}$$

Using Plancherel's identity, we get

$$\begin{aligned} \int_{-\infty}^{\infty} |u(x, t)|^2 dx &= \int_{-\infty}^{\infty} |\hat{u}(w, t)|^2 dw \\ &= e^{-4t} \int_{-\infty}^{\infty} |e^{-2tw^4}| \cdot |\hat{u}_0(w)|^2 dw, \text{ using (1.2).} \\ (1.3) \quad &\leq e^{-4t} \int_{-\infty}^{\infty} |\hat{u}_0(w)|^2 dw, \text{ since } |e^{-2tw^4}| \leq 1, \forall t > 0. \end{aligned}$$

Therefore, the relation between the solution and the initial data is

$$\int_{-\infty}^{\infty} |u(x, t)|^2 dx \leq e^{-4t} \int_{-\infty}^{\infty} |\hat{u}_0(w)|^2 dw, \quad \forall t > 0.$$

Taking limit as  $t \rightarrow \infty$ , we get

$$\begin{aligned} 0 &\leq \lim_{t \rightarrow \infty} \int_{-\infty}^{\infty} |u(x, t)|^2 dx \leq \lim_{t \rightarrow \infty} e^{-4t} \int_{-\infty}^{\infty} |\hat{u}_0(w)|^2 dw = 0. \\ \implies \lim_{t \rightarrow \infty} \int_{-\infty}^{\infty} |u(x, t)|^2 dx &= 0. \end{aligned}$$

□