

Fundamental Group Schemes

Arjun Paul

Post Doctoral Fellow,
International Centre for Theoretical Sciences,
Bengaluru, India.

May 26, 2021

- 1 Fundamental groups in algebraic geometry
- 2 Nori's Fundamental group schemes
- 3 S-fundamental group schemes
- 4 Fundamental group schemes of $\mathcal{H}ilb_X^n$

- 1 Fundamental groups in algebraic geometry
- 2 Nori's Fundamental group schemes
- 3 S-fundamental group schemes
- 4 Fundamental group schemes of $\mathcal{H}ilb_X^n$

- 1 Fundamental groups in algebraic geometry
- 2 Nori's Fundamental group schemes
- 3 S-fundamental group schemes
- 4 Fundamental group schemes of \mathcal{Hilb}_X^n

- 1 Fundamental groups in algebraic geometry
- 2 Nori's Fundamental group schemes
- 3 S-fundamental group schemes
- 4 Fundamental group schemes of \mathcal{Hilb}_X^n

Fundamental groups in algebraic geometry

- Topological fundamental group of a space X , together with a marked point $x_0 \in X$, is the group $\pi_1^{\text{top}}(X, x_0)$ of homotopy classes of loops in X based at x_0 .
- The notion of homotopy classes of loops is difficult to adopt when X is an algebraic variety or a scheme.
- However, there is a different way of looking at $\pi_1^{\text{top}}(X, x_0)$, namely the group of automorphisms of the universal cover of X .
- This approach gives a better way to extend the notion of fundamental group to the case of algebraic varieties or schemes.
- In [SGA1], Alexander Grothendieck introduced **étale fundamental group** $\pi_1^{\text{ét}}(X, x_0)$ of a scheme X with a geometric point x_0 of X as a replacement of the notion of topological fundamental group.

Fundamental groups in algebraic geometry

- Topological fundamental group of a space X , together with a marked point $x_0 \in X$, is the group $\pi_1^{\text{top}}(X, x_0)$ of homotopy classes of loops in X based at x_0 .
- The notion of homotopy classes of loops is difficult to adopt when X is an algebraic variety or a scheme.
- However, there is a different way of looking at $\pi_1^{\text{top}}(X, x_0)$, namely the group of automorphisms of the universal cover of X .
- This approach gives a better way to extend the notion of fundamental group to the case of algebraic varieties or schemes.
- In [SGA1], Alexander Grothendieck introduced **étale fundamental group** $\pi_1^{\text{ét}}(X, x_0)$ of a scheme X with a geometric point x_0 of X as a replacement of the notion of topological fundamental group.

Fundamental groups in algebraic geometry

- Topological fundamental group of a space X , together with a marked point $x_0 \in X$, is the group $\pi_1^{\text{top}}(X, x_0)$ of homotopy classes of loops in X based at x_0 .
- The notion of homotopy classes of loops is difficult to adopt when X is an algebraic variety or a scheme.
- However, there is a different way of looking at $\pi_1^{\text{top}}(X, x_0)$, namely the group of automorphisms of the universal cover of X .
- This approach gives a better way to extend the notion of fundamental group to the case of algebraic varieties or schemes.
- In [SGA1], Alexander Grothendieck introduced **étale fundamental group** $\pi_1^{\text{ét}}(X, x_0)$ of a scheme X with a geometric point x_0 of X as a replacement of the notion of topological fundamental group.

Fundamental groups in algebraic geometry

- Topological fundamental group of a space X , together with a marked point $x_0 \in X$, is the group $\pi_1^{\text{top}}(X, x_0)$ of homotopy classes of loops in X based at x_0 .
- The notion of homotopy classes of loops is difficult to adopt when X is an algebraic variety or a scheme.
- However, there is a different way of looking at $\pi_1^{\text{top}}(X, x_0)$, namely the group of automorphisms of the universal cover of X .
- This approach gives a better way to extend the notion of fundamental group to the case of algebraic varieties or schemes.
- In [SGA1], Alexander Grothendieck introduced **étale fundamental group** $\pi_1^{\text{ét}}(X, x_0)$ of a scheme X with a geometric point x_0 of X as a replacement of the notion of topological fundamental group.

Fundamental groups in algebraic geometry

- Topological fundamental group of a space X , together with a marked point $x_0 \in X$, is the group $\pi_1^{\text{top}}(X, x_0)$ of homotopy classes of loops in X based at x_0 .
- The notion of homotopy classes of loops is difficult to adopt when X is an algebraic variety or a scheme.
- However, there is a different way of looking at $\pi_1^{\text{top}}(X, x_0)$, namely the group of automorphisms of the universal cover of X .
- This approach gives a better way to extend the notion of fundamental group to the case of algebraic varieties or schemes.
- In [SGA1], Alexander Grothendieck introduced **étale fundamental group** $\pi_1^{\text{ét}}(X, x_0)$ of a scheme X with a geometric point x_0 of X as a replacement of the notion of topological fundamental group.

Definition of $\pi_1^{\text{ét}}(X, x)$

- A local homomorphism of local rings $\phi : A \rightarrow B$ is said to be **unramified** if $B/\phi(m_A)B$ is a finite separable field extension of A/m_A , where m_A is the maximal ideal of A . A morphism of schemes $f : Y \rightarrow X$ is said to be **étale** if the homomorphism $\mathcal{O}_{X,f(y)} \rightarrow \mathcal{O}_{Y,y}$ is flat and unramified.
- The definition of $\pi_1^{\text{ét}}(X, x_0)$ involves a category $\text{FÉt}(X)$ of **finite étale covers** of X , together with a fiber functor $F_{x_0} : \text{FÉt}(X) \rightarrow (\text{Set})$ given by sending a finite étale cover $f : Y \rightarrow X$ to the underlined set of points of the fiber $Y_{x_0} := Y \times_X \{x_0\}$; and then $\pi_1^{\text{ét}}(X, x_0) := \text{Aut}(F_{x_0})$.
- When X is a smooth projective variety over \mathbb{C} , it turns out that

$$\pi_1^{\text{ét}}(X, x_0) \cong \widehat{\pi_1^{\text{top}}(X_{\text{an}}, x_0)},$$

the profinite completion of the topological fundamental group of the underlined complex manifold X_{an} of X .

Definition of $\pi_1^{\text{ét}}(X, x)$

- A local homomorphism of local rings $\phi : A \rightarrow B$ is said to be **unramified** if $B/\phi(m_A)B$ is a finite separable field extension of A/m_A , where m_A is the maximal ideal of A . A morphism of schemes $f : Y \rightarrow X$ is said to be **étale** if the homomorphism $\mathcal{O}_{X,f(y)} \rightarrow \mathcal{O}_{Y,y}$ is flat and unramified.
- The definition of $\pi_1^{\text{ét}}(X, x_0)$ involves a category $\text{FÉt}(X)$ of **finite étale covers of X** , together with a fiber functor $F_{x_0} : \text{FÉt}(X) \rightarrow (\text{Set})$ given by sending a finite étale cover $f : Y \rightarrow X$ to the underlined set of points of the fiber $Y_{x_0} := Y \times_X \{x_0\}$; and then $\pi_1^{\text{ét}}(X, x_0) := \text{Aut}(F_{x_0})$.
- When X is a smooth projective variety over \mathbb{C} , it turns out that

$$\pi_1^{\text{ét}}(X, x_0) \cong \widehat{\pi_1^{\text{top}}(X_{\text{an}}, x_0)},$$

the profinite completion of the topological fundamental group of the underlined complex manifold X_{an} of X .

Definition of $\pi_1^{\text{ét}}(X, x)$

- A local homomorphism of local rings $\phi : A \rightarrow B$ is said to be **unramified** if $B/\phi(m_A)B$ is a finite separable field extension of A/m_A , where m_A is the maximal ideal of A . A morphism of schemes $f : Y \rightarrow X$ is said to be **étale** if the homomorphism $\mathcal{O}_{X,f(y)} \rightarrow \mathcal{O}_{Y,y}$ is flat and unramified.
- The definition of $\pi_1^{\text{ét}}(X, x_0)$ involves a category $\text{FÉt}(X)$ of **finite étale covers of X** , together with a fiber functor $F_{x_0} : \text{FÉt}(X) \rightarrow (\text{Set})$ given by sending a finite étale cover $f : Y \rightarrow X$ to the underlined set of points of the fiber $Y_{x_0} := Y \times_X \{x_0\}$; and then $\pi_1^{\text{ét}}(X, x_0) := \text{Aut}(F_{x_0})$.
- When X is a smooth projective variety over \mathbb{C} , it turns out that

$$\pi_1^{\text{ét}}(X, x_0) \cong \widehat{\pi_1^{\text{top}}(X_{\text{an}}, x_0)},$$

the profinite completion of the topological fundamental group of the underlined complex manifold X_{an} of X .

Nori's Fundamental group schemes

- Since the definition of $\pi_1^{\text{ét}}(X, x_0)$ involves only finite étale covers of X , **it does not take care of inseparable covers of X in positive characteristics.**
- To remedy the situation, for a connected proper reduced scheme X defined over a perfect field k , in [Nor76] Madhav Nori considered more general kind of covers of X , namely “**essentially finite vector bundles**” over X .
- Let $\text{EF}(X)$ be the category of essentially finite vector bundles on X . Fix a point $x_0 \in X$, and let $T_{x_0} : \text{EF}(X) \longrightarrow \mathcal{V}ect_k$ be the functor which sends a vector bundle $E \in \text{EF}(X)$ to its fiber $E_{x_0} \in \mathcal{V}ect_k$ at x_0 .
- The quadruple $(\text{EF}(X), \mathcal{O}_X, \otimes, T_{x_0})$ forms a “neutral Tannakian category”.
- The affine k -group scheme $\pi_1^N(X, x_0)$ representing the functor $\underline{\text{Aut}}^{\otimes}(T_{x_0})$ is called **Nori's fundamental group scheme of X with base point at $x_0 \in X$.**
- We have an equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^N(X, x_0)) \cong \text{EF}(X)$.

Nori's Fundamental group schemes

- Since the definition of $\pi_1^{\text{ét}}(X, x_0)$ involves only finite étale covers of X , **it does not take care of inseparable covers of X in positive characteristics.**
- To remedy the situation, for a connected proper reduced scheme X defined over a perfect field k , in [Nor76] Madhav Nori considered more general kind of covers of X , namely “**essentially finite vector bundles**” over X .
- Let $\text{EF}(X)$ be the category of essentially finite vector bundles on X . Fix a point $x_0 \in X$, and let $T_{x_0} : \text{EF}(X) \longrightarrow \text{Vect}_k$ be the functor which sends a vector bundle $E \in \text{EF}(X)$ to its fiber $E_{x_0} \in \text{Vect}_k$ at x_0 .
- The quadruple $(\text{EF}(X), \mathcal{O}_X, \otimes, T_{x_0})$ forms a “neutral Tannakian category”.
- The affine k -group scheme $\pi_1^N(X, x_0)$ representing the functor $\underline{\text{Aut}}^{\otimes}(T_{x_0})$ is called **Nori's fundamental group scheme of X with base point at $x_0 \in X$.**
- We have an equivalence of categories $\text{Rep}_k^{\text{fd}}(\pi_1^N(X, x_0)) \cong \text{EF}(X)$.

Nori's Fundamental group schemes

- Since the definition of $\pi_1^{\text{ét}}(X, x_0)$ involves only finite étale covers of X , **it does not take care of inseparable covers of X in positive characteristics.**
- To remedy the situation, for a connected proper reduced scheme X defined over a perfect field k , in [Nor76] Madhav Nori considered more general kind of covers of X , namely “**essentially finite vector bundles**” over X .
- Let $\text{EF}(X)$ be the category of essentially finite vector bundles on X . Fix a point $x_0 \in X$, and let $T_{x_0} : \text{EF}(X) \longrightarrow \mathcal{V}ect_k$ be the functor which sends a vector bundle $E \in \text{EF}(X)$ to its fiber $E_{x_0} \in \mathcal{V}ect_k$ at x_0 .
- The quadruple $(\text{EF}(X), \mathcal{O}_X, \otimes, T_{x_0})$ forms a “neutral Tannakian category”.
- The affine k -group scheme $\pi_1^N(X, x_0)$ representing the functor $\underline{\text{Aut}}^{\otimes}(T_{x_0})$ is called **Nori's fundamental group scheme of X with base point at $x_0 \in X$.**
- We have an equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^N(X, x_0)) \cong \text{EF}(X)$.

Nori's Fundamental group schemes

- Since the definition of $\pi_1^{\text{ét}}(X, x_0)$ involves only finite étale covers of X , **it does not take care of inseparable covers of X in positive characteristics.**
- To remedy the situation, for a connected proper reduced scheme X defined over a perfect field k , in [Nor76] Madhav Nori considered more general kind of covers of X , namely “**essentially finite vector bundles**” over X .
- Let $\text{EF}(X)$ be the category of essentially finite vector bundles on X . Fix a point $x_0 \in X$, and let $T_{x_0} : \text{EF}(X) \longrightarrow \mathcal{V}ect_k$ be the functor which sends a vector bundle $E \in \text{EF}(X)$ to its fiber $E_{x_0} \in \mathcal{V}ect_k$ at x_0 .
- The quadruple $(\text{EF}(X), \mathcal{O}_X, \otimes, T_{x_0})$ forms a “neutral Tannakian category”.
- The affine k -group scheme $\pi_1^N(X, x_0)$ representing the functor $\underline{\text{Aut}}^{\otimes}(T_{x_0})$ is called **Nori's fundamental group scheme of X with base point at $x_0 \in X$.**
- We have an equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^N(X, x_0)) \cong \text{EF}(X)$.

Nori's Fundamental group schemes

- Since the definition of $\pi_1^{\text{ét}}(X, x_0)$ involves only finite étale covers of X , **it does not take care of inseparable covers of X in positive characteristics.**
- To remedy the situation, for a connected proper reduced scheme X defined over a perfect field k , in [Nor76] Madhav Nori considered more general kind of covers of X , namely “**essentially finite vector bundles**” over X .
- Let $\text{EF}(X)$ be the category of essentially finite vector bundles on X . Fix a point $x_0 \in X$, and let $T_{x_0} : \text{EF}(X) \longrightarrow \mathcal{V}ect_k$ be the functor which sends a vector bundle $E \in \text{EF}(X)$ to its fiber $E_{x_0} \in \mathcal{V}ect_k$ at x_0 .
- The quadruple $(\text{EF}(X), \mathcal{O}_X, \otimes, T_{x_0})$ forms a “neutral Tannakian category”.
- The affine k -group scheme $\pi_1^N(X, x_0)$ representing the functor $\underline{\text{Aut}}^{\otimes}(T_{x_0})$ is called **Nori's fundamental group scheme of X with base point at $x_0 \in X$.**
- We have an equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^N(X, x_0)) \cong \text{EF}(X)$.

S-Fundamental group schemes

- Let C be a smooth projective curve over a perfect field k . In [BPS06], Indranil Biswas, A. J. Parameswaran and S. Subramanian introduced a new group scheme $\pi_1^S(C, x_0)$ by considering more general kind of covers of C , namely **numerically flat vector bundles over C** . This group scheme $\pi_1^S(C, x_0)$ is called the **S-fundamental group scheme** of C with base point at $x_0 \in C$.
- The definition of $\pi_1^S(C, x_0)$ is extended for higher dimensional connected proper k -varieties X independently by V. Mehta and A. Langer [Lan11, Lan12].
- We have an exact k -linear tensor equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^S(X, x_0)) \cong \mathcal{C}_X^{\text{nf}}$, where $\mathcal{C}_X^{\text{nf}}$ is the category of **numerically flat vector bundles on X** .
- In general $\pi_1^S(X, x_0)$ carries more geometric information about X than that of $\pi_1^N(X, x_0)$ and $\pi_1^{\text{ét}}(X, x_0)$. There are faithfully flat homomorphisms

$$\pi_1^S(X, x_0) \longrightarrow \pi_1^N(X, x_0) \longrightarrow \pi_1^{\text{ét}}(X, x_0).$$

S-Fundamental group schemes

- Let C be a smooth projective curve over a perfect field k . In [BPS06], Indranil Biswas, A. J. Parameswaran and S. Subramanian introduced a new group scheme $\pi_1^S(C, x_0)$ by considering more general kind of covers of C , namely **numerically flat vector bundles over C** . This group scheme $\pi_1^S(C, x_0)$ is called the **S-fundamental group scheme** of C with base point at $x_0 \in C$.
- The definition of $\pi_1^S(C, x_0)$ is extended for higher dimensional connected proper k -varieties X independently by V. Mehta and A. Langer [Lan11, Lan12].
- We have an exact k -linear tensor equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^S(X, x_0)) \cong \mathcal{C}_X^{\text{nf}}$, where $\mathcal{C}_X^{\text{nf}}$ is the category of **numerically flat vector bundles on X** .
- In general $\pi_1^S(X, x_0)$ carries more geometric information about X than that of $\pi_1^N(X, x_0)$ and $\pi_1^{\text{ét}}(X, x_0)$. There are faithfully flat homomorphisms

$$\pi_1^S(X, x_0) \longrightarrow \pi_1^N(X, x_0) \longrightarrow \pi_1^{\text{ét}}(X, x_0).$$

S-Fundamental group schemes

- Let C be a smooth projective curve over a perfect field k . In [BPS06], Indranil Biswas, A. J. Parameswaran and S. Subramanian introduced a new group scheme $\pi_1^S(C, x_0)$ by considering more general kind of covers of C , namely **numerically flat vector bundles over C** . This group scheme $\pi_1^S(C, x_0)$ is called the **S-fundamental group scheme** of C with base point at $x_0 \in C$.
- The definition of $\pi_1^S(C, x_0)$ is extended for higher dimensional connected proper k -varieties X independently by V. Mehta and A. Langer [Lan11, Lan12].
- We have an exact k -linear tensor equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^S(X, x_0)) \cong \mathcal{C}_X^{\text{nf}}$, where $\mathcal{C}_X^{\text{nf}}$ is the category of **numerically flat vector bundles on X** .
- In general $\pi_1^S(X, x_0)$ carries more geometric information about X than that of $\pi_1^N(X, x_0)$ and $\pi_1^{\text{ét}}(X, x_0)$. There are faithfully flat homomorphisms

$$\pi_1^S(X, x_0) \longrightarrow \pi_1^N(X, x_0) \longrightarrow \pi_1^{\text{ét}}(X, x_0).$$

S-Fundamental group schemes

- Let C be a smooth projective curve over a perfect field k . In [BPS06], Indranil Biswas, A. J. Parameswaran and S. Subramanian introduced a new group scheme $\pi_1^S(C, x_0)$ by considering more general kind of covers of C , namely **numerically flat vector bundles over C** . This group scheme $\pi_1^S(C, x_0)$ is called the **S-fundamental group scheme** of C with base point at $x_0 \in C$.
- The definition of $\pi_1^S(C, x_0)$ is extended for higher dimensional connected proper k -varieties X independently by V. Mehta and A. Langer [Lan11, Lan12].
- We have an exact k -linear tensor equivalence of categories $\mathcal{R}ep_k^{\text{fd}}(\pi_1^S(X, x_0)) \cong \mathcal{C}_X^{\text{nf}}$, where $\mathcal{C}_X^{\text{nf}}$ is the category of **numerically flat vector bundles on X** .
- In general $\pi_1^S(X, x_0)$ carries more geometric information about X than that of $\pi_1^N(X, x_0)$ and $\pi_1^{\text{ét}}(X, x_0)$. There are faithfully flat homomorphisms

$$\pi_1^S(X, x_0) \longrightarrow \pi_1^N(X, x_0) \longrightarrow \pi_1^{\text{ét}}(X, x_0).$$

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n / S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n / S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n / S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n / S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n / S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

\mathcal{Hilb}_X^n , the Hilbert scheme of n points on X

- Let k be an algebraically closed field of characteristic $p \geq 0$.
- Let X be an irreducible smooth projective k -variety of dimension $d \in \{1, 2\}$.
- Fix an integer $n \geq 2$, and let \mathcal{Hilb}_X^n be the Hilbert scheme parametrizing 0-cycles of length n on X .
- \mathcal{Hilb}_X^n is a smooth projective k -variety of dimension nd , for $d = 1, 2$.
- \mathcal{Hilb}_X^n is very important and studied by many authors; see e.g., [FGA05].
- It is an interesting problem to find the S -fundamental group scheme and Nori's fundamental group schemes of \mathcal{Hilb}_X^n .
- Let $S^n(X) := X^n/S_n$ be the n -fold symmetric product of X . This is a normal projective k -variety of dimension nd .

Hilbert-Chow morphism

- Let X be an irreducible smooth projective surface over $k = \bar{k}$.
- Let $n \geq 2$. Then there is a morphism of k -schemes $\varphi : \text{Hilb}_X^n \longrightarrow S^n(X)$, known as the Hilbert-Chow morphism, which sends

$$Z \in \text{Hilb}_X^n \longmapsto \sum_{x \in \text{Supp}(Z)} \ell(\mathcal{O}_{Z,x})[x] \in S^n(X),$$

where $\text{Supp}(Z) = \{x \in X : \mathcal{O}_{Z,x} \neq 0\}$ and $\ell(\mathcal{O}_{Z,x})$ is the length of $\mathcal{O}_{Z,x}$.

- It is known that φ is a proper morphism.
- Fix a closed point $x_0 \in X$, and let $\widetilde{nx_0} \in \text{Hilb}_X^n$ be such that $\varphi(\widetilde{nx_0}) = nx_0 \in S^n(X)$, where $\varphi : \text{Hilb}_X^n \rightarrow S^n(X)$ is the Hilbert-Chow morphism.

Hilbert-Chow morphism

- Let X be an irreducible smooth projective surface over $k = \bar{k}$.
- Let $n \geq 2$. Then there is a morphism of k -schemes $\varphi : \mathcal{Hilb}_X^n \longrightarrow S^n(X)$, known as the **Hilbert-Chow morphism**, which sends

$$Z \in \mathcal{Hilb}_X^n \longmapsto \sum_{x \in \text{Supp}(Z)} \ell(\mathcal{O}_{Z,x})[x] \in S^n(X),$$

where $\text{Supp}(Z) = \{x \in X : \mathcal{O}_{Z,x} \neq 0\}$ and $\ell(\mathcal{O}_{Z,x})$ is the length of $\mathcal{O}_{Z,x}$.

- It is known that φ is a proper morphism.
- Fix a closed point $x_0 \in X$, and let $\widetilde{nx_0} \in \mathcal{Hilb}_X^n$ be such that $\varphi(\widetilde{nx_0}) = nx_0 \in S^n(X)$, where $\varphi : \mathcal{Hilb}_X^n \rightarrow S^n(X)$ is the Hilbert-Chow morphism.

Hilbert-Chow morphism

- Let X be an irreducible smooth projective surface over $k = \bar{k}$.
- Let $n \geq 2$. Then there is a morphism of k -schemes $\varphi : \mathcal{H}ilb_X^n \longrightarrow S^n(X)$, known as the **Hilbert-Chow morphism**, which sends

$$Z \in \mathcal{H}ilb_X^n \longmapsto \sum_{x \in \text{Supp}(Z)} \ell(\mathcal{O}_{Z,x})[x] \in S^n(X),$$

where $\text{Supp}(Z) = \{x \in X : \mathcal{O}_{Z,x} \neq 0\}$ and $\ell(\mathcal{O}_{Z,x})$ is the length of $\mathcal{O}_{Z,x}$.

- It is known that φ is a proper morphism.
- Fix a closed point $x_0 \in X$, and let $\widetilde{nx_0} \in \mathcal{H}ilb_X^n$ be such that $\varphi(\widetilde{nx_0}) = nx_0 \in S^n(X)$, where $\varphi : \mathcal{H}ilb_X^n \rightarrow S^n(X)$ is the Hilbert-Chow morphism.

Hilbert-Chow morphism

- Let X be an irreducible smooth projective surface over $k = \bar{k}$.
- Let $n \geq 2$. Then there is a morphism of k -schemes $\varphi : \mathcal{Hilb}_X^n \longrightarrow S^n(X)$, known as the **Hilbert-Chow morphism**, which sends

$$Z \in \mathcal{Hilb}_X^n \longmapsto \sum_{x \in \text{Supp}(Z)} \ell(\mathcal{O}_{Z,x})[x] \in S^n(X),$$

where $\text{Supp}(Z) = \{x \in X : \mathcal{O}_{Z,x} \neq 0\}$ and $\ell(\mathcal{O}_{Z,x})$ is the length of $\mathcal{O}_{Z,x}$.

- It is known that φ is a proper morphism.
- Fix a closed point $x_0 \in X$, and let $\widetilde{nx_0} \in \mathcal{Hilb}_X^n$ be such that $\varphi(\widetilde{nx_0}) = nx_0 \in S^n(X)$, where $\varphi : \mathcal{Hilb}_X^n \rightarrow S^n(X)$ is the Hilbert-Chow morphism.

Fundamental group schemes of \mathcal{Hilb}_X^n

- **Notation:** For an affine k -group scheme G , denote by G_{ab} the abelianization of G .

This is the largest abelian affine quotient k -group scheme of G .

Theorem (Paul, Sebastian [PS20])

Assume that $\text{char}(k) = p > 3$. Let X be an irreducible smooth projective surface over k . Then there is an isomorphism of affine k -group schemes

$$f^? : \pi_1^?(X, x_0)_{\text{ab}} \xrightarrow{\cong} \pi_1^?(\mathcal{Hilb}_X^n, \widetilde{nx_0}),$$

where $? = S, N, \text{ét}$. In particular, $\pi_1^?(\mathcal{Hilb}_X^n, \widetilde{nx_0})$ is abelian.

- In [PS21], we have proved similar results for the case $\dim_k(X) = 1$ and $\text{char}(k) = p > 0$.
- **Remark:** The case of étale fundamental groups was known from the work of Biswas and Hogadi in [BH15]; however, our approach is different.

Fundamental group schemes of $\mathcal{H}ilb_X^n$

- **Notation:** For an affine k -group scheme G , denote by G_{ab} the abelianization of G .

This is the largest abelian affine quotient k -group scheme of G .

Theorem (Paul, Sebastian [PS20])

Assume that $\text{char}(k) = p > 3$. Let X be an irreducible smooth projective surface over k . Then there is an isomorphism of affine k -group schemes

$$f^? : \pi_1^?(X, x_0)_{\text{ab}} \xrightarrow{\cong} \pi_1^?(\mathcal{H}ilb_X^n, \widetilde{nx_0}),$$

where $? = S, N, \text{ét}$. In particular, $\pi_1^?(\mathcal{H}ilb_X^n, \widetilde{nx_0})$ is abelian.

- In [PS21], we have proved similar results for the case $\dim_k(X) = 1$ and $\text{char}(k) = p > 0$.
- **Remark:** The case of étale fundamental groups was known from the work of Biswas and Hogadi in [BH15]; however, our approach is different.

Fundamental group schemes of \mathcal{Hilb}_X^n

- **Notation:** For an affine k -group scheme G , denote by G_{ab} the abelianization of G .

This is the largest abelian affine quotient k -group scheme of G .

Theorem (Paul, Sebastian [PS20])

Assume that $\text{char}(k) = p > 3$. Let X be an irreducible smooth projective surface over k . Then there is an isomorphism of affine k -group schemes

$$f^? : \pi_1^?(X, x_0)_{\text{ab}} \xrightarrow{\cong} \pi_1^?(\mathcal{Hilb}_X^n, \widetilde{nx_0}),$$

where $? = S, N, \text{ét}$. In particular, $\pi_1^?(\mathcal{Hilb}_X^n, \widetilde{nx_0})$ is abelian.

- In [PS21], we have proved similar results for the case $\dim_k(X) = 1$ and $\text{char}(k) = p > 0$.
- **Remark:** The case of étale fundamental groups was known from the work of Biswas and Hogadi in [BH15]; however, our approach is different.

Outline of proof

- A point of $S^n(X)$ is of the form $\bar{x} = r_1 x_1 + \cdots + r_p x_p$, with $x_j \in X$ and $r_1 \geq \cdots \geq r_p$ positive integers with $\sum_{j=1}^p r_j = n$. We call $\langle r_1, \dots, r_p \rangle$ the **type** of \bar{x} .
- Let $W \subset S^n(X)$ be the open subscheme consisting of points of types $\langle 1, 1, \dots, 1 \rangle$ and $\langle 2, 1, 1, \dots, 1 \rangle$, and let $V = \varphi^{-1}(W) \subset \mathcal{Hilb}_X^n$.
- Consider the diagram

$$\begin{array}{ccccc}
 & & V & \xrightarrow{\quad} & \mathcal{Hilb}_X^n \\
 & & \downarrow \varphi & & \downarrow \varphi \\
 X^n & \xleftarrow{j} \psi^{-1}(W) & \xrightarrow{\psi_0 := \psi|_{\psi^{-1}(W)}} & W & \xrightarrow{\quad} & S^n(X) \\
 & \searrow \psi & & & & \nearrow
 \end{array}$$

- Then the functor $\mathcal{G} : \mathcal{C}_{\mathcal{Hilb}_X^n}^{\text{nf}} \longrightarrow \mathcal{C}_{X^n}^{\text{nf}}, E \longmapsto (j_* \psi_0^* \varphi_*(E|_V))^{\vee\vee}$ defines a homomorphism $f : \pi_1^S(X^n, (x_0, \dots, x_0)) \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widehat{nx_0})$.

Outline of proof

- A point of $S^n(X)$ is of the form $\bar{x} = r_1 x_1 + \cdots + r_p x_p$, with $x_j \in X$ and $r_1 \geq \cdots \geq r_p$ positive integers with $\sum_{j=1}^p r_j = n$. We call $\langle r_1, \dots, r_p \rangle$ the **type** of \bar{x} .
- Let $W \subset S^n(X)$ be the open subscheme consisting of points of types $\langle 1, 1, \dots, 1 \rangle$ and $\langle 2, 1, 1, \dots, 1 \rangle$, and let $V = \varphi^{-1}(W) \subset \mathcal{Hilb}_X^n$.
- Consider the diagram

$$\begin{array}{ccccc}
 & & V & \hookrightarrow & \mathcal{Hilb}_X^n \\
 & & \downarrow \varphi & & \downarrow \varphi \\
 X^n & \xleftarrow{j} \psi^{-1}(W) & \xrightarrow{\psi_0 := \psi|_{\psi^{-1}(W)}} & W & \hookrightarrow S^n(X) \\
 & \searrow \psi & & & \nearrow
 \end{array}$$

- Then the functor $\mathcal{G} : \mathcal{C}_{\mathcal{Hilb}_X^n}^{\text{nf}} \longrightarrow \mathcal{C}_{X^n}^{\text{nf}}, E \longmapsto (j_* \psi_0^* \varphi_*(E|_V))^{\vee\vee}$ defines a homomorphism $f : \pi_1^S(X^n, (x_0, \dots, x_0)) \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widehat{nx_0})$.

Outline of proof

- A point of $S^n(X)$ is of the form $\bar{x} = r_1 x_1 + \cdots + r_p x_p$, with $x_j \in X$ and $r_1 \geq \cdots \geq r_p$ positive integers with $\sum_{j=1}^p r_j = n$. We call $\langle r_1, \dots, r_p \rangle$ the **type** of \bar{x} .
- Let $W \subset S^n(X)$ be the open subscheme consisting of points of types $\langle 1, 1, \dots, 1 \rangle$ and $\langle 2, 1, 1, \dots, 1 \rangle$, and let $V = \varphi^{-1}(W) \subset \mathcal{Hilb}_X^n$.
- Consider the diagram

$$\begin{array}{ccccc}
 & & V & \xrightarrow{\quad} & \mathcal{Hilb}_X^n \\
 & & \downarrow \varphi & & \downarrow \varphi \\
 X^n & \xleftarrow{j} \psi^{-1}(W) & \xrightarrow{\psi_0 := \psi|_{\psi^{-1}(W)}} & W & \xrightarrow{\quad} S^n(X) \\
 & \searrow \psi & & & \nearrow
 \end{array}$$

- Then the functor $\mathcal{G} : \mathcal{C}_{\mathcal{Hilb}_X^n}^{\text{nf}} \longrightarrow \mathcal{C}_{X^n}^{\text{nf}}, E \longmapsto (j_* \psi_0^* \varphi_*(E|_V))^{\vee\vee}$ defines a homomorphism $f : \pi_1^S(X^n, (x_0, \dots, x_0)) \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widehat{nx_0})$.

Outline of proof

- A point of $S^n(X)$ is of the form $\bar{x} = r_1 x_1 + \cdots + r_p x_p$, with $x_j \in X$ and $r_1 \geq \cdots \geq r_p$ positive integers with $\sum_{j=1}^p r_j = n$. We call $\langle r_1, \dots, r_p \rangle$ the **type** of \bar{x} .
- Let $W \subset S^n(X)$ be the open subscheme consisting of points of types $\langle 1, 1, \dots, 1 \rangle$ and $\langle 2, 1, 1, \dots, 1 \rangle$, and let $V = \varphi^{-1}(W) \subset \mathcal{Hilb}_X^n$.
- Consider the diagram

$$\begin{array}{ccccc}
 & & V & \xrightarrow{\quad} & \mathcal{Hilb}_X^n \\
 & & \downarrow \varphi & & \downarrow \varphi \\
 X^n & \xleftarrow{j} \psi^{-1}(W) & \xrightarrow{\psi_0 := \psi|_{\psi^{-1}(W)}} & W & \xrightarrow{\quad} S^n(X)
 \end{array}$$

ψ

- Then the functor $\mathcal{G} : \mathcal{C}_{\mathcal{Hilb}_X^n}^{\text{nf}} \longrightarrow \mathcal{C}_{X^n}^{\text{nf}}, \quad E \longmapsto (j_* \psi_0^* \varphi_*(E|_V))^{\vee\vee}$ defines a homomorphism $f : \pi_1^S(X^n, (x_0, \dots, x_0)) \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0})$.

Outline of proof (cont.)

- The homomorphism $\prod_{i=1}^n \pi_1^S(X, x_0) \cong \pi_1^S(X^n, (x_0, \dots, x_0)) \xrightarrow{f} \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0})$ is S_n -invariant, and gives rise to a homomorphism of affine k -group schemes

$$\widetilde{f}^S : \pi_1^S(X, x_0)_{\text{ab}} \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0}).$$

- Then we use the following result of Deligne and Milne [DM82] to show that \widetilde{f}^S is a faithfully flat closed immersion, and hence is an isomorphism of k -group schemes.
- Let $\theta : G \rightarrow G'$ be a homomorphism of affine algebraic groups over k . Then
 - ① θ is faithfully flat if and only if the functor $\widetilde{\theta} : \mathcal{R}ep_k(G') \rightarrow \mathcal{R}ep_k(G)$ is fully faithful and given any subobject $W \subset \widetilde{\theta}(V')$, with $V' \in \mathcal{R}ep_k(G')$, there is a subobject $W' \subset V'$ in $\mathcal{R}ep_k(G')$ such that $\widetilde{\theta}(W') \cong W$ in $\mathcal{R}ep_k(G)$.
 - ② θ is a closed immersion if and only if every object W of $\mathcal{R}ep_k(G)$ is isomorphic to a subquotient of an object of the form $\widetilde{\theta}(V')$, for some $V' \in \mathcal{R}ep_k(G')$.

Outline of proof (cont.)

- The homomorphism $\prod_{i=1}^n \pi_1^S(X, x_0) \cong \pi_1^S(X^n, (x_0, \dots, x_0)) \xrightarrow{f} \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0})$ is S_n -invariant, and gives rise to a homomorphism of affine k -group schemes

$$\widetilde{f}^S : \pi_1^S(X, x_0)_{\text{ab}} \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0}).$$

- Then we use the following result of Deligne and Milne [DM82] to show that \widetilde{f}^S is a faithfully flat closed immersion, and hence is an isomorphism of k -group schemes.
- Let $\theta : G \rightarrow G'$ be a homomorphism of affine algebraic groups over k . Then
 - ① θ is faithfully flat if and only if the functor $\widetilde{\theta} : \mathcal{R}ep_k(G') \rightarrow \mathcal{R}ep_k(G)$ is fully faithful and given any subobject $W \subset \widetilde{\theta}(V')$, with $V' \in \mathcal{R}ep_k(G')$, there is a subobject $W' \subset V'$ in $\mathcal{R}ep_k(G')$ such that $\widetilde{\theta}(W') \cong W$ in $\mathcal{R}ep_k(G)$.
 - ② θ is a closed immersion if and only if every object W of $\mathcal{R}ep_k(G)$ is isomorphic to a subquotient of an object of the form $\widetilde{\theta}(V')$, for some $V' \in \mathcal{R}ep_k(G')$.

Outline of proof (cont.)

- The homomorphism $\prod_{i=1}^n \pi_1^S(X, x_0) \cong \pi_1^S(X^n, (x_0, \dots, x_0)) \xrightarrow{f} \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0})$ is S_n -invariant, and gives rise to a homomorphism of affine k -group schemes

$$\widetilde{f}^S : \pi_1^S(X, x_0)_{\text{ab}} \longrightarrow \pi_1^S(\mathcal{Hilb}_X^n, \widetilde{nx_0}).$$

- Then we use the following result of Deligne and Milne [DM82] to show that \widetilde{f}^S is a faithfully flat closed immersion, and hence is an isomorphism of k -group schemes.
- Let $\theta : G \rightarrow G'$ be a homomorphism of affine algebraic groups over k . Then
 - ① θ is faithfully flat if and only if the functor $\widetilde{\theta} : \mathcal{R}ep_k(G') \rightarrow \mathcal{R}ep_k(G)$ is fully faithful and given any subobject $W \subset \widetilde{\theta}(V')$, with $V' \in \mathcal{R}ep_k(G')$, there is a subobject $W' \subset V'$ in $\mathcal{R}ep_k(G')$ such that $\widetilde{\theta}(W') \cong W$ in $\mathcal{R}ep_k(G)$.
 - ② θ is a closed immersion if and only if every object W of $\mathcal{R}ep_k(G)$ is isomorphic to a subquotient of an object of the form $\widetilde{\theta}(V')$, for some $V' \in \mathcal{R}ep_k(G')$.

References I

- [BH15] Indranil Biswas and Amit Hogadi, *On the fundamental group of a variety with quotient singularities*, Int. Math. Res. Not. IMRN (2015), no. 5, 1421–1444.
- [BPS06] Indranil Biswas, A. J. Parameswaran, and S. Subramanian, *Monodromy group for a strongly semistable principal bundle over a curve*, Duke Math. J. **132** (2006), no. 1, 1–48.
- [DM82] Pierre Deligne and James S. Milne, *Tannakian Categories*, Chapter 2, pages 101–228. In: *Hodge Cycles, Motives, and Shimura Varieties*. Lecture Notes in Mathematics, vol 900. Springer-Verlag, Berlin-New York, 1982.
- [FGA05] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Steven L. Kleiman, Nitin Nitsure, and Angelo Vistoli, *Fundamental algebraic geometry*, Mathematical Surveys and Monographs, vol. 123, American Mathematical Society, Providence, RI, 2005, Grothendieck’s FGA explained.

References II

- [Lan11] Adrian Langer, *On the S -fundamental group scheme*, Ann. Inst. Fourier (Grenoble) **61** (2011), no. 5, 2077–2119 (2012).
- [Lan12] ———, *On the S -fundamental group scheme. II*, J. Inst. Math. Jussieu **11** (2012), no. 4, 835–854.
- [Nor76] Madhav V. Nori, *On the representations of the fundamental group*, Compositio Math. **33** (1976), no. 1, 29–41.
- [PS20] Arjun Paul and Ronnie Sebastian, *Fundamental group schemes of Hilbert scheme of n points on a smooth projective surface*, Bull. Sci. Math. **164** (2020), 102898.
- [PS21] Arjun Paul and Ronnie Sebastian, *Fundamental Group Schemes of n -fold Symmetric Product of a Smooth Projective Curve*, accepted for publication in Proc. Indian Acad. Sci. Math. Sci. (2021).
- [SGA1] *Revêtements étales et groupe fondamental (SGA 1)*, Séminaire de géométrie algébrique du Bois Marie 1960–61.

Thank you!