

FUNDAMENTAL GROUPS OF MODULI OF PRINCIPAL BUNDLES ON CURVES

INDRANIL BISWAS, SWARNAVA MUKHOPADHYAY, AND ARJUN PAUL

ABSTRACT. Let X be a compact connected Riemann surface of genus g , with $g \geq 2$, and let G be a connected semisimple affine algebraic group defined over \mathbb{C} . Given any $\delta \in \pi_1(G)$, we prove that the moduli space of semistable principal G -bundles over X of topological type δ is simply connected. More generally, if G is a connected reductive complex affine algebraic group, then the fundamental group of the moduli space is isomorphic to \mathbb{Z}^{2gd} , where d is the complex dimension of the center of G . In contrast, the fundamental group of the moduli stack of principal G -bundles over X of topological type δ is shown to be isomorphic to $H^1(X, \pi_1(G))$, when G is semisimple. We also compute the fundamental group of the moduli stack of principal G -bundles when G is reductive.

CONTENTS

1. Introduction	2
2. Uniformization and fundamental group of the moduli stack	4
2.1. Uniformization	4
2.2. Fundamental Groups	5
3. Twisted moduli stack and fundamental group of its smooth locus	7
3.1. Fundamental group of the twisted moduli stack	7
3.2. Notation	10
3.3. Fundamental group of the regularly stable locus	10
3.4. Presentation as quotient stacks	11
3.5. Proof of Proposition 2.8	12
3.6. Fundamental group of $\mathcal{M}_G^{\delta, rs}$	12
3.7. Twisted Moduli spaces	13
4. Fundamental group of a moduli space of principal bundles	14
4.1. Simply connected simple groups	14
4.2. All simple groups	15
4.3. The case of reductive groups	18
Acknowledgements	20

2010 *Mathematics Subject Classification.* 14D23, 14D20, 14H30.

Key words and phrases. Moduli stack of principal bundles, uniformization, moduli space, almost commuting triples, fundamental group.

Corresponding author: Swarnava Mukhopadhyay.

1. INTRODUCTION

Let X be an irreducible smooth complex projective curve, or, equivalently, a compact connected Riemann surface. Let G be a connected reductive affine algebraic group defined over \mathbb{C} . The topological types of holomorphic principal G -bundles over X are parametrized by $\pi_1(G)$ (see [BLS, p. 186, Proposition 1.3(a)], [Ho, Section 5]). For any $\delta \in \pi_1(G)$, let M_G^δ denote the moduli space of semistable principal G -bundles over X of topological type δ . These moduli spaces have been extensively studied for the last twenty years. Our aim here is to compute the fundamental group of M_G^δ .

When $\text{genus}(X) = 0$, then M_G^δ is a point; this follows from the facts that any holomorphic principal G -bundle over \mathbb{CP}^1 admits a reduction of structure group to a maximal torus of G [Gr, p. 122, Théorème 1.1], and the holomorphic line bundles on \mathbb{CP}^1 are classified by their degree. When $\text{genus}(X) = 1$, there are explicit descriptions of M_G^δ [FMW], [FM], [La]. So we assume that $g := \text{genus}(X) > 1$.

There is a short exact sequence of groups

$$1 \longrightarrow [G, G] \longrightarrow G \xrightarrow{q} Q := G/[G, G] \cong (\mathbb{G}_m)^d \longrightarrow 1,$$

where d is the dimension of the center of G . Let

$$J_Q^\alpha(X) \cong \text{Pic}^0(X)^d$$

be the moduli space of all holomorphic principal Q -bundles on X of topological type $\alpha = q_*(\delta)$. The above homomorphism q induces a morphism of moduli spaces

$$\tilde{q} : M_G^\delta \longrightarrow J_Q^\alpha(X)$$

which is in fact an étale locally trivial fibration (see the proof of Corollary 4.5). We prove the following (see Corollary 4.5).

Theorem 1.1. *The homomorphism of fundamental groups*

$$\tilde{q}_* : \pi_1(M_G^\delta) \longrightarrow \pi_1(J_Q^\alpha(X)) \cong \mathbb{Z}^{2gd}$$

induced by the above projection \tilde{q} is an isomorphism.

Theorem 1.1 actually extends to the more general case of any connected complex affine algebraic group (see Remark 4.6).

Theorem 1.1 has the following immediate consequence:

Corollary 1.2. *For a semisimple G the moduli space M_G^δ is simply connected.*

We note that Theorem 1.1 was proved earlier in [BLR] under the assumption that $\delta = 1$. The method of [BLR] does not extend when δ is nontrivial; the crucial Lemma 2.4 in [BLR] fails to extend (also Corollary 2.2 in [BLR] does not extend).

The proof of Theorem 1.1 uses uniformization theorems [BLS, DS, KNR], for moduli stack of bundles and unirationality of M_G^δ for a semi-simple group G . For example, if we take $G = \text{SL}(r)$ and δ be an integer coprime to r , then it is well known that the corresponding moduli space is a projective, smooth, unirational Fano variety [BL, BLS,

[KNR] and hence simply connected [Ser, Kol]. However the varieties M_G^δ for general G and δ are not always smooth and hence we need use different methods to address these issues.

We first consider, the fundamental group of the moduli stack \mathcal{M}_G^δ (see Sections 3.1 for a definition) of principal G -bundles over X of topological type δ . We prove the following (see Theorem 2.10):

Theorem 1.3. *For a semisimple G the fundamental group $\pi_1(\mathcal{M}_G^\delta)$ is isomorphic to $H^1(X, \pi_1(G))$.*

It should be mentioned that more generally, when G is reductive, the fundamental group of the moduli stack of principal G -bundles over X is computed in Corollary 2.11. As an example if we take $G = \mathrm{PGL}(r)$, then for any δ , the fundamental group of the moduli stack is $(\mathbb{Z}/r\mathbb{Z})^{2g}$, where as the corresponding moduli space is simply connected.

To give a rough reason why $\pi_1(M_G^\delta)$ vanishes for G semisimple, first consider the action of the group $H^1(X, \pi_1(G))$ on any twisted moduli space (see Sections 3.1 and 3.7 for definitions) of semistable principal \tilde{G} -bundles on X , where \tilde{G} denotes the universal cover of G . This action has the property that the subgroup of $H^1(X, \pi_1(G))$ generated by all the isotropy subgroups is $H^1(X, \pi_1(G))$ itself. As a consequence of a general result of [Am], this makes the corresponding quotient by $H^1(X, \pi_1(G))$, of the twisted moduli space under consideration, a simply connected space, because the twisted moduli space is simply connected. Finally, the quotient by $H^1(X, \pi_1(G))$ of a twisted moduli space of semistable principal \tilde{G} -bundles is isomorphic to the moduli space M_G^δ , where $\delta \in \pi_1(G)$ is the element used in the construction of the twisted moduli space under consideration.

We now give an application of Theorem 1.1. If Y is a proper variety over an algebraically closed field, there is an isomorphism

$$\mathrm{Hom}(\pi_1^{\acute{e}t}(Y), \mathbb{Z}/n\mathbb{Z}) \xrightarrow{\cong} H_{\acute{e}t}^1(Y, \mathbb{Z}/n\mathbb{Z})$$

for any n . From the long exact sequence of cohomologies associated to the short exact sequence of groups

$$0 \longrightarrow \mathbb{Z}/n\mathbb{Z} \longrightarrow \mathbb{G}_m \xrightarrow{z \mapsto z^n} \mathbb{G}_m \longrightarrow 0,$$

it follows that $H_{\acute{e}t}^1(Y, \mathbb{Z}/n\mathbb{Z})$ is isomorphic to the n -torsion part

$$H_{\acute{e}t}^1(Y, \mathbb{G}_m)[n].$$

Consequently, using a generalization of Hilbert Theorem 90 ([Mi, p. 124, Proposition 4.9]), it follows that

$$\mathrm{Hom}(\pi_1^{\acute{e}t}(Y), \mathbb{Z}/n\mathbb{Z}) \simeq \mathrm{Pic}(Y)[n].$$

Now setting $Y = M_G^\delta$, where G is connected semisimple affine algebraic group over \mathbb{C} , the following corollary of Theorem 1.1 is obtained.

Corollary 1.4. *For a connected semisimple affine algebraic group G over \mathbb{C} the Picard group of M_G^δ is torsion-free.*

If G is simply connected, the Picard group of M_G is known to be \mathbb{Z} [KNR]. A result of [BLS] says that the Picard group of M_G^δ is torsion-free if G is a classical semisimple group.

2. UNIFORMIZATION AND FUNDAMENTAL GROUP OF THE MODULI STACK

Let G be a connected, reductive affine algebraic group defined over \mathbb{C} . Let X be an irreducible smooth complex projective curve. The moduli stack of principal G -bundles on X will be denoted by \mathcal{M}_G . It is well known that the stack \mathcal{M}_G is algebraic [LM].

2.1. Uniformization. Let G be a connected, semi-simple affine algebraic group defined over \mathbb{C} . We now recall the uniformization theorem that describes \mathcal{M}_G as a quotient of the affine Grassmannian [BL], [Fa], [KNR]. Let LG denote the loop group viewed as an ind-scheme over \mathbb{C} ; we note that the set of \mathbb{C} -points of LG is just $G(\mathbb{C}((t)))$. The group of positive loops (respectively, the \mathbb{C} -valued points of the groups of positive loops) will be denoted by L^+G (respectively, $G(\mathbb{C}[[t]])$). The quotient

$$\mathcal{Q}_G := LG / L^+G \quad (2.1)$$

is the affine Grassmannian. The universal cover of G will be denoted by \tilde{G} . The kernel of the projection map $\tilde{G} \rightarrow G$ is isomorphic to the fundamental group $\pi_1(G)$.

Fix a point $p \in X$. Let $L_X G$ denote the ind-sub group of LG whose set of \mathbb{C} -valued points is

$$G(\mathcal{O}_X(*p)) = G(\mathcal{O}_{X \setminus \{p\}}) \subset G(\mathbb{C}((t))).$$

The first part of the following result is standard and can be found in ([KNR], [BL], [Fa]), while the second part is proved in [DS].

Proposition 2.1. *There is a canonical isomorphism between the stacks \mathcal{M}_G and $L_X G \backslash \mathcal{Q}_G$. Moreover, the quotient map $\mathcal{Q}_G \rightarrow \mathcal{M}_G$ is locally trivial in the étale topology.*

We now recall some well known results on the objects described above; see Lemma 1.2 in [BLS, p. 185].

Proposition 2.2 ([BLS]). *Let X be an irreducible smooth complex projective curve and G a connected semisimple complex affine algebraic group. Then the following four hold:*

- (1) $\pi_0(LG) = \pi_1(G)$.
- (2) *The quotient morphism $LG \rightarrow \mathcal{Q}_G$ induces a bijection $\pi_0(LG) \rightarrow \pi_0(\mathcal{Q}_G)$. Each connected component of \mathcal{Q}_G is isomorphic to $\mathcal{Q}_{\tilde{G}}$ (defined as in (2.1) by substituting \tilde{G} in place of G). As before, \tilde{G} denotes the simply connected cover of G .*
- (3) *The group $\pi_0(L_X G)$ is canonically isomorphic to $H^1(X, \pi_1(G))$, i.e.*

$$\pi_0(L_X G) \cong H^1(X, \pi_1(G)).$$

Further via the universal coefficients theorem in cohomology, we get

$$H^1(X; \pi_1(G)) \cong \text{Hom}(H_1(X, \mathbb{Z}), \pi_1(G)) = \text{Hom}(\mathbb{Z}^{2g}, \pi_1(G)) \cong (\pi_1(G))^{2g}.$$

- (4) *The group $L_X G$ is contained in the neutral component $(LG)^0$ of LG .*

By Proposition 2.2 (cf. [BLS, p. 186, Proposition 1.3]), the set of connected components $\pi_0(\mathcal{M}_G)$ has a canonical bijection with the fundamental group $\pi_1(G)$.

Definition 2.3. For any $\delta \in \pi_1(G)$, let \mathcal{M}_G^δ denote the connected component of \mathcal{M}_G corresponding to δ . The component of $LG(\mathbb{C})$ corresponding to $\delta \in \pi_1(G)$ will be denoted by $LG^\delta(\mathbb{C})$.

Let ζ be any element in the component $\mathrm{LG}^\delta(\mathbb{C})$. By Proposition 2.2(2), we get an action of $\zeta^{-1} \mathrm{L}_X G \zeta$ on $\mathcal{Q}_{\tilde{G}}$. We now recall the uniformization theorem for each component \mathcal{M}_G^δ [BLS, Proposition 1.3(b)]. The second statement of the following proposition is derived from [DS].

Proposition 2.4. *For each $\delta \in \pi_1(G)$, let ζ be any element in the component $\mathrm{LG}^\delta(\mathbb{C})$ (see Definition 2.3). There is a canonical isomorphism of stacks*

$$\mathcal{M}_G^\delta \simeq (\zeta^{-1} \mathrm{L}_X G \zeta) \backslash \mathcal{Q}_{\tilde{G}}.$$

Moreover the quotient map $\pi : \mathcal{Q}_{\tilde{G}} \longrightarrow \mathcal{M}_G^\delta$ is locally trivial in the étale topology.

2.2. Fundamental Groups. The quotient \mathbb{C} -space \mathcal{Q}_G in (2.1) as constructed in the works of Beauville-Laszlo, Kumar and Laszlo-Sorger, [BL], [Ku], [LS], is an ind-scheme, which is a direct limit of a sequence of projective schemes. It turns out that when G is simply connected, the ind-scheme \mathcal{Q}_G is both reduced and irreducible, hence it is integral [LS, p. 508, Proposition 4.6], [BL, p. 406–407, Lemma 6.3]. The affine Grassmannian $\mathcal{Q}_G = \mathrm{LG} / \mathrm{L}^+ G$ can be realized as an inductive limit of reduced projective Schubert varieties [Ku], [Ma].

Remark 2.5. We do not need to assume that G is semisimple for defining LG . The same definition works for any reductive group G .

We now recall a lemma (Lemma 2.6) whose proof can be found in Section 8 of [PS] for $G = \mathrm{GL}_n$. The general case follows from more general results in Section 4 of [Na]. We also refer the reader to Theorem 1.6.1 and the paragraph after Theorem 1.6.1 in [Zhu] for a more comprehensive discussion.

Lemma 2.6. *The affine Grassmannian \mathcal{Q}_G is homotopic to the based loop group $\Omega_e(K_G)$, where K_G is a compact form of G .*

The following lemma is a direct consequence of Lemma 2.6.

Lemma 2.7. *Assume that G is semisimple and simply-connected. Then $\pi_1(\mathcal{Q}_G)$ is trivial.*

2.2.1. Topological Stacks. We refer the reader to papers of Behrang Noohi [No1, No2, No3] for the notion of topological stacks and its associated homotopy theory. Topological stacks are defined in Section 13.2 in [No1] and homotopy groups of topological stacks are discussed in Section 17 in [No1]. We also refer the reader to Section 5.1 in [No3] for more discussion of higher homotopy groups.

In [No1, Section 20], the author constructs a functor that takes an algebraic stack over \mathbb{C} to a topological stack (see Proposition 20.2 in [No1]). Moreover this functor has nice properties — it sends smooth morphisms to local fibrations and étale morphisms to local homeomorphisms. The stacks \mathcal{M}_G^δ are all algebraic (admitting locally finite presentation over \mathbb{C}) and hence in particular topological. Moreover, the affine Grassmannian $\mathcal{Q}_{\tilde{G}}$ has a natural topology coming from its ind-variety structure. We refer the reader to Section 3.5 for a proof of the following proposition.

Proposition 2.8. *The natural morphism $\pi : \mathcal{Q}_{\tilde{G}} \longrightarrow \mathcal{M}_G^\delta$ in Proposition 2.4 gives a morphism between the corresponding topological stacks.*

The long exact sequence in homotopy associated to a “Serre fibration” of topological stacks can be found in Section 5.2 in [No3]. We also refer the reader to Section 4.2 in [No3] for discussions on quotient stacks. Throughout this paper, we consider the fundamental group of an algebraic stack to be the fundamental group of the associated topological stack. The following lemma is due to Behrang Noohi.

Lemma 2.9. *Let \mathfrak{X} be a filtered topological stack with filtration given by $\{\mathfrak{X}_i\}_{i \in \mathbb{N}}$ and $\mathfrak{X} = \cup_{i \in \mathbb{N}} \mathfrak{X}_i$, then $\pi_1(\mathfrak{X}) = \lim \pi_1(\mathfrak{X}_i)$.*

Proof. We shall use the notion of the classifying space $f : X' \rightarrow \mathfrak{X}$ for any topological stack [No2]. This X' is a topological space, and f is a (representable) morphism with the property that the base extension f_T of f , along any morphism $T \rightarrow \mathfrak{X}$ with T a topological space T , is a weak equivalence of topological spaces. We refer the reader to [No2] for all these notions and the existence of such a topological space X' .

So we choose one classifying space, and let $\{X'_i\}_{i \in \mathbb{N}}$ be the filtration induced on X' via pull back. Since each $X'_i \rightarrow \mathfrak{X}_i$ is a weak equivalence, the result now reduces to the same statement of the lemma for topological spaces. \square

Theorem 2.10. *Assume the group G to be a semisimple affine algebraic group but not necessarily simply connected. For any $\delta \in \pi_1(G)$, there is a natural isomorphism*

$$\pi_1(\mathcal{M}_G^\delta) \cong \pi_0(L_X G) \cong H^1(X, \pi_1(G)).$$

Proof. Consider the quotient map π in Proposition 2.4. By the Proposition 2.8, this induces a map between the underlying topological stacks. Since this fibration is locally trivial with respect to the étale topology, we have a long exact sequence of homotopy groups

$$\pi_1(\mathcal{Q}_{\tilde{G}}) \rightarrow \pi_1(\mathcal{M}_G^\delta) \xrightarrow{\eta} \pi_0(\zeta^{-1} L_X G \zeta) \rightarrow \pi_0(\mathcal{Q}_{\tilde{G}}) \rightarrow 0. \quad (2.2)$$

associated to the Serre-fibration π (see Theorem 5.2 in [No3]). Now, from Lemma 2.7 it follows that the homomorphism η in (2.2) is injective, and from Proposition 2.2 we conclude that η is surjective. Consequently, the homomorphism η is an isomorphism.

Since $L_X G$ and $\zeta^{-1} L_X G \zeta$ are conjugate (by ζ), it follows that the two sets $\pi_0(\zeta^{-1} L_X G \zeta)$ and $\pi_0(L_X G)$ are bijective. Now the theorem follows from Proposition 2.2. \square

A consequence of Theorem 2.10 is the following corollary on the fundamental group of the moduli stacks of principal bundles with a reductive group as a structure group.

Corollary 2.11. *Let G be a reductive complex affine algebraic group, and let \mathcal{M}_G^δ denote a component of the moduli stack of principal G -bundles on the smooth complex projective curve X , where $\delta \in \pi_1(G)$. Then the fundamental group $\pi_1(\mathcal{M}_G^\delta)$ is a subgroup of the (abelian) group $H^1(X, \pi_1(G/Z(G))) \times H^1(X, \pi_1(G/[G, G]))$ such that the quotient group*

$$(H^1(X, \pi_1(G/Z(G))) \times H^1(X, \pi_1(G/[G, G]))) / \pi_1(\mathcal{M}_G^\delta)$$

is $H^1(X, Z([G, G]))$, where $Z([G, G])$ is the center of $[G, G]$.

Proof. Let $Z(G)$ denote the center of G , and let $[G, G]$ be the commutator subgroup of G . Consider the natural group homomorphism

$$f : G \rightarrow (G/Z(G)) \times (G/[G, G]).$$

It is easy to see that the kernel K of f is $Z(G) \cap [G, G]$ which also coincides with the center $Z([G, G])$. Now, since $[G, G]$ is semisimple, we conclude that $K = \ker(f)$ is a finite group. The corresponding morphism of moduli stacks

$$\mathcal{M}_f : \mathcal{M}_G^\delta \longrightarrow \mathcal{M}_{G/Z(G)}^{\delta_1} \times \mathcal{M}_{G/[G, G]}^{\delta_2}$$

is an étale Galois cover with Galois group $H^1(X, K)$; here \mathcal{M}_G^δ denotes a particular component of the moduli stack \mathcal{M}_G , while δ_1 and δ_2 are the images of δ in $\pi_1(G/Z(G))$ and $\pi_1(G/[G, G])$ respectively under the quotient maps. Hence from the long exact sequence of homotopy groups associated to the above fibration \mathcal{M}_f we see that $\pi_1(\mathcal{M}_G^\delta)$ injects into $\pi_1(\mathcal{M}_{G/Z(G)}^{\delta_1}) \times \pi_1(\mathcal{M}_{G/[G, G]}^{\delta_2})$ with quotient $H^1(X, Z([G, G]))$.

Since $G/Z(G)$ is semisimple, Theorem 2.10 says that

$$\pi_1(\mathcal{M}_{G/Z(G)}^{\delta_1}) = H^1(X, \pi_1(G/Z(G))).$$

On the other hand, since $G/[G, G]$ is a product of copies of the multiplicative group \mathbb{G}_m , it follows that

$$\pi_1(\mathcal{M}_{G/[G, G]}^{\delta_2}) = H^1(X, \pi_1(G/[G, G])).$$

This completes the proof. \square

The following consequence of Corollary 2.11 was observed by an anonymous referee and we thank him for his comment.

Corollary 2.12. *The rank (as an abelian group) of $\pi_1(\mathcal{M}_G^\delta)$ is $2gd$, where $d = \dim Z(G)$. In particular the fundamental groups of the moduli space \mathcal{M}_G^δ (see Theorem 1.1) and that of the moduli stack \mathcal{M}_G^δ differ only on their torsion parts.*

Proof. The result follows from the following short exact sequence obtained from Corollary 2.11, the additivity of rank in such sequences and the vanishing of the ranks of $H^1(X; \pi_1(\mathrm{Ad}(G)))$ and $H^1(X; Z([G, G]))$. We have

$$0 \longrightarrow \pi_1(\mathcal{M}_G^\delta) \longrightarrow H^1(X; \pi_1(\mathrm{Ad}(G))) \oplus H^1(X; \pi_1(\mathbb{G}_m^d)) \longrightarrow H^1(X; Z([G, G])) \longrightarrow 0.$$

Here $\mathrm{Ad}(G) = G/Z(G)$ denotes the adjoint group of G . \square

3. TWISTED MODULI STACK AND FUNDAMENTAL GROUP OF ITS SMOOTH LOCUS

In this section, we compute fundamental group of some twisted moduli stacks. We consider moduli stacks of certain reductive group $C_A \tilde{G}$ associated to a central subgroup A of \tilde{G} . The idea to consider moduli stacks for these groups $C_A \tilde{G}$ comes from the work of Beauville-Laszlo-Sorger [BLS].

3.1. Fundamental group of the twisted moduli stack. As before, let \tilde{G} be a semi-simple and simply connected affine complex algebraic group. Given a subgroup A of the center of \tilde{G} , define

$$G := \tilde{G}/A.$$

Take any $\delta \in \pi_1(G)$. We shall now recall from [BLS] the construction of a “twisted” moduli stack \mathcal{M}_G^δ dominating \mathcal{M}_G .

For any positive integer n , the group of n -th roots of unity will be denoted by μ_n . We identify μ_n with $\mathbb{Z}/n\mathbb{Z}$ using the generator $\exp(2\pi\sqrt{-1}/n)$ of μ_n . Fix an isomorphism

$$A \simeq \prod_{j=1}^s \mu_{n_j}. \quad (3.1)$$

Since $\prod_{j=1}^s \mu_{n_j}$ is canonically a subgroup of $T := (\mathbb{G}_m)^s$, the isomorphism in (3.1) identifies A with a subgroup of T . Next we identify the quotient \mathbb{G}_m/μ_n with \mathbb{G}_m via the endomorphism $z \mapsto z^n$ of \mathbb{G}_m . Using these, the quotient T/A gets identified with T . Let

$$C_A(\tilde{G}) = (\tilde{G} \times T)/A \quad (3.2)$$

be the quotient by the diagonal subgroup A . The projection to the second factor

$$C_A(\tilde{G}) \longrightarrow (\tilde{G}/A) \times (T/A) = G \times (T/A) \longrightarrow T/A = T$$

induces a morphism of the moduli stacks

$$\det : \mathcal{M}_{C_A(\tilde{G})} \longrightarrow \mathcal{M}_T. \quad (3.3)$$

Now, since \tilde{G} is simply connected, there is an isomorphism

$$\rho : \pi_1(G) \cong A.$$

Take any $\vec{d} = (d_1, \dots, d_s) \in \mathbb{Z}^s$ (see (3.1)) such that $0 \leq d_i < n_i$ for all $1 \leq i \leq s$. We set

$$\delta := \rho^{-1}(\exp(2\pi\sqrt{-1}(-d_1/n_1)), \dots, \exp(2\pi\sqrt{-1}(-d_s/n_s))). \quad (3.4)$$

Let

$$\mathcal{M}_{\tilde{G},A}^\delta := \det^{-1}((\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))) \subset \mathcal{M}_{C_A\tilde{G}} \quad (3.5)$$

be the sub-stack, where δ and $\vec{d} = (d_1, \dots, d_s)$ are related by (3.4). Following [BLS, Section 2], we shall call the stack $\mathcal{M}_{\tilde{G},A}^\delta$ the *twisted moduli stack parametrizing $C_A(\tilde{G})$ -bundles with “determinant” $(\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))$* .

It should be mentioned that the twisted principal $(C_A\tilde{G})$ -bundles, described above, can be realized as parahoric G -torsors on X [BS], [He]. So $\mathcal{M}_{\tilde{G},A}^\delta$ is also a moduli stack of parahoric G -torsors.

Remark 3.1. In [BLS], $\mathcal{M}_{\tilde{G},A}^\delta$ is defined for arbitrary semi-simple groups (not necessarily simply connected) and is denoted by $\mathcal{M}_{\tilde{G},A}^d$. The notation $\mathcal{M}_{\tilde{G},A}^\delta$ is used in [BLS] for an open and closed substack of $\mathcal{M}_{\tilde{G},A}^d$. It was observed [BLS] that for simply connected groups these two substacks coincide.

The natural projection $C_A(\tilde{G}) \longrightarrow \tilde{G}/A = G$ induces a surjective morphism of stacks

$$\mathcal{M}_{\tilde{G},A}^\delta \longrightarrow \mathcal{M}_G.$$

We now recall from [BLS, Proposition 1.3 and Example 2.4], [BS] and [He] the uniformization theorem for twisted moduli stacks.

Proposition 3.2. *Let A denote a subgroup of the center $Z(\tilde{G})$ of \tilde{G} , and consider the group $G = \tilde{G}/A$. Let ζ be any element of $\mathrm{LG}^\delta(\mathbb{C})$. Then there is a canonical isomorphism*

$$\mathcal{M}_{\tilde{G},A}^\delta \simeq (\zeta^{-1}(\mathrm{L}_X \tilde{G})\zeta) \backslash \mathcal{Q}_{\tilde{G}},$$

and moreover the natural fibration $\pi : \mathcal{Q}_{\tilde{G}} \rightarrow \mathcal{M}_{\tilde{G},A}^\delta$ is locally trivial in the étale topology.

Remark 3.3. In the statement of Proposition 3.2, note that ζ is an element of LG^δ . We explain the notation of conjugation by ζ in $\mathrm{L}_X \tilde{G}$. Consider the short exact sequence

$$0 \rightarrow T/A \rightarrow C_A(\tilde{G}) \rightarrow \tilde{G}/A \rightarrow 0,$$

where $G = \tilde{G}/A$. Moreover T/A is in the center of $C_A(\tilde{G})$. Any two lifts of ζ to $\mathrm{L} C_A(\tilde{G})$ will differ by a central element. Consequently, conjugation in $\mathrm{L} C_A(\tilde{G})$ by any lift of ζ is independent of the lift.

Proof of Proposition 3.2. We just sketch the main step to reduce to the untwisted case. First observe that \tilde{G} is the kernel of the natural homomorphism $C_A \tilde{G} \rightarrow T$. Now by construction,

$$\mathcal{M}_{\tilde{G},A}^\delta = \det^{-1}((\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))),$$

where δ and (d_1, \dots, d_s) are related by (3.4). Observe that $(\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))$ restricted to $X \setminus \{p\}$ is just $(\mathcal{O}_{X \setminus \{p\}}, \dots, \mathcal{O}_{X \setminus \{p\}})$. Thus any principal $C_A(\tilde{G})$ -bundle with determinant $(\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))$ restricted to the punctured curve $X \setminus \{p\}$, is a principal \tilde{G} -bundle on $X \setminus \{p\}$. This construction is clearly functorial, in the sense that if a scheme S parametrizes a family of $C_A(\tilde{G})$ -bundles on X with determinant $(\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))$, then the restriction of the family to $(X \setminus \{p\}) \times S$ gives a family of principal \tilde{G} -bundles on $X \setminus \{p\}$ parametrized by S .

Now the proof follows as in the untwisted case by using [DS] and the proof of Proposition 1.3 in [BLS] (see also Remark 3.6 in [BL]), but we outline the key steps for completeness. First consider the natural homomorphism $C_A(\tilde{G}) \rightarrow T$ which in turn gives a homomorphism of the corresponding loop groups $\det : \mathrm{L} C_A(\tilde{G}) \rightarrow \mathrm{L} T$. Let us consider the ind-subscheme of $\mathrm{L} C_A(\tilde{G})$ given by $\mathrm{L} \tilde{G}^\delta := \det^{-1}(z^{-d_1}, \dots, z^{-d_s})$. The discussion in the above paragraph and the uniformization theorem [DS] together give the following isomorphism of stacks:

$$\mathcal{M}_{\tilde{G},A}^\delta \simeq \mathrm{L}_X \tilde{G} \backslash \mathrm{L} \tilde{G}^\delta / \mathrm{L}^+ \tilde{G}.$$

Let ζ be any element of $\mathrm{L} G^\delta(\mathbb{C})$. Take any lift $\tilde{\zeta}$ of ζ in $\mathrm{L} \tilde{G}^\delta$. Observe that multiplication by $\tilde{\zeta}^{-1}$ gives an isomorphism of $\mathrm{L} \tilde{G}^\delta$ with $\mathrm{L} \tilde{G}$. Hence the result on uniformization follows. Local triviality follows directly from [DS]. \square

Recall that \tilde{G} is simply connected, and $G = \tilde{G}/A$, where A is a subgroup of the center of \tilde{G} isomorphic to $\pi_1(G)$. Now as in Section 2.2.1, we apply the homotopy exact sequence to the above Serre-fibration π , to get the following:

Corollary 3.4. *For any $\delta \in \pi_1(G)$, the above moduli stack $\mathcal{M}_{\tilde{G},A}^\delta$ is simply connected.*

Proof. Since \tilde{G} is simply connected, it follows from Proposition 2.2 that $\pi_0(\zeta^{-1}(L_X \tilde{G})\zeta)$ is trivial. Hence the above mentioned homotopy exact sequence gives that

$$\pi_1(\mathcal{M}_{\tilde{G},A}^\delta) \simeq \pi_0(\zeta^{-1}(L_X \tilde{G})\zeta) = \{1\}.$$

This completes the proof. \square

3.2. Notation. Let G be a connected semisimple complex affine algebraic group, and let \tilde{G} be its universal cover. For a central subgroup A of \tilde{G} isomorphic to $\pi_1(G)$, henceforth we drop the subscript A and denote by $\mathcal{M}_{\tilde{G}}^\delta$ the twisted moduli stack $\mathcal{M}_{\tilde{G},A}^\delta$.

3.3. Fundamental group of the regularly stable locus. Henceforth, we assume that $\text{genus}(X) = g \geq 2$. Take an element δ of the center of a simple and simply connected group \tilde{G} . If $g = 2$, then in this section, we assume that either $G \neq \text{SL}(2, \mathbb{C})$ or $\delta \neq 1$.

We shall recall the definition of a regularly stable principal bundle [BLS], [BH1]; for this we need the definition of a stable principal bundle which we also recall below [Ra1].

Definition 3.5. Let H be a connected reductive affine algebraic group over \mathbb{C} . A principal H -bundle E_H on X is said to be *semistable* (respectively, *stable*) if for any given reduction $E_P \subset E_H$ of the structure group of E_H to any proper parabolic subgroup $P \subsetneq H$ (not necessarily maximal), and any nontrivial dominant character $\chi : P \rightarrow \mathbb{G}_m$ which is trivial on the center of H , we have $\text{degree}(\chi_* E_P) \leq 0$ (respectively, $\text{degree}(\chi_* E_P) < 0$), where $\chi_* E_P = E_P \times^\chi \mathbb{G}_a$ is the line bundle on X associated to the principal P -bundle E_P for the character χ .

It is known that a principal H -bundle E_H is semistable (respectively, stable) if and only if for any maximal parabolic subgroup $P \subsetneq H$, and any section s of the projection $E_H/P \rightarrow X$, we have $\text{degree}(s^* T_{\text{rel}}) \geq 0$ (respectively, $\text{degree}(s^* T_{\text{rel}}) > 0$), where T_{rel} is the relative tangent bundle for the above projection $E_H/P \rightarrow X$ [Ra1, Lemma 2.1].

A principal H -bundle E on X is called *regularly stable* if

- E is stable, and
- the natural homomorphism from the center of H to $\text{Aut}(E)$, given by the action of H on E , is an isomorphism.

As before, $\mathcal{M}_{\tilde{G}}^\delta$ denotes the twisted moduli stack associated to the triple (X, G, δ) (see Section 3.2). Let

$$\mathcal{M}_{\tilde{G}}^{\delta,rs} \subset \mathcal{M}_{\tilde{G}}^\delta$$

be the open sub-stack defined by the regularly stable locus. Then there are the following natural inclusions

$$\mathcal{M}_{\tilde{G}}^{\delta,rs} \subset \mathcal{M}_{\tilde{G}}^{\delta,s} \subseteq \mathcal{M}_{\tilde{G}}^{\delta,ss} \subset \mathcal{M}_{\tilde{G}}^\delta, \quad (3.6)$$

where the $\mathcal{M}_{\tilde{G}}^{\delta,s}$ (respectively, $\mathcal{M}_{\tilde{G}}^{\delta,ss}$) denotes the open sub-stack of $\mathcal{M}_{\tilde{G}}^\delta$ given by the stable (respectively, semistable) locus.

3.4. Presentation as quotient stacks. In this section, we recall following [Ra2], a presentation of $\mathcal{M}_{\tilde{G}}^{\delta, ss}$ as a quotient stack. We closely follow and recall the constructions given in the proof of Lemma 7.3 in [BLS] (see also Proposition 3.4 in [KN]). As mentioned before, it is assumed that $g(X) \geq 2$.

Let G be any semisimple group. Choose a faithful representation $\rho : G \rightarrow \mathrm{SL}_r$. For any principal G -bundle P , let $\rho_*(P) := P_\rho = P \times^\rho \mathbb{C}^r$ be the vector bundle associated to P for the representation ρ .

Fix a closed point p on the curve X . For any integer n sufficiently large, we have $H^1(X, P_\rho(np)) = 0$ for all semistable principal G -bundles P . Indeed, this follows from semicontinuity of cohomology and boundedness of semistable principal G -bundles. Take n to be sufficiently large. Set $m(n) = r(n+1-g)$, and consider the functor parametrizing locally free quotients E of $\mathcal{O}_X^{\oplus m(n)}$ of rank r and degree rn . This is clearly representable [Ra2] by a scheme $R(n)$ along with a universal family \mathcal{E} . Moreover $R(n)$ is smooth for all n sufficiently large. By [Ra2, Sections 4.8. 4.13.3], we get a scheme $R_G(n)$ that represents the functor of global sections of the fiber bundle \mathcal{E}/G on $X \times R(n)$ which is equivalent to the functor parametrizing principal G -bundles P whose associated vector bundle $P_\rho(np)$ is a locally free quotient of $\mathcal{O}_X^{\oplus m(n)}$. By the discussion in the proof of Lemma 4.13.3 in [Ra2] we get that $R_G(n)$ is smooth for n large enough and supports an universal family of principal G -bundles. Moreover the group $\Gamma_n = \mathrm{GL}(m(n))$ acts on $R_G(n)$ and $R(n)$ and the morphism $R_G(n) \rightarrow R(n)$ is Γ_n equivariant.

Now assume as before that \tilde{G} is simply connected and $A = \prod_{j=1}^s \mu_{n_j} \subset T$ is a central subgroup of \tilde{G} such that $\tilde{G}/A = G$. The group $C_A \tilde{G} = (\tilde{G} \times T)/A$ is reductive; we first embed $C_A \tilde{G}$ into a reductive group $S = \prod_{i=1}^s \mathrm{GL}_{N_i} \times T/A$ such that the center of $C_A \tilde{G}$ goes to the center of S (see the proof of Lemma 7.3 in [BLS] for the construction of S). Now as before we have a map $\det : \mathcal{M}_S \rightarrow \mathcal{M}_{Z(S)}$, where $Z(S)$ is the center of S . For any element $d' = (d'_1, \dots, d'_{2s}) \in \mathbb{Z}^{2s}$ and a closed point p , consider the element $(\mathcal{O}_X(d'_1 p), \dots, \mathcal{O}_X(d'_{2s} p))$ of $\mathcal{M}_{Z(S)}$. We denote by $\mathcal{M}_S^{d'}$ the closed sub-stack $\det^{-1}(\mathcal{O}_X(d'_1 p), \dots, \mathcal{O}_X(d'_{2s} p))$. In particular, we have the diagram

$$\begin{array}{ccc} \mathcal{M}_{C_A \tilde{G}} & \longrightarrow & \mathcal{M}_S \\ \uparrow & & \uparrow \\ \mathcal{M}_{\tilde{G}}^\delta & \longrightarrow & \mathcal{M}_S^{d'} \end{array} \quad (3.7)$$

Here δ and d' are related by the map between the centers of $C_A \tilde{G}$ and S and equation (3.4). Since Ramanathan's construction works for arbitrary reductive group, the above construction goes through with the role of SL_r being replaced by S . Thus we get a scheme $R_{C_A(\tilde{G})}(n)$ along with a universal family of principal $C_A(\tilde{G})$ -bundles. The projection $C_A \tilde{G} \rightarrow T/A \simeq T$ induces a map

$$\det : R_{C_A \tilde{G}}(n) \rightarrow \mathcal{M}_T,$$

where $T := \mathbb{G}_m^s$. Fixing \vec{d} and δ related by equation (3.4), we define the scheme $R_{\tilde{G}}^\delta(n) = \det^{-1}(\mathcal{O}_X(d_1 p), \dots, \mathcal{O}_X(d_s p))$. Similarly we also define the scheme $R_S^{d'}$. Since $R_{C_A \tilde{G}}(n)$

is smooth for n large enough, and the morphism \det is smooth, this implies that $R_{\tilde{G}}^{\delta}(n)$ is also smooth.

For a faithful representation $\rho : C_A \tilde{G} \rightarrow S$, consider the stack $\mathcal{M}_{\tilde{G}}^{\delta}(n)$ parametrizing δ -twisted \tilde{G} -bundles E such that the corresponding vector bundles $E_{\rho}(np)$ are generated by global sections and $H^1(X, E_{\rho}(np)) = 0$. By the above discussion, we get the following:

Proposition 3.6. *The stacks $\mathcal{M}_{\tilde{G}}^{\delta}(n)$ can be presented as the quotient stack $[R_{\tilde{G}}^{\delta}(n)/\Gamma_n]$, where $R_{\tilde{G}}^{\delta}(n)$ is a smooth scheme and Γ_n is a reductive group. Moreover $R_{\tilde{G}}^{\delta}(n)$ supports a family \mathcal{W} of δ -twisted \tilde{G} -bundles along with a lift of the action of Γ_n .*

3.5. Proof of Proposition 2.8. We now observe that these spaces $R_{\tilde{G}}^{\delta}(n)$ can be used to give the morphism of the underlying topological stacks $\pi^{\text{top}} : \mathcal{Q}_{\tilde{G}} \rightarrow \mathcal{M}_{\tilde{G}}^{\delta}$ induced by the morphism of stacks $\pi : \mathcal{Q}_{\tilde{G}} \rightarrow \mathcal{M}_{\tilde{G}}^{\delta}$ in Proposition 2.4. Since $\mathcal{M}_{\tilde{G}}^{\delta}$ is a quotient of $\mathcal{M}_{\tilde{G}}^{\delta}$ by the finite group $H^1(X, \pi_1(G))$, it is enough to consider the simply connected case. For each n , we define $X_n \subset \mathcal{Q}_{\tilde{G}}$ by

$$X_n = \{\mathbf{g}\mathcal{P} \in \mathcal{Q}_{\tilde{G}} : H^1(X, \rho_*(\pi(\mathbf{g})) \otimes \mathcal{O}(np)) = 0\},$$

where $\mathcal{P} = L^+ \tilde{G}$. It follows that $X_n \subset X_{n+1}$. The affine Grassmannian $\mathcal{Q}_{\tilde{G}}$ has the structure of an ind-variety, and hence X_n acquires a natural topology. By the proof of Lemma 3.2 in [KN], we see that each X_n is open in $\mathcal{Q}_{\tilde{G}}$ and that $\bigcup_{n \geq 0} X_n = \mathcal{Q}_{\tilde{G}}$. By definition

$$X_n = \pi^{-1}(\mathcal{M}_{\tilde{G}}^{\delta}(n)),$$

and it parametrizes a family of δ -twisted \tilde{G} -bundles. By the universality of $R_{\tilde{G}}^{\delta}(n)$ (see Section 7.8 in [KNR] and Section 3 in [KN]), we get a family \mathcal{F}_n of Γ_n -bundles on X_n and Γ_n -equivariant morphism $\mathcal{F}_n \rightarrow R_{\tilde{G}}^{\delta}(n)$. Taking quotients, we get a morphism of their underlying topological quotient stacks $X_n \rightarrow \mathcal{M}_{\tilde{G}}^{\delta}(n)$. Taking the limit, we get the required morphism of topological stacks $\pi^{\text{top}} : \mathcal{Q}_{\tilde{G}} \rightarrow \mathcal{M}_{\tilde{G}}^{\delta}$. This completes the proof of Proposition 2.8.

3.6. Fundamental group of $\mathcal{M}_{\tilde{G}}^{\delta,rs}$. Let $R_{\tilde{G}}^{\delta,ss}(n)$ (respectively, $R_{\tilde{G}}^{\delta,rs}(n)$) be an open subscheme of $R_{\tilde{G}}^{\delta}(n)$ such that the associated family of δ twisted principal \tilde{G} -bundles is semistable (respectively, regularly stable). Since our representation ρ takes the center of $C_A \tilde{G}$ to the center of S , it follows from [RR, Theorem 3.18] (see also [Ra2]), that the canonical map $R_{C_A \tilde{G}}(n) \rightarrow R_S(n)$ preserves semistability.

For n large enough, $\mathcal{M}_{\tilde{G}}^{\delta,ss} \hookrightarrow \mathcal{M}_{\tilde{G}}^{\delta}(n)$ and we get that $\mathcal{M}_{\tilde{G}}^{\delta,ss}$ coincides with the quotient stack $[R_{\tilde{G}}^{\delta,ss}(n)/\Gamma_n]$.

Lemma 3.7. *For n large enough, the codimension of the complement of $R_{\tilde{G}}^{\delta,ss}(n)$ in $R_{\tilde{G}}^{\delta}(n)$ is at least two. In particular $\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,ss}) = \pi_1(\mathcal{M}_{\tilde{G}}^{\delta}(n))$.*

Proof. By [BH1, Lemma 2.1] (see also [Fa, Theorem II.6]), the codimension of $\mathcal{M}_{\tilde{G}}^{\delta,ss}$ in $\mathcal{M}_{\tilde{G}}^{\delta}(n)$ is at least two. Since $R_{\tilde{G}}^{\delta}(n)$ and $R_{\tilde{G}}^{\delta,ss}(n)$ are Γ_n torsors, this implies that the

codimension of the complement of $R_{\tilde{G}}^{\delta,ss}(n)$ in $R_{\tilde{G}}^{\delta}(n)$ is at least two. Now by construction $R_{\tilde{G}}^{\delta}(n)$ is smooth and hence $\pi_1(R_{\tilde{G}}^{\delta,ss}(n)) = \pi_1(R_{\tilde{G}}^{\delta}(n))$. Thus the result follows. \square

Lemma 3.8. *The fundamental group $\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,rs})$ is trivial if*

- *either $g(X) \geq 3$, or*
- *$g = 2$ with either $\tilde{G} \neq \mathrm{SL}_2$ or $\delta \neq 1$.*

Proof. By [BH2, Theorem 2.5], we get that the codimension of the complement of $\mathcal{M}_{\tilde{G}}^{\delta,rs}$ in $\mathcal{M}_{\tilde{G}}^{\delta,ss}$ is at least 2 if $g(X) \geq 3$ and $\tilde{G} \neq \mathrm{SL}_2$ or $\delta \neq 1$ if $g(X) = 2$. Since $R_{\tilde{G}}^{\delta,rs}(n)$ (respectively $R_{\tilde{G}}^{\delta,ss}(n)$) are both Γ_n -torsors, it implies that for n large enough the codimension of the complement of $R_{\tilde{G}}^{\delta,rs}(n)$ in $R_{\tilde{G}}^{\delta,ss}(n)$ is at least two. Moreover, both $R_{\tilde{G}}^{\delta,rs}(n)$ and $R_{\tilde{G}}^{\delta,ss}(n)$ are smooth. Thus $\pi_1(R_{\tilde{G}}^{\delta,rs}(n)) = \pi_1(R_{\tilde{G}}^{\delta,ss}(n))$. Thus for n large enough, we get

$$\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,rs}) = \pi_1(\mathcal{M}_{\tilde{G}}^{\delta,ss}) = \pi_1(\mathcal{M}_{\tilde{G}}^{\delta}(n)).$$

Now by taking limits and applying Lemma 2.9, we get that $\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,rs}) \simeq \pi_1(\mathcal{M}_{\tilde{G}}^{\delta})$. \square

3.7. Twisted Moduli spaces. The twisted moduli space $M_{\tilde{G}}^{\delta}$ associated to the triple (X, G, δ) is defined just as the twisted moduli stack is defined (see Equation (3.5)). As before, let A be the subgroup of the center of \tilde{G} isomorphic to $\pi_1(G)$.

Definition 3.9. The space $M_{\tilde{G},A}^{\delta}$ is defined to be the moduli space of semistable principal $C_A \tilde{G}$ -bundles E on X (see Equation (3.2)) such that the associated principal T -bundle obtained by extending the structure group of E using the homomorphism \det in (3.3) is the principal T -bundle corresponding to $(\mathcal{O}_X(d_1p), \dots, \mathcal{O}_X(d_sp))$, where δ and $\vec{d} = (d_1, \dots, d_s)$ are related by (3.4).

For notational simplicity, we drop the subscript A and write $M_{\tilde{G}}^{\delta}$ for $M_{\tilde{G},A}^{\delta}$.

Let

$$M_{\tilde{G}}^{\delta,rs} \subset M_{\tilde{G}}^{\delta}$$

be the twisted moduli space of regularly stable principal G -bundles associated to the triple (X, G, δ) . As before, we assume that $g \geq 3$ and for $g = 2$, either $G \neq \mathrm{SL}_2(\mathbb{C})$ or $\delta \neq 1$.

3.7.1. Presentation of moduli spaces. We continue with the same notations as in Section 3.4. By our constructions in Section 3.4, we get a map $C_A \tilde{G} \rightarrow S$ which preserves the center. This induces a morphism $\mathcal{M}_{C_A \tilde{G}} \rightarrow \mathcal{M}_S$, which, in turn, give the morphisms $\mathcal{M}_{\tilde{G}}^{\delta} \rightarrow \mathcal{M}_S^{d'}$ (see Diagram 3.7). By the discussion in Section 3.4, the stack $\mathcal{M}_{\tilde{G}}^{\delta,ss}$ (respectively, $\mathcal{M}_S^{d',ss}$) is represented as a stack quotient of $R_{\tilde{G}}^{\delta,ss}$ (respectively, $R_S^{d',ss}$) by a reductive group Γ_n . From classical theory of existence of good quotients of moduli spaces of vector bundles on a curve, it follows that $M_S^{d'}$ is a good quotient of $R_S^{d',ss}$ by Γ_n . Now since semistability is preserved ([RR, Theorem 3.18]), the construction of $M_{\tilde{G}}^{\delta}$ as a good quotient of $R_{\tilde{G}}^{\delta,ss}$ by Γ_n follows from Lemma 5.1 in [Ra2].

Corollary 3.10. *The variety $M_{\tilde{G}}^{\delta,rs}$ is simply connected.*

Proof. The coarse moduli space for $\mathcal{M}_{\tilde{G}}^{\delta,rs}$ is $M_{\tilde{G}}^{\delta,rs}$. The morphism to the coarse moduli space

$$\mathcal{M}_{\tilde{G}}^{\delta,rs} \longrightarrow M_{\tilde{G}}^{\delta,rs} \quad (3.8)$$

is a gerbe banded by the center $Z(\tilde{G})$ of \tilde{G} . A typical fiber over a point $x \in M_{\tilde{G}}^{\delta,rs}(\mathbb{C})$ is given by the classifying stack $\Gamma_x := B(Z(\tilde{G}))$, whose associated topological stack is connected. Moreover, banded gerbes are weak Serre fibrations [No3, Section 4.4]. Therefore, using the homotopy exact sequence for the morphism (3.8), we conclude that the homomorphism

$$\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,rs}) \longrightarrow \pi_1(M_{\tilde{G}}^{\delta,rs})$$

induced by the morphism (3.8) is surjective. Finally, $\pi_1(\mathcal{M}_{\tilde{G}}^{\delta,rs}) = 1$ by Lemma 3.8. \square

4. FUNDAMENTAL GROUP OF A MODULI SPACE OF PRINCIPAL BUNDLES

4.1. Simply connected simple groups. Let X be a compact connected Riemann surface of genus $g \geq 2$. Let G be a simple group with simply connected cover \tilde{G} . Consider $\pi_1(G)$ as a subgroup A of the center of \tilde{G} . As before, for any $\delta \in \pi_1(G)$, let $M_{\tilde{G}}^{\delta} = M_{\tilde{G},A}^{\delta}$ be the twisted moduli space (see Definition 3.9) of semistable bundles associated to (X, G, δ) . Recall that G is isomorphic to \tilde{G}/A .

Proposition 4.1. *The moduli space $M_{\tilde{G}}^{\delta}$ is simply connected.*

Proof. First we consider the case where $g = 2$, $G = \mathrm{SL}(2, \mathbb{C})$ and $\delta = 1$. In this case, it follows from [NR, p. 27, Lemma 6.2 (ii) and p. 33, Theorem 2] that $M_{\tilde{G}}^{\delta} = \mathbb{CP}^3$, so this moduli space is simply connected.

Therefore, we assume that either $G \neq \mathrm{SL}(2, \mathbb{C})$ or $\delta \neq 1$ whenever $g = 2$. The Zariski open subset

$$M_{\tilde{G}}^{\delta,rs} \subset M_{\tilde{G}}^{\delta} \quad (4.1)$$

is simply connected (Corollary 3.10). First observe that $M_{\tilde{G}}^{\delta}$ is a subspace of $M_{C_A \tilde{G}}^{\delta}$ realized as a fiber of the determinant map $M_{C_A \tilde{G}} \longrightarrow M_T$ in (3.3). Since $C_A \tilde{G}$ is reductive, we know by Corollary 3.4 of [BH1] that $M_{C_A \tilde{G}}^{\delta,rs}$ is the smooth locus of $M_{C_A \tilde{G}}^{\delta}$. Since $M_{C_A \tilde{G}}^{\delta,rs}$ is an étale locally trivial fiber bundle over a smooth variety with $M_{\tilde{G}}^{\delta,rs}$ as the typical fiber, it follows that $M_{\tilde{G}}^{\delta,rs}$ is the smooth locus of $M_{\tilde{G}}^{\delta}$.

We note that if Z is a normal projective variety, and $U_Z \subset Z$ is its smooth locus, then the homomorphism $\pi_1(U_Z) \longrightarrow \pi_1(Z)$ induced by the inclusion map is surjective. To prove this, take any desingularization

$$\sigma : \hat{Z} \longrightarrow Z.$$

By Zariski's main theorem (cf. [Ha, p. 280, Ch. III, Corollary 11.4]) the fibers of σ are all connected. Therefore, the homomorphism

$$\sigma_* : \pi_1(\hat{Z}) \longrightarrow \pi_1(Z)$$

induced by σ is surjective. Furthermore, the homomorphism $\pi_1(\sigma^{-1}(U_Z)) \rightarrow \pi_1(\widehat{Z})$ induced by the inclusion map is surjective, because \widehat{Z} is smooth. But

$$\sigma|_{\sigma^{-1}(U_Z)} : \sigma^{-1}(U_Z) \rightarrow U_Z$$

is an isomorphism. Hence combining the above observations we conclude that the homomorphism $\pi_1(U_Z) \rightarrow \pi_1(Z)$ is surjective.

Now, by construction ([BLS, Lemma 7.3] and [Ra2, Theorem 5.9]) the variety M_G^δ is a good quotient (see Section 3.7.1) of a smooth scheme by a reductive group; hence it is normal. So the homomorphism of fundamental groups induced by the inclusion in (4.1) is surjective. This implies that M_G^δ is simply connected, because $M_G^{\delta,rs}$ is simply connected by Corollary 3.10. \square

4.2. All simple groups. As before, assume that $g \geq 2$. Let G be any simple group. Fix an element

$$\delta \in \pi_1(G). \quad (4.2)$$

As before, M_G^δ denotes the moduli space of semistable principal G -bundles on X of topological type δ .

Theorem 4.2. *The moduli space M_G^δ is simply connected.*

Proof. Let $\gamma : \widetilde{G} \rightarrow G$ be the universal covering. The subgroup $\ker(\gamma) \subset \widetilde{G}$ will be denoted by A . This subgroup A is contained in the center of \widetilde{G} , and

$$A = \pi_1(G). \quad (4.3)$$

Let

$$\Gamma := \text{Hom}(\pi_1(X), A) = H^1(X, A) \quad (4.4)$$

be the isomorphism classes of principal A -bundles on X . We note that Γ is a finite abelian group. The group structure on A produces a group structure on Γ because A is abelian.

Let M_G^δ be the twisted moduli space of semistable principal bundles associated to (X, G, δ) , where δ is the element in (4.2). We will construct an action of Γ on M_G^δ . The homomorphism

$$\widetilde{G} \times A \rightarrow \widetilde{G}, \quad (z, a) \mapsto za$$

produces a homomorphism

$$\tau : C_A \widetilde{G} \times A \rightarrow C_A \widetilde{G},$$

where $C_A \widetilde{G}$ is the quotient group in (3.2). Given a principal $C_A \widetilde{G}$ -bundle E and a principal A -bundle F on X , we have a principal $C_A \widetilde{G}$ -bundle $\tau_*(E \times_X F)$, which is the extension of structure group of the principal $(C_A \widetilde{G} \times A)$ -bundle $E \times_X F$, using the above homomorphism τ . Clearly, $\tau_*(E \times_X F)$ is semistable if and only if E is semistable. Consequently, we get an action on M_G^δ of the group Γ in (4.4)

$$\widehat{\tau} : M_G^\delta \times \Gamma \rightarrow M_G^\delta. \quad (4.5)$$

Consider the projection to the second factor

$$C_A \widetilde{G} \rightarrow (\widetilde{G}/A) \times (T/A) = G \times (T/A) \rightarrow G,$$

where $C_A\tilde{G}$ is defined in (3.2). Given a principal $C_A\tilde{G}$ -bundle on X , we have a principal G -bundle obtained by extending the structure group using this homomorphism. This produces a morphism $M_G^\delta \longrightarrow M_{\tilde{G}}^\delta$. This morphism clearly factors through the quotient $M_{\tilde{G}}^\delta/\Gamma$ for the above action of $\hat{\tau}$ on $M_{\tilde{G}}^\delta$. The resulting morphism

$$M_{\tilde{G}}^\delta/\Gamma \longrightarrow M_G^\delta$$

is an isomorphism.

The homomorphism $\Gamma \longrightarrow \text{Aut}(M_G^\delta)$ given by the above action of $\hat{\tau}$ on M_G^δ is injective. To prove this, take any nontrivial element $h_0 \in \text{Hom}(\pi_1(X), A) = \Gamma$. Let $F \longrightarrow X$ be the principal A -bundle corresponding to h_0 . Let

$$h : \tilde{X} \longrightarrow X$$

be the étale Galois covering corresponding to $\text{kernel}(h_0) \subset \pi_1(X)$. The pullback $h^*F \longrightarrow \tilde{X}$ is a trivial principal A -bundle. Take any principal $C_A\tilde{G}$ -bundle E on X such that

- the pullback h^*E is regularly stable, and
- E lies in the moduli space $M_{\tilde{G}}^\delta$.

Since h^*F is a trivial principal A -bundle, it follows that an isomorphism between E and $\tau_*(E \times_X F)$ produces an automorphism of h^*E ; such an automorphism of h^*E is not given by an element of the center of $C_A\tilde{G}$ because F is nontrivial. Since h^*E is regularly stable, it follows that the point of $M_{\tilde{G}}^\delta$ given by E is not fixed by the action of h on $M_{\tilde{G}}^\delta$. Therefore, the above homomorphism

$$\Gamma \longrightarrow \text{Aut}(M_G^\delta)$$

is injective.

The fundamental group of the quotient of a path connected, simply connected, locally compact metric space by a faithful action of a finite group B is the quotient of B by the normal subgroup of it generated by all the isotropy subgroups [Am, p. 299, Theorem]. We shall apply this result to the action in (4.5). Note that the moduli space M_G^δ is simply connected by Proposition 4.1.

Since A is abelian, the group Γ in (4.4) is generated by the homomorphisms $\pi_1(X) \longrightarrow A$ such that the image is a cyclic subgroup of A . Take any

$$\theta : \pi_1(X) \longrightarrow A \tag{4.6}$$

such that $\theta(\pi_1(X))$ is a cyclic subgroup of A ; the order of θ will be denoted by m_0 . In view of the above mentioned result of [Am], to prove that M_G^δ is simply connected it suffices to show that the action of θ on M_G^δ has a fixed point. This result was proved in [BLS, Lemma 7.4(b)]. We give another proof of this fact and also recall the proof in [BLS].

4.2.1. First Proof. It can be shown that there is a set of generators $\{a_1, \dots, a_g, b_1, \dots, b_g\}$ of standard type of $\pi_1(X)$ with a single relation

$$\prod_{i=1}^g a_i b_i a_i^{-1} b_i^{-1} = 1$$

such that

- (1) $\theta(b_i) = 1$ for all $1 \leq i \leq g$, and
- (2) $\theta(a_i) = 1$ for all $2 \leq i \leq g$.

Indeed, for any element $\underline{\ell} := (\ell_1, \dots, \ell_{2g}) \in (\mathbb{Z}/m_0\mathbb{Z})^{2g}$ (recall that m_0 is the order of the image of θ), there is an element $A \in \mathrm{Sp}(2g, \mathbb{Z})$ such that

$$\underline{\ell}A = (\ell_1, 0, 0, \dots, 0).$$

The image of the natural homomorphism from the mapping class group for X to the automorphism group $\mathrm{Aut}(H_1(X, \mathbb{Z}))$ is the symplectic group associated to the symplectic form on $H_1(X, \mathbb{Z})$ defined by the cap product. Combining these it follows that given any standard presentation of $\pi_1(X)$, there is an element of the mapping class group that takes it to a presentation of $\pi_1(X)$ satisfying the above conditions. Clearly, the above presentation of $\pi_1(X)$ depends on θ .

Fix a maximal compact subgroup

$$\tilde{K} \subset \tilde{G}$$

Let F_{2g} denote the free group generated by $\{a_1, \dots, a_g, b_1, \dots, b_g\}$, so $\pi_1(X)$ is a quotient of F_{2g} . Let \mathcal{R} denote the space of all homomorphisms

$$\beta : F_{2g} \longrightarrow \tilde{K}$$

such that $\beta(\prod_{i=1}^g a_i b_i a_i^{-1} b_i^{-1}) = \delta'$, where $\delta' \in A$ is the element corresponding to $\delta \in \pi_1(G)$ (see (4.3), (4.2)). The group \tilde{K} acts on \mathcal{R} through the conjugation action of \tilde{K} on itself. A theorem of Ramanathan [Ral] shows that

$$M_G^\delta = \mathcal{R}/\tilde{K}.$$

The action $\hat{\tau}$ (see (4.5)) of θ (see (4.6)) on M_G^δ sends any homomorphism β as above to the homomorphism defined as follows:

- $b_i \mapsto \beta(b_i)$ for all $1 \leq i \leq g$,
- $a_i \mapsto \beta(a_i)$ for all $2 \leq i \leq g$, and
- $a_1 \mapsto \beta(a_1)\theta(a_1)$.

Define the subset of \tilde{K}^3

$$\tilde{\mathcal{S}} := \{(x_1, x_2, x_3) \in \tilde{K}^3 \mid [x_1, x_2] = \delta', [x_3, x_1] = \theta(a_1), [x_3, x_2] = 1\}.$$

The number of connected components of $\tilde{\mathcal{S}}$ coincides with that of the quotient space

$$\mathcal{S} := \tilde{\mathcal{S}}/\tilde{K}$$

because \tilde{K} is connected. The set of connected components of \mathcal{S} is described in [BFM, p. 6, Theorem 1.5.1(3)]; if we set

$$C = \begin{pmatrix} 1 & \delta' & \theta(a_1)^{-1} \\ (\delta')^{-1} & 1 & 1 \\ \theta(a_1) & 1 & 1 \end{pmatrix}$$

in [BFM, p. 6, Theorem 1.5.1], and G in [BFM, p. 6, Theorem 1.5.1] to be \tilde{K} , then the above quotient \mathcal{S} coincides with the space $\mathcal{T}_G(C)$ in [BFM, p. 6, Theorem 1.5.1]. Setting $k = 1$ in [BFM, p. 6, Theorem 1.5.1(3)] we conclude that $\mathcal{T}_G(C) = \mathcal{S}$ is nonempty because the Euler φ -function sends 1 to 1.

Take any triple $(x_1, x_2, x_3) \in \tilde{\mathcal{S}}$. Let

$$\beta_0 \in \text{Hom}(F_{2g}, \tilde{K})$$

be the homomorphism defined by

- $b_i \mapsto 1$ for all $2 \leq i \leq g$,
- $a_i \mapsto 1$ for all $2 \leq i \leq g$,
- $a_1 \mapsto x_1$, and
- $b_1 \mapsto x_2$.

Note that $\beta_0 \in \mathcal{R}$. Let

$$\beta'_0 \in \mathcal{R}/\tilde{K} = M_{\tilde{G}}^\delta$$

be the image of β_0 under the quotient map. We have $x_3 x_1 x_3^{-1} = x_1 \theta(a_1)$, because $[x_3, x_1] = \theta(a_1)$. In view of this and the third condition that $[x_3, x_2] = 1$, we conclude from the above description of the action of Γ on $\mathcal{R}/\tilde{K} = M_{\tilde{G}}^\delta$ that the above point β'_0 is fixed by the action of θ . As noted before, this completes the proof using [Am, p. 299, Theorem]. \square

4.2.2. Second Proof. The following proof is well known [BLS, Lemma 7.2(b)], but we recall it for completeness of the exposition:

Recall that $\Gamma = H^1(X, A)$ acts on $M_{\tilde{G}}^\delta$, where \tilde{G} is simply connected. Thus every element of $\gamma \in \Gamma$ gives an automorphism of $M_{\tilde{G}}^\delta$ of finite order. First following an argument in [KNR, Corollary 6.3], we show that $M_{\tilde{G}}^\delta$ is unirational. By uniformization theorem, Proposition 2.2, we get a surjection from the affine Grassmannian $\mathcal{Q}_{\tilde{G}}$ to the moduli stack $\mathcal{M}_{\tilde{G}}^\delta$. In particular there is a surjection from an open subset of $\mathcal{Q}_{\tilde{G}}$ parametrizing semistable bundle to $M_{\tilde{G}}^\delta$. Since $\mathcal{Q}_{\tilde{G}}$ is a direct limit of an increasing sequence of generalized Schubert varieties, it follows that $M_{\tilde{G}}^\delta$ is unirational.

Lemma 4.3. *Let Y be an unirational, projective variety over \mathbb{C} . Then any finite order automorphism of Y must have a fixed point.*

Proof. Let us assume that Y is smooth. Since $H^i(Y, \mathcal{O}_Y) = 0$ for all $i > 0$, by the holomorphic Lefschetz fixed-point formula, any finite order automorphism of Y must have a fixed point. Thus we are done. If Y is singular, let C be the cyclic group generated by the finite order automorphism. Let \tilde{Y} be a C -equivariant resolution of singularities, [BM], [EV], of \tilde{Y} . By the previous step, we get a fixed point of \tilde{Y} under the action of any element $c \in C$. Since the resolution is C -equivariant, we get a fixed point of Y under the action of c . This completes the proof. \square

4.3. The case of reductive groups. First assume that G is any connected semisimple affine algebraic group defined over \mathbb{C} . Take any $\delta \in \pi_1(G)$. Let M_G^δ denote the moduli space of semistable principal G -bundles on X of topological type δ .

Corollary 4.4. *The moduli space M_G^δ is simply connected.*

Proof. Let $Z \subset G$ be the center. The quotient G/Z is isomorphic to $\prod_{i=1}^d G_i$, where each G_i is simple with trivial center. The image of δ in $\pi_1(G_i)$ under the quotient map

$$G/Z = \prod_{i=1}^d G_i \longrightarrow G_i$$

will be denoted by δ_i . Let $M_{G_i}^{\delta_i}$ be the moduli space of semistable principal G_i -bundles on X of topological type δ_i .

The isomorphism classes of principal Z -bundles on X will be denoted by Γ . The homomorphism $G \times Z \longrightarrow G$, $(x, z) \longmapsto xz$, produces an action of the abelian group Γ on M_G^δ . We have

$$\prod_{i=1}^d M_{G_i}^{\delta_i} = M_G^\delta / \Gamma. \quad (4.7)$$

Now $\prod_{i=1}^d M_{G_i}^{\delta_i}$ is simply connected by Theorem 4.2. Therefore, from (4.7) we conclude that M_G^δ is simply connected. \square

Finally, let G be any connected reductive affine algebraic group defined over \mathbb{C} . The commutator subgroup $[G, G]$ is connected semisimple, and there is a short exact sequence of groups

$$1 \longrightarrow [G, G] \longrightarrow G \xrightarrow{q} Q := G/[G, G] \longrightarrow 1, \quad (4.8)$$

where the quotient Q is a product of copies of the multiplicative group \mathbb{G}_m .

Take any $\delta \in \pi_1(G)$. The image of δ in $\pi_1(Q)$ under the above projection q will be denoted by α . Let M_G^δ denote the moduli space of semistable principal G -bundles on X of topological type δ . The moduli space of principal Q -bundles on X of topological type α will be denoted by $J_Q^\alpha(X)$. We note that $J_Q^\alpha(X)$ is isomorphic to $(\text{Pic}^0(X))^d$, where d is the dimension of Q . Therefore, we have

$$\pi_1(J_Q^\alpha(X)) = \mathbb{Z}^{2gd}.$$

The projection q in (4.8) induces a morphism

$$\tilde{q} : M_G^\delta \longrightarrow J_Q^\alpha(X). \quad (4.9)$$

Corollary 4.5. *The homomorphism*

$$\tilde{q}_* : \pi_1(M_G^\delta) \longrightarrow \pi_1(J_Q^\alpha(X))$$

induced by the projection \tilde{q} in (4.9) is an isomorphism.

Proof. Let Z denote the center of $[G, G]$. The moduli space of principal Z -bundles on X will be denoted by Γ .

The projection \tilde{q} in (4.9) is surjective. It can be shown that \tilde{q} is étale locally trivial as follows. For that, let $Z_0 \subset Z$ be the connected component containing the identity element. Let

$$q' : G \longrightarrow G/Z_0$$

be the natural projection. Let

$$\alpha' := q'_*(\delta) \in \pi_1(G/Z_0)$$

be the image under the homomorphism $q'_* : \pi_1(G) \longrightarrow \pi_1(G/Z_0)$ induced by q' . Let $M_{G/Z_0}^{\alpha'}$ be the corresponding moduli space of semistable principal G/Z_0 -bundles on X . Let

$$\varpi : M_G^\delta \longrightarrow M_{G/Z_0}^{\alpha'} \times J_Q^\alpha(X)$$

be the morphism of moduli spaces corresponding to the surjective homomorphism

$$G \longrightarrow (G/Z_0) \times (G/[G, G]), \quad z \longmapsto (q'(z), q(z)).$$

It is straightforward to check that

$$\tilde{q} = p_2 \circ \varpi, \tag{4.10}$$

where $p_2 : M_{G/Z_0}^{\alpha'} \times J_Q^\alpha(X) \longrightarrow J_Q^\alpha(X)$ is the natural projection, and \tilde{q} is the map in (4.9).

Consider the finite abelian group $Z_1 := [G, G] \cap Z_0 \subset G$. Let M_{Z_1} be the group of principal Z_1 -bundles on X . The group M_{Z_1} acts on M_G^δ , and this action of M_{Z_1} on M_G^δ takes any fiber of \tilde{q} to itself. In fact, we have

$$\tilde{q}^{-1}(t)/M_{Z_1} = M_{G/Z_0}^{\alpha'}.$$

Therefore, from (4.10) it is deduced that \tilde{q} is étale locally trivial.

Hence by Corollary 4.4, we get that M is simply connected. Now from the long exact sequence of homotopy groups associated to the fiber bundle in (4.9) we conclude that the homomorphism \tilde{q}_* is an isomorphism. \square

Remark 4.6. Take \mathcal{G} to be any connected complex affine algebraic group. Let G be the quotient of \mathcal{G} by the unipotent radical of \mathcal{G} , so G is a connected complex reductive affine algebraic group. For any $\delta \in \pi_1(\mathcal{G}) = \pi_1(G)$, the natural projection

$$M_{\mathcal{G}}^\delta \longrightarrow M_G^\delta$$

is surjective with contractible fibers, in particular this map $M_{\mathcal{G}}^\delta \longrightarrow M_G^\delta$ induces an isomorphism of fundamental groups. Consequently, Theorem 1.1 computes the fundamental group of $M_{\mathcal{G}}^\delta$.

ACKNOWLEDGEMENTS

We thank the two referees for very helpful comments. We are very grateful to Behrang Noohi for Lemma 2.9. We thank S. Lawton and D. Ramras for making us aware of the question addressed here. We thank the Institute for Mathematical Sciences in the National University of Singapore for hospitality while this work was being completed. The first author is supported by a J. C. Bose Fellowship. The second author was supported in part by a Simons Travel Grant and by NSF grant DMS-1361159 (PI: Patrick Brosnan) and also by the Science and Engineering Research Board, India (SRG/2019/000513). The first and second author were also supported by the Department of Atomic Energy, India, under project no. 12-R&D-TFR-5.01-0500.

REFERENCES

- [Am] M. A. Armstrong, The fundamental group of the orbit space of a discontinuous group, *Proc. Camb. Phil. Soc.* **64** (1968), 299–301. doi: [10.1017/s0305004100042845](https://doi.org/10.1017/s0305004100042845). [[3](#), [16](#), and [18](#).]
- [BL] A. Beauville and Y. Laszlo, Conformal blocks and generalized theta functions, *Comm. Math. Phys.* **164** (1994), 385–419. url: <https://projecteuclid.org/euclid.cmp/1104270837>. [[3](#), [4](#), [5](#), and [9](#).]
- [BLS] A. Beauville, Y. Laszlo and C. Sorger, The Picard group of the moduli of G -bundles on a curve, *Compositio Math.* **112** (1998), 183–216. doi: [10.1023/A:1000477122220](https://doi.org/10.1023/A:1000477122220). [[2](#), [3](#), [4](#), [5](#), [7](#), [8](#), [9](#), [10](#), [11](#), [15](#), [16](#), and [18](#).]
- [BM] E. Bierstone and P. D. Milman, Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant, *Invent. Math.* **128** (1997), 207–302. doi: [10.1007/s002220050141](https://doi.org/10.1007/s002220050141). [[18](#).]
- [BH1] I. Biswas and N. Hoffmann, A Torelli theorem for moduli spaces of principal bundles over a curve, *Ann. Inst. Four.* **62** (2012), 87–106. doi: [10.5802/aif.2700](https://doi.org/10.5802/aif.2700). [[10](#), [12](#), and [14](#).]
- [BH2] I. Biswas and N. Hoffmann, Poincaré families of G -bundles on a curve, *Math. Ann.* **352** (2012), 133–154. doi: [10.1007/s00208-010-0628-x](https://doi.org/10.1007/s00208-010-0628-x). [[13](#).]
- [BS] V. Balaji and C. S. Seshadri, Moduli of parahoric \mathcal{G} -torsors on a compact Riemann surface, *J. Algebraic Geom.* **24** (2015), 1–49. doi: [10.1090/S1056-3911-2014-00626-3](https://doi.org/10.1090/S1056-3911-2014-00626-3). [[8](#).]
- [BLR] I. Biswas, S. Lawton and D. Ramras, Fundamental groups of character varieties: surfaces and tori, *Math. Zeit.* **281** (2015), 415–425. doi: [10.1007/s00209-015-1492-x](https://doi.org/10.1007/s00209-015-1492-x). [[2](#).]
- [BFM] A. Borel, R. Friedman and J. M. Morgan, Almost commuting elements in compact Lie groups, *Mem. Amer. Math. Soc.* **157** (2002), no. 747. doi: [10.1090/memo/0747](https://doi.org/10.1090/memo/0747). [[17](#).]
- [DS] V. G. Drinfeld and C. T. Simpson, B-structures on G -bundles and local triviality, *Math. Res. Lett.* **2** (1995), 823–829. doi: [10.4310/MRL.1995.v2.n6.a13](https://doi.org/10.4310/MRL.1995.v2.n6.a13). [[2](#), [4](#), [5](#), and [9](#).]
- [EV] S. Encinas and O. Villamayor, Good points and constructive resolution of singularities, *Acta Math.* **181** (1998), 109–158. doi: [10.1007/BF02392749](https://doi.org/10.1007/BF02392749). [[18](#).]
- [Fa] G. Faltings, A proof of the Verlinde formula, *Jour. Alg. Geom.* **3** (1994), 347–374. [[4](#) and [12](#).]
- [FM] R. Friedman and J. W. Morgan, Holomorphic principal bundles over elliptic curves. II. The parabolic construction, *Jour. Diff. Geom.* **56** (2000), 301–379. doi: [10.4310/jdg/1090347646](https://doi.org/10.4310/jdg/1090347646). [[2](#).]
- [FMW] R. Friedman, J. W. Morgan and E. Witten, Principal G -bundles over elliptic curves, *Math. Res. Lett.* **5** (1998), 97–118. doi: [10.4310/MRL.1998.v5.n1.a8](https://doi.org/10.4310/MRL.1998.v5.n1.a8). [[2](#).]
- [Gr] A. Grothendieck, Sur la classification des fibrés holomorphes sur la sphère de Riemann, *Amer. Jour. Math.* **79** (1957), 121–138. doi: [10.2307/2372388](https://doi.org/10.2307/2372388). [[2](#).]
- [Ha] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977. doi: [10.1007/978-1-4757-3849-0](https://doi.org/10.1007/978-1-4757-3849-0). [[14](#).]
- [He] J. Heinloth, Uniformization of \mathcal{G} -bundles, *Math. Ann.* **347** (2010), 499–528. doi: [10.1007/s00208-009-0443-4](https://doi.org/10.1007/s00208-009-0443-4). [[8](#).]
- [Ho] N. Hoffmann, On Moduli Stacks of G -Bundles over a Curve, *Affine flag manifolds and principal bundles*, 155–163, Trends in Mathematics, Birkhäuser/Springer Basel AG, Basel, 2010. doi: [10.1007/978-3-0346-0288-4_5](https://doi.org/10.1007/978-3-0346-0288-4_5). [[2](#).]
- [KN] S. Kumar and M. S. Narasimhan, Picard group of the moduli spaces of G -bundles, *Math. Ann.* **308** (1997), 155–173. doi: [10.1007/s002080050070](https://doi.org/10.1007/s002080050070). [[11](#) and [12](#).]
- [KNR] S. Kumar, M. S. Narasimhan and A. Ramanathan, Infinite Grassmannians and moduli spaces of G -bundles, *Math. Ann.* **300** (1994), 41–75. doi: [10.1007/BF01450475](https://doi.org/10.1007/BF01450475). [[2](#), [3](#), [4](#), [12](#), and [18](#).]
- [Kol] J. Kollár, Fundamental groups of rationally connected varieties, *Michigan Math. J.* **48** (2000), 359–368. doi: [10.1307/mmj/1030132724](https://doi.org/10.1307/mmj/1030132724). [[3](#).]
- [Ku] S. Kumar, Demazure character formula in arbitrary Kac-Moody setting, *Invent. Math.* **89** (1987), 395–423. doi: [10.1007/BF01389086](https://doi.org/10.1007/BF01389086). [[5](#).]
- [La] Y. Laszlo, About G -bundles over elliptic curves, *Ann. Inst. Fourier* **48** (1998), 413–424. doi: [10.5802/aif.1623](https://doi.org/10.5802/aif.1623). [[2](#).]
- [LM] G. Laumon and L. Moret-Bailly, *Champs algebriques*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3, Folge, A Series of Modern Surveys in Mathematics, 39 Springer-Verlag, Berlin, 2000. doi: [10.1007/978-3-540-24899-6](https://doi.org/10.1007/978-3-540-24899-6). [[4](#).]

- [LS] Y. Laszlo and C. Sorger, The line bundles on the moduli of parabolic G -bundles over curves and their sections, *Ann. Sci. École Norm. Sup.* **30** (1997), 499–525. doi: [10.1016/S0012-9593\(97\)89929-6](https://doi.org/10.1016/S0012-9593(97)89929-6). [↑ 5.]
- [Ma] O. Mathieu, Formules de caractères pour les algèbres de Kac–Moody générales, *Astérisque*, No. 159–160 (1988). [↑ 5.]
- [Mi] J. S. Milne, *Etale Cohomology*, Princeton Mathematical Series, No 33. Princeton University Press, Princeton, N.J., 1980. [↑ 3.]
- [Na] D. Nadler, Matsuki correspondence for the affine Grassmannian, *Duke Math. J.* **124** (2004), 421–457. doi: [10.1215/S0012-7094-04-12431-5](https://doi.org/10.1215/S0012-7094-04-12431-5). [↑ 5.]
- [NR] M. S. Narasimhan and S. Ramanan, Moduli of vector bundles on a compact Riemann surface, *Ann. Math.* **89** (1969), 14–51. doi: [10.2307/1970807](https://doi.org/10.2307/1970807). [↑ 14.]
- [No1] B. Noohi, Foundations of Topological Stacks I, arXiv:[math/0503247v1](https://arxiv.org/abs/math/0503247v1). [↑ 5.]
- [No2] B. Noohi, Homotopy types of topological stacks, *Adv. Math.* **230** (2012), 2014–2047. doi: [10.1016/j.aim.2012.04.001](https://doi.org/10.1016/j.aim.2012.04.001). [↑ 5 and 6.]
- [No3] B. Noohi, Fibrations of topological stacks, *Adv. Math.* **252** (2014), 612–640. doi: [10.1016/j.aim.2013.11.008](https://doi.org/10.1016/j.aim.2013.11.008). [↑ 5, 6, and 14.]
- [PS] A. Pressley and G. Segal, Loop groups, *Oxford Mathematical Monographs*, The Clarendon Press, Oxford University Press, New York, 1986. [↑ 5.]
- [RR] A. Ramathan and S. Ramanan, Some remarks on the instability flag, *The Tohoku Math. Jour.* **36** (1984), 269–291. doi: [10.2748/tmj/1178228852](https://doi.org/10.2748/tmj/1178228852). [↑ 12 and 13.]
- [Ra1] A. Ramanathan, Stable principal bundles on a compact Riemann surface, *Math. Ann.* **213** (1975), 129–152. doi: [10.1007/BF01343949](https://doi.org/10.1007/BF01343949). [↑ 10 and 17.]
- [Ra2] A. Ramanathan, Moduli for principal bundles over algebraic curves. II, *Proc. Indian Acad. Sci. Math. Sci.* **106** (1996), 421–449. doi: [10.1007/BF02837697](https://doi.org/10.1007/BF02837697). [↑ 11, 12, 13, and 15.]
- [Ser] J.-P. Serre, On the fundamental group of a unirational variety, *J. London Math. Soc.* **34** (1959), 481–484. doi: [10.1112/jlms/s1-34.4.481](https://doi.org/10.1112/jlms/s1-34.4.481). [↑ 3.]
- [Zhu] X. Zhu, An introduction to affine Grassmannians and the geometric Satake equivalence, *Geometry of moduli spaces and representation theory*, 59–154, IAS/Park City Math. Ser., 24, Amer. Math. Soc., Providence, RI, 2017. doi: [10.1090/pcms/024/02](https://doi.org/10.1090/pcms/024/02). [↑ 5.]

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

Email address: indranil@math.tifr.res.in

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

Email address: swarnavaster@gmail.com, swarnava@math.tifr.res.in

DEPARTMENT OF MATHEMATICS, INTERNATIONAL CENTRE FOR THEORETICAL SCIENCES, SURVEY NO. 151, SHIVAKOTE, HESARAGHATTA HOBLI, BENGALURU - 560 089, INDIA

Email address: arjun.math.tifr@gmail.com