

---

# Algebra I: Group Theory

---

Dr. Arjun Paul

Assistant Professor  
Department of Mathematics and Statistics  
Indian Institute of Science Education and Research Kolkata,  
Mohanpur - 741 246, Nadia,  
West Bengal, India.  
Email: [arjun.paul@iiserkol.ac.in](mailto:arjun.paul@iiserkol.ac.in).  
Homepage: <https://sites.google.com/site/arjunpaultifr/>

Version: November 4, 2023 at 11:51pm (IST).  
Available at: <https://arjunpaul29.github.io/home/notes/Algebra-I.pdf>

*Note: This note will be updated from time to time.  
If you find any potential mistakes/typos, please bring it to **my notice**.*



*To my students ...*



# Contents

<b>List of Symbols</b>	<b>vii</b>
<b>1 Foundation of Arithmetic</b>	<b>1</b>
1.1 What is a Natural Number? . . . . .	1
1.2 Integers: Construction & Basic Operations . . . . .	8
1.3 Division Algorithm . . . . .	10
<b>2 Group Theory</b>	<b>13</b>
2.1 Group . . . . .	13
2.2 Subgroup . . . . .	22
2.3 Cyclic group . . . . .	25
2.4 Product of subgroups . . . . .	28
2.4.1 Lattice diagram . . . . .	30
2.5 Permutation Groups . . . . .	31
2.6 Group homomorphism . . . . .	39
2.7 Notion of Quotient & Cosets . . . . .	46
2.8 Normal Subgroup & Quotient Group . . . . .	50
2.9 Isomorphism Theorems . . . . .	53
2.9.1 Inner Automorphisms . . . . .	55
2.10 Direct Product & Direct Sum of Groups . . . . .	59
2.11 Group Action . . . . .	65
2.12 Conjugacy Action & Class Equations . . . . .	72
2.12.1 $p$ -groups . . . . .	75
2.13 Simple Groups . . . . .	77
2.13.1 Simplicity of $A_n$ , for $n \geq 5$ . . . . .	78
2.13.2 *Simplicity of $\text{PSL}_n(F)$ , for $n \geq 3$ . . . . .	79
2.14 Sylow's Theorems . . . . .	84
2.15 Miscellaneous Exercises . . . . .	89
2.16 Applications of Sylow's Theorems . . . . .	93
2.16.1 Quaternion group $Q_8$ . . . . .	96
2.17 Finitely Generated Abelian Groups . . . . .	98

2.18 Free Group . . . . .	99
2.19 Solvable & Nilpotent Groups . . . . .	99
2.20 Semi-direct product . . . . .	99
2.21 Linear Groups . . . . .	100

# List of Symbols

$\emptyset$	Empty set
$\mathbb{Z}$	The set of all integers
$\mathbb{Z}_{\geq 0}$	The set of all non-negative integers
$\mathbb{N}$	The set of all natural numbers (i.e., positive integers)
$\mathbb{Q}$	The set of all rational numbers
$\mathbb{R}$	The set of all real numbers
$\mathbb{C}$	The set of all complex numbers
$<$	Less than
$\leq$	Less than or equal to
$>$	Greater than
$\geq$	Greater than or equal to
$\subset$	Proper subset
$\subseteq$	Subset or equal to
$\subsetneq$	Subset but not equal to (c.f. proper subset)
$\exists$	There exists
$\nexists$	Does not exist
$\forall$	For all
$\in$	Belongs to
$\notin$	Does not belong to
$\sum$	Sum
$\prod$	Product
$\pm$	Plus and minus
$\infty$	Infinity
$\sqrt{a}$	Square root of $a$
$\cup$	Union
$\sqcup$	Disjoint union
$\cap$	Intersection
$A \rightarrow B$	$A$ mapping into $B$
$a \mapsto b$	$a$ maps to $b$
$\hookrightarrow$	Inclusion map
$A \setminus B$	$A$ setminus $B$
$\cong$	Isomorphic to
$A := \dots$	$A$ is defined to be ...
$\square$	End of a proof

Symbol	Name	Symbol	Name
$\alpha$	alpha	$\beta$	beta
$\gamma$	gamma	$\delta$	delta
$\pi$	pi	$\phi$	phi
$\varphi$	var-phi	$\psi$	psi
$\epsilon$	epsilon	$\varepsilon$	var-epsilon
$\zeta$	zeta	$\eta$	eta
$\theta$	theta	$\iota$	iota
$\kappa$	kappa	$\lambda$	lambda
$\mu$	mu	$\nu$	nu
$\upsilon$	upsilon	$\rho$	rho
$\varrho$	var-rho	$\xi$	xi
$\sigma$	sigma	$\tau$	tau
$\chi$	chi	$\omega$	omega
$\Omega$	Capital omega	$\Gamma$	Capital gamma
$\Theta$	Capital theta	$\Delta$	Capital delta
$\Lambda$	Capital lambda	$\Xi$	Capital xi
$\Sigma$	Capital sigma	$\Pi$	Capital pi
$\Phi$	Capital phi	$\Psi$	Capital psi

Some of the useful Greek alphabets



## Chapter 1

# Foundation of Arithmetic

### 1.1 What is a Natural Number?

We begin with axiomatic definition of the set of all *natural numbers*, known as *Peano's axioms*, also known as *Dedekind–Peano axioms*. This was originally proposed by Richard Dedekind in 1888, and was published in a simplified version as a collection of axioms in 1989 by Giuseppe Peano in his book *Arithmetices principia, nova methodo exposita* (in English: *The principles of arithmetic presented by a new method*). We define addition and multiplication of natural numbers, and briefly discuss their useful arithmetic properties (with outline of proofs) that we are familiar with from elementary mathematics courses, without possibly thinking *why and how these work*? The purpose of this section is to provide a *logical foundation of natural numbers and their arithmetic*.

**Axiom 1.1.1** (Peano's axioms). *There is a set  $\mathbb{N}$  satisfying the following axioms.*

(P1)  $1 \in \mathbb{N}$  (so  $\mathbb{N} \neq \emptyset$ ); the element 1 is called one.

(P2) Axiom of equality: There is a relation “=” on  $\mathbb{N}$ , called the equality, satisfying the following properties.

- (i)  $a = a, \forall a \in \mathbb{N}$ ,
- (ii) given  $a, b \in \mathbb{N}$ , we have  $a = b \Rightarrow b = a$ , and
- (iii) given  $a, b, c \in \mathbb{N}$ , if  $a = b$  and  $b = c$ , then  $a = c$ .

In other words, the relation “=” on  $\mathbb{N}$  is an equivalence relation on  $\mathbb{N}$ . If “ $a = b$ ”, we say that “ $a$  is equal to  $b$ ”. If “ $a = b$ ” is not true, we say that “ $a$  is not equal to  $b$ ”, expressed symbolically as “ $a \neq b$ ”.

**(Remark:** The axiom (P2) was included in the original list of axioms published by Peano in 1889. However, since the axiom (P2) is logically valid in first-order logic with equality, this is always accepted, and is not considered to be a part of Axiom 1.1.1 in modern treatments.)

(P3) For each  $n \in \mathbb{N}$ , there is a unique  $s(n) \in \mathbb{N}$ , called the successor of  $n$ .

(P4) 1 is not a successor of any element of  $\mathbb{N}$ .

(P5) Given  $m, n \in \mathbb{N}$  with  $m \neq n$ , we have  $s(m) \neq s(n)$ .

(P6) Principle of Mathematical Induction: If a subset  $S \subseteq \mathbb{N}$  has properties that

- (i)  $1 \in S$ , and
- (ii)  $n \in S \Rightarrow s(n) \in S$ ,

then  $S = \mathbb{N}$ .

The elements of  $\mathbb{N}$  are called *natural numbers*, and hence  $\mathbb{N}$  is called the *set of all natural numbers*.

**Exercise 1.1.2.** Verify that  $s : \mathbb{N} \rightarrow \mathbb{N}$ ,  $n \mapsto s(n)$  is injective but not surjective.

**Remark 1.1.3.** In contrast to our naive intuition, the properties (P1)–(P5) in Peano's Axioms 1.1.1 do not guarantee that the successor function generates all natural numbers (we are familiar with) except for 1. To make our naive intuition works, we need the assumption (P6), known as the Principle of Mathematical Induction.

**Lemma 1.1.4.** If  $n \in \mathbb{N}$  with  $n \neq 1$ , then there is a unique element  $p(n) \in \mathbb{N}$ , called the *predecessor* of  $n$ , such that  $s(p(n)) = n$ .

*Proof.* Since  $s : \mathbb{N} \rightarrow \mathbb{N}$ ,  $n \mapsto s(n)$  is injective by (P5), uniqueness of  $p(n)$  follows. To show existence of  $p(n)$ , for each  $n \in \mathbb{N} \setminus \{1\}$ , it is enough to show that

$$s(\mathbb{N}) := \{s(n) : n \in \mathbb{N}\} = \mathbb{N} \setminus \{1\}.$$

Since  $1 \notin s(\mathbb{N}) := \{s(n) : n \in \mathbb{N}\}$  by (P4), to show that  $s(\mathbb{N}) = \mathbb{N} \setminus \{1\}$ , it is enough to show that

$$T := s(\mathbb{N}) \cup \{1\} = \mathbb{N}.$$

Clearly  $T \subseteq \mathbb{N}$  and  $1 \in T$ . If  $m \in T$ , then  $m = 1$  or  $m = s(n)$ , for some  $n \in \mathbb{N}$ , and so in both cases,  $s(m) \in T$  by construction of  $T$ . Then (P6) tells us that  $T = \mathbb{N}$ . This completes the proof.  $\square$

**Definition 1.1.5.** A *binary operation* on a set  $S$  is a map  $S \times S \rightarrow S$ .

**Definition 1.1.6.** On the set  $\mathbb{N}$ , we define two binary operations

$$(1.1.7) \quad \text{Addition} \quad + : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}, \quad (m, n) \mapsto m + n,$$

$$(1.1.8) \quad \text{and Multiplication} \quad \cdot : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}, \quad (m, n) \mapsto m \cdot n.$$

using the following rules given by the *recurrence relations*<sup>1</sup>:

*Rule for addition of natural numbers:*

$$(1.1.9) \quad n + 1 := s(n), \quad \forall n \in \mathbb{N}, \quad \text{and}$$

$$(1.1.10) \quad n + s(m) := s(n + m), \quad \forall n, m \in \mathbb{N}.$$

*Rule for multiplication of natural numbers:*

$$(1.1.11) \quad n \cdot 1 := n, \quad \forall n \in \mathbb{N}, \quad \text{and}$$

$$(1.1.12) \quad n \cdot s(m) := (n \cdot m) + n, \quad \forall n, m \in \mathbb{N}.$$

**Lemma 1.1.13.** The above rules (1.1.9)–(1.1.10) defines a unique binary operation on  $\mathbb{N}$ , called *addition of natural numbers* satisfying those properties.

*Proof.* To check uniqueness of the binary operation  $+$  satisfying the properties (1.1.9)–(1.1.10), let  $\oplus : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  be any binary operation on  $\mathbb{N}$  satisfying the following properties:

$$(A') \quad n \oplus 1 = s(n), \quad \forall n \in \mathbb{N}, \quad \text{and}$$

$$(B') \quad n \oplus s(m) = s(n \oplus m), \quad \forall n, m \in \mathbb{N}.$$

Let  $m \in \mathbb{N}$  be arbitrary but fixed after choice. Let  $A := \{n \in \mathbb{N} : m + n = m \oplus n\} \subseteq \mathbb{N}$ . Since  $m + 1 = s(m) = m \oplus 1$ ,  $1 \in A$ . If  $n \in A$ , then  $m + n = m \oplus n$ , and so  $m + s(n) = s(m + n) =$

<sup>1</sup>A relation that recalls itself repeatedly to generate its complete meaning.

$s(m \oplus n) = m \oplus s(n)$ . Therefore,  $s(n) \in A$ , and hence by principle of mathematical induction (see (P6) in Axiom 1.1.1) we have  $A = \mathbb{N}$ . This proves uniqueness of the binary operation  $+$  on  $\mathbb{N}$ .

Let  $n \in \mathbb{N}$  be arbitrary but fixed after choice. Let

$$T_n := \{m \in \mathbb{N} : n + m \text{ is defined}\}.$$

Clearly  $T_n \subseteq \mathbb{N}$ . We want to show that  $T_n = \mathbb{N}$ . Now  $1 \in T_n$  by axiom (1.1.9). If  $m \in T_n$ , then  $n + m$  is defined, and so by axiom 1.1.10  $n + s(m)$  is defined. So  $s(m) \in T_n$ . Then by principle of mathematical induction (see (P6) in Peano's Axiom 1.1.1) we have  $T_n = \mathbb{N}$ .  $\square$

**Lemma 1.1.14.** *The above rules (1.1.11)–(1.1.12) defines a unique binary operation on  $\mathbb{N}$ , called the multiplication of natural numbers satisfying those properties.*

*Proof.* Left as an exercise.  $\square$

Now you know why and how you could add and multiply any two natural numbers!

**Definition 1.1.15.** Let  $*$  :  $S \times S \rightarrow S$  be a binary operation on a set  $S$ . We say that  $*$

- (i) is *associative* if  $(a * b) * c = a * (b * c)$ ,  $\forall a, b, c \in S$ ;
- (ii) is *commutative* if  $a * b = b * a$ ,  $\forall a, b \in S$ ;
- (iii) *distributes* over a binary operation  $\boxplus : S \times S \rightarrow S$  if for all  $a, b, c \in S$  we have

$$\begin{aligned} a * (b \boxplus c) &= (a * b) \boxplus (a * c), \\ (a \boxplus b) * c &= (a * c) \boxplus (b * c). \end{aligned}$$

The following result is well-known, however, it is strongly recommended to verify these in details purely using Peano's Axioms 1.1.1, and the axioms (or, definition) for addition and multiplication (1.1.9)–(1.1.12).

**Theorem 1.1.16.** *For all  $a, b, c \in \mathbb{N}$ , the following statements hold.*

- (i) *Associativity for addition:*  $(a + b) + c = a + (b + c)$ .
- (ii) *Commutativity for addition:*  $a + b = b + a$ .
- (iii) *Left distribution of multiplication over addition:*  $a \cdot (b + c) = (a \cdot b) + (a \cdot c)$ .
- (iv) *Right distribution of multiplication over addition:*  $(a + b) \cdot c = (a \cdot c) + (b \cdot c)$ .
- (v) *Commutativity for multiplication:*  $a \cdot b = b \cdot a$ .
- (vi) *Associativity for multiplication:*  $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ .

*Proof.* (i) *Proof of associativity of addition:* Let  $a, b \in \mathbb{N}$  be arbitrary but fixed after choices. Let

$$T_{a,b} := \{c \in \mathbb{N} : a + (b + c) = (a + b) + c\}.$$

Clearly  $T_{a,b} \subseteq \mathbb{N}$ . To prove associativity for addition, we need to show that  $T_{a,b} = \mathbb{N}$ . Since

$$\begin{aligned} a + (b + 1) &= a + s(b), \text{ by axiom (1.1.9).} \\ &= s(a + b), \text{ by axiom 1.1.10.} \\ &= (a + b) + 1, \text{ by axiom 1.1.9,} \end{aligned}$$

we conclude that  $1 \in T_{a,b}$ . Suppose that  $c \in T_{a,b}$  be arbitrary. Then

$$\begin{aligned} a + (b + s(c)) &= a + s(b + c), \text{ by axiom (1.1.10).} \\ &= s(a + (b + c)), \text{ by axiom (1.1.10).} \\ &= s((a + b) + c), \text{ by axiom (1.1.10).} \\ &= (a + b) + s(c), \text{ by axiom (1.1.10).} \end{aligned}$$

Therefore,  $s(c) \in T_{a,b}$ . Then by principle of mathematical induction (see (P6) in Peano's Axiom 1.1.1) we have  $T_{a,b} = \mathbb{N}$ . (Now you know why  $1 + (2 + 3) = (1 + 2) + 3$ .)

- (ii) *Proof of commutativity of addition:* For each  $a \in \mathbb{N}$ , let  $S_a := \{b \in \mathbb{N} : a + b = b + a\} \subseteq \mathbb{N}$ . We first show that  $S_1 = \mathbb{N}$ . Clearly  $1 \in S_1$ . If  $b \in S_1$ , then

$$\begin{aligned} s(b) + 1 &= s(s(b)), \text{ by axiom 1.1.9.} \\ &= s(b + 1), \text{ by axiom 1.1.9.} \\ &= s(1 + b), \text{ since } b \in S_1 \text{ by assumption.} \\ &= 1 + s(b), \text{ by axiom 1.1.10.} \end{aligned}$$

Therefore,  $s(b) \in S_1$ , and hence  $S_1 = \mathbb{N}$  by (P6) in Axiom 1.1.1. Now let  $a \in \mathbb{N}$  be arbitrary but fixed after choice. Since  $S_1 = \mathbb{N}$ , we have  $1 \in S_a$ . If  $b \in S_a$ , then  $a + b = b + a$ , and so we have

$$\begin{aligned} a + s(b) &= s(a + b), \text{ by axiom (1.1.10).} \\ &= s(b + a), \text{ since } b \in S_a \text{ by assumption.} \\ &= (b + a) + 1, \text{ by axiom (1.1.9).} \\ &= 1 + (b + a), \text{ since } b + a \in \mathbb{N} = S_1. \\ &= (1 + b) + a, \text{ using associativity of addition.} \\ &= (b + 1) + a, \text{ since } b \in \mathbb{N} = S_1. \\ &= s(b) + a, \text{ by axiom (1.1.9).} \end{aligned}$$

Then  $s(b) \in S_a$ , and hence by (P6) in Peano's Axiom 1.1.1 we have  $S_a = \mathbb{N}$ .

- (iii) *Proof of left distribution of multiplication over addition:* Let  $a, b \in \mathbb{N}$  be arbitrary but fixed after choices. Let

$$D_{a,b} := \{c \in \mathbb{N} : a \cdot (b + c) = (a \cdot b) + (a \cdot c)\} \subseteq \mathbb{N}.$$

We need to show that  $D_{a,b} = \mathbb{N}$ . Note that,

$$\begin{aligned} a \cdot (b + 1) &= a \cdot s(b), \text{ by axiom (1.1.9);} \\ &= (a \cdot b) + a, \text{ by axiom (1.1.12);} \\ &= (a \cdot b) + (a \cdot 1), \text{ by axiom (1.1.9).} \end{aligned}$$

Therefore,  $1 \in D_{a,b}$ . Suppose that  $c \in D_{a,b}$ . Then

$$\begin{aligned} a \cdot (b + s(c)) &= a \cdot s(b + c), \text{ by axiom (1.1.10);} \\ &= a \cdot (b + c) + a, \text{ by axiom (1.1.12);} \\ &= ((a \cdot b) + (a \cdot c)) + a, \text{ since } c \in D_{a,b} \text{ by assumption;} \\ &= (a \cdot b) + ((a \cdot c) + a), \text{ by associativity for addition;} \\ &= (a \cdot b) + (a \cdot s(c)), \text{ by axiom (1.1.12).} \end{aligned}$$

So  $s(c) \in D_{a,b}$ . Therefore, by (P6) of Axiom 1.1.1 we have  $D_{a,b} = \mathbb{N}$ .

- (iv) *Proof of right distribution of multiplication over addition:* Left as an exercise.

(v) *Proof of commutativity of multiplication:* Given  $a \in \mathbb{N}$ , let

$$S_a := \{b \in \mathbb{N} : a \cdot b = b \cdot a\}.$$

We first consider the case  $a = 1$ . Clearly  $1 \in S_1$ . If  $b \in S_1$ , then

$$\begin{aligned} 1 \cdot s(b) &= 1 \cdot (b + 1), \text{ by axiom (1.1.9).} \\ &= (1 \cdot b) + (1 \cdot 1), \text{ by left distribution of multiplication over addition.} \\ &= (b \cdot 1) + (1 \cdot 1), \text{ since } b \in S_1. \\ &= (b + 1) \cdot 1, \text{ by right distribution of multiplication over addition.} \\ &= s(b) \cdot 1, \text{ by axiom (1.1.9).} \end{aligned}$$

Thus,  $s(b) \in S_1$ . Therefore, by principle of mathematical induction we have  $S_1 = \mathbb{N}$ . Now assume that  $a \neq 1$ . Since  $S_1 = \mathbb{N}$ , we have  $1 \in S_a$ . Suppose that  $b \in S_a$ . Then

$$\begin{aligned} a \cdot s(b) &= (a \cdot b) + a, \text{ by axiom (1.1.12).} \\ &= (b \cdot a) + (1 \cdot a), \text{ since } 1 \in S_a \Rightarrow 1 \cdot a = a \cdot 1 = a. \\ &= (b + 1) \cdot a, \text{ by Theorem 1.1.16 (iv).} \\ &= s(b) \cdot a, \text{ by axiom 1.1.9.} \end{aligned}$$

So  $s(b) \in S_a$ , and hence  $S_a = \mathbb{N}$  by principle of mathematical induction.

(vi) *Proof of associativity of multiplication:* Left as an exercise! Let  $a, b \in \mathbb{N}$  be arbitrary but fixed after choice. Let

$$M_{a,b} := \{c \in \mathbb{N} : a \cdot (b \cdot c) = (a \cdot b) \cdot c\}.$$

Clearly  $M_{a,b} \subseteq \mathbb{N}$ . To prove associativity for multiplication, we need to show that  $M_{a,b} = \mathbb{N}$ . Since  $n \cdot 1 = n$ ,  $\forall n \in \mathbb{N}$  by axiom (1.1.11), we have  $a \cdot (b \cdot 1) = a \cdot b = (a \cdot b) \cdot 1$ . So  $1 \in M_{a,b}$ . Suppose that  $c \in M_{a,b}$ . Then

$$\begin{aligned} a \cdot (b \cdot s(c)) &= a \cdot ((b \cdot c) + b), \text{ by axiom (1.1.12).} \\ &= a \cdot (b \cdot c) + (a \cdot b), \text{ by Theorem 1.1.16 (iii).} \\ &= (a \cdot b) \cdot c + (a \cdot b), \text{ by Theorem 1.1.16 (v).} \\ &= (a \cdot b) \cdot s(c), \text{ by axiom 1.1.12.} \end{aligned}$$

Therefore,  $s(c) \in M_{a,b}$ , and hence by principle of mathematical induction we have  $M_{a,b} = \mathbb{N}$ . □

**Proposition 1.1.17.** *For each  $n, a \in \mathbb{N}$ , we have  $s^n(a) = a + n$ , where  $s^n : \mathbb{N} \rightarrow \mathbb{N}$  is the  $n$ -times composition of  $s$  with itself (e.g.,  $s^2 = s \circ s$ ,  $s^3 = s \circ s \circ s$  etc.).*

*Proof.* Let  $T := \{n \in \mathbb{N} : s^n(a) = a + n, \forall a \in \mathbb{N}\}$ . Clearly  $T \subseteq \mathbb{N}$ , and  $1 \in T$  by axiom 1.1.9. Assume that  $n \in T$ . Then  $s^{s(n)}(a) = s^{n+1}(a) = s(s^n(a)) = s(a + n) = (a + n) + 1 = a + (n + 1) = a + s(n)$ . So  $s(n) \in T$ . Then by principle of mathematical induction we have  $T = \mathbb{N}$ . □

**Lemma 1.1.18.** *Let  $a, b, n \in \mathbb{N}$ . If  $a + n = b + n$ , then  $a = b$ .*

*Proof.* Note that the successor map  $s : \mathbb{N} \rightarrow \mathbb{N}$  is injective by (P5) in Axiom 1.1.1. Since  $s^n(a) = a + n = b + n = s^n(b)$  by Proposition 1.1.17, and composition of injective maps is injective, we have  $a = b$ . □

**Exercise 1.1.19** (Cancellation for multiplication). Let  $a, b, r, \ell \in \mathbb{N}$ .

(i) If  $\ell a = \ell b$ , show that  $a = b$ .

(ii) If  $ar = br$ , show that  $a = b$ .

**Exercise 1.1.20.** Let  $a \in \mathbb{N}$ . Show that the equation  $x + a = 1$  has no solution for  $x$  in  $\mathbb{N}$ .

**Theorem 1.1.21** (Law of trichotomy for natural numbers). *Given  $a, b \in \mathbb{N}$ , exactly one of the following three conditions holds:*

(i)  $a = b$ ,

(ii)  $a = b + c$ , for some  $c \in \mathbb{N}$ , or

(iii)  $b = a + d$ , for some  $d \in \mathbb{N}$ .

*Proof.* We first show that no two conditions among (i)–(iii) can hold simultaneously. If (i) and (ii) holds simultaneously, then  $b = b + c$  implies  $s(b) = s(b + c) \Rightarrow b + 1 = (b + c) + 1 = b + (c + 1) \Rightarrow 1 = c + 1 = s(c)$ , which contradicts axiom (P4) in Peano's Axioms 1.1.1. The same argument shows that (i) and (iii) cannot hold simultaneously. If (ii) and (iii) hold simultaneously, then we have  $a = b + c = (a + c) + d = a + (c + d)$ , for some  $c, d \in \mathbb{N}$ . Then applying successor map we see that  $a + 1 = (a + (c + d)) + 1 = a + ((c + d) + 1)$ . Then by Lemma 1.1.18 we have  $1 = (c + d) + 1 = s(c + d)$ , which contradicts (P4) in Peano's Axioms 1.1.1. Therefore, no two conditions among (i)–(iii) can hold simultaneously.

We now show that at least one of (i)–(iii) holds. For each  $a \in \mathbb{N}$ , let

$$S_a := \{b \in \mathbb{N} : \text{at least one of (i) or (ii) or (iii) holds}\}.$$

Consider the case  $a = 1$ . Clearly  $1 \in S_1$ . Suppose that  $b \in S_1$ . Then  $s(b) = b + 1 = a + b$  satisfies condition (iii), and so  $s(b) \in S_1$ . Then by (P6) in Peano's Axioms we have  $S_1 = \mathbb{N}$ . Suppose that  $a \in \mathbb{N} \setminus \{1\}$  be arbitrary but fixed after choice. Since  $b = 1$  satisfies  $a = s(p(a)) = p(a) + 1 = p(a) + b$ , with  $p(a) \in \mathbb{N}$ , the condition (ii) holds for  $b = 1$ , and so  $1 \in S_a$ . Suppose that  $b \in S_a$ . Then we have the following cases:

- (I) If  $a = b$ , then  $s(b) = b + 1 = a + 1$ , and so  $s(b)$  satisfies condition (iii). So  $s(b) \in S_a$ .
- (II) If  $a = b + c$ , for some  $c \in \mathbb{N}$ , then  $a = s(b)$  or  $a = s(b) + p(c)$  depending on whether  $c = 1$  or  $c \in \mathbb{N} \setminus \{1\}$ , respectively. So in both cases,  $s(b) \in S_a$ .
- (III) If  $b = a + d$ , for some  $d \in \mathbb{N}$ , then  $s(b) = b + 1 = a + (d + 1)$  satisfies condition (iii), and hence  $s(b) \in S_a$ .

Therefore,  $S_a = \mathbb{N}$  by principle of mathematical induction.  $\square$

The law of trichotomy in Theorem 1.1.21 allow us to define usual order relation “ $<$ ” on  $\mathbb{N}$  as follow.

**Definition 1.1.22.** Given  $a, b \in \mathbb{N}$ , we define  $a < b$  if  $\exists c \in \mathbb{N}$  such that  $a + c = b$ . If  $a < b$ , we say that “ $a$  is strictly less than  $b$ ”.

Note that “ $<$ ” is a relation on  $\mathbb{N}$  which is neither reflexive nor symmetric or anti-symmetric. We show that it is a transitive relation on  $\mathbb{N}$ . If  $a < b$  and  $b < c$ , then  $a + r = b$  and  $b + s = c$ , for some  $r, s \in \mathbb{N}$ , and then  $a + (r + s) = (a + r) + s = b + s = c$  shows that  $a < c$ . If  $a < b$  we say that “ $a$  is less than  $b$ ”. The relation “ $<$ ” is called the *usual ordering relation* on  $\mathbb{N}$ . Define another relation “ $\leq$ ” on  $\mathbb{N}$  by setting

$$a \leq b \text{ if either } a = b \text{ or } a < b.$$

If  $a \leq b$ , we say that “ $a$  is less than or equal to  $b$ ”. It is easy to see that “ $\leq$ ” is reflexive and transitive. We show that “ $\leq$ ” is anti-symmetric, and hence is a partial order relation on  $\mathbb{N}$ .

Suppose that  $a, b \in \mathbb{N}$  with  $a \leq b$  and  $b \leq a$ . We want to show that  $a = b$ . Suppose on the contrary that  $a \neq b$ . Then we must have  $a < b$  and  $b < a$ . Then there exist  $c, d \in \mathbb{N}$  such that  $a + c = b$  and  $b + d = a$ . Then

$$\begin{aligned}
 & (b + d) + c = a + c = b \\
 \Rightarrow & b + (d + c) = b, \text{ using associativity of addition.} \\
 \Rightarrow & s(b + (d + c)) = s(b), \text{ applying successor map.} \\
 \Rightarrow & (b + (d + c)) + 1 = b + 1, \text{ using axiom 1.1.10.} \\
 \Rightarrow & b + ((d + c) + 1) = b + 1, \text{ using associativity of addition.} \\
 \Rightarrow & (d + c) + 1 = 1, \text{ using Lemma 1.1.18.} \\
 \Rightarrow & s(d + c) = 1.
 \end{aligned}$$

This contradicts axiom (P4) in Peano's Axioms 1.1.1. Therefore, we must have  $a = b$  as required.

**Definition 1.1.23.** A partial order relation  $\rho$  on a set  $S$  is called a *total order* if for any two elements  $a, b \in S$ , at least one of  $a \rho b$  and  $b \rho a$  holds. A non-empty set  $S$  together with a total order relation is called a *well-ordered set*.

As an immediate consequence of the law of trichotomy of natural numbers (Theorem 1.1.21) we see that " $\leq$ " is a total order relation on  $\mathbb{N}$ , and hence  $(\mathbb{N}, \leq)$  is a well-ordered set.

**Theorem 1.1.24.** *The following are equivalent.*

(i) *Principle of mathematical induction (regular version): Let  $S \subseteq \mathbb{N}$  be such that*

- (a)  $1 \in S$ , and
- (b) *for each  $n \in \mathbb{N}$ ,  $n \in S$  implies  $s(n) \in S$ .*

*Then  $S = \mathbb{N}$ .*

(ii) *Principle of mathematical induction (strong version): Let  $T \subseteq \mathbb{N}$  be such that*

- (a')  $1 \in T$ , and
- (b') *for each  $n \in \mathbb{N}$ ,  $J_n := \{k \in \mathbb{N} : k \leq n\} \subseteq T$  implies  $s(n) \in T$ .*

*Then  $T = \mathbb{N}$ .*

*Proof.* (i)  $\Rightarrow$  (ii): Suppose that the conditions (a') and (b') holds for  $T \subseteq \mathbb{N}$ . Since  $1 \in T$  by (a'), to show  $T = \mathbb{N}$  using the regular version of principle of mathematical induction (i), it is enough to show that for each  $n \in \mathbb{N}$ , the statement

$$P_n : "n \in T \text{ implies } s(n) \in T."$$

holds. Consider the set

$$S := \{n \in \mathbb{N} : P_k \text{ holds, } \forall k \leq n\} \subseteq \mathbb{N}.$$

Since  $1 \in T$  by (a'), we have  $J_1 = \{1\} \subseteq T$ , and hence by (b') we have  $s(1) \in T$ . Therefore,  $P_1$  holds, and so  $1 \in S$ . Let  $n \in S$  be arbitrary but fixed after choice. Then  $P_1, \dots, P_n$  hold, and hence we have  $J_{s(n)} = \{k \in \mathbb{N} : k \leq s(n)\} \subseteq T$ . Then by the condition (b') we have  $s(s(n)) \in T$ , and hence  $P_{s(n)}$  holds. Therefore,  $P_k$  holds,  $\forall k \leq s(n)$ , and hence  $s(n) \in S$ . Then by (i) we have  $S = \mathbb{N}$ . Thus,  $T = \mathbb{N}$ .

(ii)  $\Rightarrow$  (i): Let  $S \subseteq \mathbb{N}$  be such that  $1 \in S$ , and  $n \in S$  implies  $s(n) \in S$ . To show  $S = \mathbb{N}$  using the strong version of principle of mathematical induction (ii), we just need to ensure that for each  $n \in \mathbb{N}$ , if  $J_n \subseteq S$  then  $s(n) \in S$ . But this follows because  $n \in J_n$  implies that  $n \in S$ , and so  $s(n) \in S$  by (a). Then by (ii) we have  $S = \mathbb{N}$ . This proves (i).  $\square$

**Theorem 1.1.25.** *The following are equivalent.*

(i) *Principle of Mathematical Induction (strong version):* Let  $S \subseteq \mathbb{N}$  be such that

(a)  $1 \in S$ , and

(b) for each  $n \in \mathbb{N}$  with  $n > 1$ , if  $\{k \in \mathbb{N} : k < n\} \subseteq S$  then  $n \in S$ .

Then  $S = \mathbb{N}$ .

(ii) *Well-ordering principle of  $(\mathbb{N}, \leq)$ :* Any non-empty subset of  $\mathbb{N}$  has a least element.

*Proof.* (i)  $\Rightarrow$  (ii): Suppose on the contrary that there is a non-empty subset  $S \subseteq \mathbb{N}$  which has no least element. Let

$$T := \mathbb{N} \setminus S = \{n \in \mathbb{N} : n \notin S\}.$$

Since 1 is the least element of  $\mathbb{N}$ , we have  $1 \notin S$ ; for otherwise 1 would be the least element of  $S$ . Therefore,  $1 \in T$  and hence  $T$  is a non-empty subset of  $\mathbb{N}$ . Let  $n \in \mathbb{N}$  with  $n > 1$ , and suppose that for any  $k \in \mathbb{N}$  with  $k < n$ , we have  $k \in T$ . Then  $n \notin S$ , for otherwise  $n$  would be the least element of  $S$ . So  $n \in T$ . Then by principle of mathematical induction (strong version), we have  $T = \mathbb{N}$ . This contradicts our assumption that  $S$  is non-empty. So  $S$  must have a least element.

(ii)  $\Rightarrow$  (i): Let  $S \subseteq \mathbb{N}$  be such that

(a)  $1 \in S$ , and

(b) for each  $n \in \mathbb{N}$  with  $n > 1$ , if  $\{k \in \mathbb{N} : k < n\} \subseteq S$  then  $n \in S$ .

Assuming well-ordering principle of  $(\mathbb{N}, \leq)$ , we want to show that  $S = \mathbb{N}$ . Suppose on the contrary that  $S \neq \mathbb{N}$ . Then  $T := \mathbb{N} \setminus S$  is a non-empty subset of  $\mathbb{N}$ , and so by (i) it has a least element, say  $n \in T$ . Since  $1 \in S$  by assumption,  $n > 1$ . Since  $n$  is the least element of  $T$ , for any  $k \in \mathbb{N}$  with  $k < n$ , we have  $k \in \mathbb{N} \setminus T = S$ . Then by property (b) of  $S$  we have  $n \in S$ , which is a contradiction. This completes the proof.  $\square$

## 1.2 Integers: Construction & Basic Operations

Let  $a, b \in \mathbb{N}$ . Suppose that we want to solve the equation

$$(1.2.1) \quad x + a = b$$

to find  $x$ . If  $a < b$  in  $\mathbb{N}$ , then there is  $r \in \mathbb{N}$  such that  $b = r + a$ . If there is another number  $s \in \mathbb{N}$  such that  $b = s + a$ , then  $r + a = s + a$  implies  $r = s$ . So the solution of the equation (1.2.1) exists and is unique; we denote this solution by  $a - b \in \mathbb{N}$ . Now the problem is if  $a \leq b$ , we don't have any solution of this equation in  $\mathbb{N}$ . This forces us to enlarge our natural number system to a bigger number system where we can find solutions to such linear equations.

Define a relation  $\sim$  on the Cartesian product  $\mathbb{N} \times \mathbb{N}$  by setting

$$(1.2.2) \quad (a, b) \sim (c, d), \quad \text{if } a + d = b + c.$$

It is an easy exercise to show that  $\sim$  is an equivalence relation on  $\mathbb{N} \times \mathbb{N}$ . The  $\sim$ -equivalence class of  $(a, b) \in \mathbb{N} \times \mathbb{N}$  is the subset

$$(1.2.3) \quad [(a, b)] := \{(c, d) \in \mathbb{N} \times \mathbb{N} \mid (a, b) \sim (c, d)\}.$$

Let  $\mathbb{Z} := \{[(a, b)] : a, b \in \mathbb{N}\}$  be the associated set of all  $\sim$ -equivalence classes. The idea is to think of the equivalence class  $[(a, b)]$  to be the solution of the equation  $x + b = a$ . The elements of  $\mathbb{Z}$  are called *integers*, and  $\mathbb{Z}$  is called the *set of all integers*.

Define a map

$$\iota : \mathbb{N} \rightarrow \mathbb{Z}$$



by

$$\iota(n) = [(s(n), 1)], \quad \forall n \in \mathbb{N}.$$

Then  $\iota(n) = \iota(m) \Rightarrow [(s(n), 1)] = [(s(m), 1)] \Rightarrow s(n) + 1 = s(m) + 1 \Rightarrow s(s(n)) = s(s(m)) \Rightarrow s(n) = s(m) \Rightarrow n = m$ , since  $s : \mathbb{N} \rightarrow \mathbb{N}$  is an injective map. Therefore,  $\iota : \mathbb{N} \rightarrow \mathbb{Z}$  is an injective map, and hence we can use it to identify  $\mathbb{N}$  as a subset of  $\mathbb{Z}$ . For notational simplicity, we may denote by  $n$  the element  $[(s(n), 1)] \in \mathbb{Z}$ , for all  $n \in \mathbb{N}$ .

Define a binary operation on  $\mathbb{Z}$ , called *addition of integers*, by

$$(1.2.4) \quad [(a, b)] + [(c, d)] := [(a + c, b + d)], \quad \forall [(a, b)], [(c, d)] \in \mathbb{Z}.$$

Note that, if  $(a, b) \sim (a', b')$  and  $(c, d) \sim (c', d')$ , then  $(a + c, b + d) \sim (a' + c', b' + d')$ . Therefore, we have a well-defined binary operation  $+$  on  $\mathbb{Z}$ .

**Exercise 1.2.5.** Show that the addition of integers is associative and commutative.

**Exercise 1.2.6.** Verify that,

$$\iota(m + n) = \iota(m) + \iota(n), \quad \forall m, n \in \mathbb{N}.$$

Therefore, the addition operation on integers preserves the addition operation on natural numbers defined earlier.

Note that, the element  $[(1, 1)] \in \mathbb{Z}$  satisfies

$$[(a, b)] + [(1, 1)] = [(a, b)] = [(1, 1)] + [(a, b)].$$

We denote by 0 (pronounced as *zero*) the element  $[(1, 1)] \in \mathbb{Z}$ . Since

$$[(s(n), 1)] + [(1, s(n))] = [(1, 1)] = 0, \quad \forall n \in \mathbb{N},$$

for notational simplicity (for peaceful working notations), we denote by  $-n$  the element  $[(1, s(n))] \in \mathbb{Z}$ , for all  $n \in \mathbb{N}$ . The element of  $\mathbb{Z}$  of the form  $n$  and  $-n$  are called *positive integers* and *negative integers*, respectively.

**Exercise 1.2.7.** The subsets  $\mathbb{Z}^- := \{[(1, s(n))] : n \in \mathbb{N}\}$ ,  $\{0\} := \{[(1, 1)]\}$  and  $\mathbb{Z}^+ := \{[(s(n), 1)] : n \in \mathbb{N}\}$  are mutually disjoint, and their union is  $\mathbb{Z}$ . As a result, we may write the set  $\mathbb{Z}$  as

$$\mathbb{Z} = \{-n : n \in \mathbb{N}\} \cup \{0\} \cup \mathbb{N}.$$

The elements of  $\mathbb{Z}^-$  and  $\mathbb{Z}^+$  are called the *negative integers* and the *positive integers*, respectively.

We define another binary operation on  $\mathbb{Z}$ , called the *product operation*, by

$$(1.2.8) \quad [(a, b)] \cdot [(c, d)] := [(ac + bd, ad + bc)], \quad \forall [(a, b)], [(c, d)] \in \mathbb{Z}.$$

It is easy to check that, if  $(a, b) \sim (a', b')$  and  $(c, d) \sim (c', d')$ , then  $(ac + bd, ad + bc) \sim (a'c' + b'd', a'd' + b'c')$ , and hence the product operation is well-defined. One can easily check that,

- (i)  $[(a, b)] \cdot [(c, d)] = [(c, d)] \cdot [(a, b)],$
- (ii)  $[(s(m), 1)] \cdot [(s(n), 1)] = [(s(mn), 1)].$

**Remark 1.2.9.** With the above definitions and notations, one can check that the binary operations addition and multiplication of integers are associative, commutative, and multiplication distributes over addition. In other words, the following properties hold.

- (i)  $(a + b) + c = a + (b + c), \quad \forall a, b, c \in \mathbb{Z};$

- (ii)  $(ab)c = a(bc), \forall a, b, c \in \mathbb{Z};$
- (iii)  $a + b = b + a, \forall a, b \in \mathbb{Z};$
- (iv)  $ab = ba, \forall a, b \in \mathbb{Z};$
- (v)  $a(b + c) = (ab) + (ac), \forall a, b, c \in \mathbb{Z};$
- (vi)  $(a + b)c = (ac) + (bc), \forall a, b, c \in \mathbb{Z}.$

**Exercise 1.2.10.** Let  $n \in \mathbb{Z}$ . If  $a + n = b + n$ , for some  $a, b \in \mathbb{Z}$ , show that  $a = b$ .

We define the *usual ordering relation* “ $\leq$ ” on  $\mathbb{Z}$  as follow: given  $m, n \in \mathbb{Z}$ , we define

$$m \leq n \text{ if } \exists r \in \mathbb{N} \cup \{0\} \text{ such that } m + r = n.$$

**Exercise 1.2.11.** Verify that  $(\mathbb{Z}, \leq)$  is a well-ordered set.

### 1.3 Division Algorithm

Recall that the *well-ordering principle of natural numbers* says that any non-empty subset  $S$  of  $\mathbb{N}$  has a least element. This means, there exists  $n \in S$  such that  $n \leq m$ , for all  $m \in S$ . This statement is equivalent to the *principle of mathematical induction*, which says that if  $S \subseteq \mathbb{N}$  is such that  $1 \in S$ , and for each  $n \in \mathbb{N}, n \in S \Rightarrow n + 1 \in S$ , then  $S = \mathbb{N}$ .

**Theorem 1.3.1** (Division algorithm). *Given  $a, d \in \mathbb{Z}$  with  $d > 0$ , there exists unique  $q, r \in \mathbb{Z}$  with  $0 \leq r < d$  such that  $a = qd + r$ .*

*Proof.* We first show uniqueness of  $q$  and  $r$ . Suppose that we have another pair of integers  $q', r' \in \mathbb{Z}$  such that  $0 \leq r' < d$  and  $a = q'd + r'$ . Without loss of generality we may assume that  $r \leq r'$ . Then  $qd + r = a = q'd + r'$  implies  $r' - r = (q - q')d$ . Since  $0 \leq r \leq r' < d$ , we have  $0 \leq (q - q')d = r' - r < d$ . Therefore,  $(q - q')d$  is a non-negative integer which is strictly less than  $d$  and is a multiple of  $d$ . This is possible only if  $(q - q')d = 0$ . Since  $d \neq 0$ , we must have  $q = q'$ , and hence  $r = r'$ . This proves uniqueness part.

To show existence, consider the set

$$S := \{a - dq : q \in \mathbb{N}\} \cap \mathbb{N}.$$

Since  $d > 0$ , choosing  $q$  sufficiently small we can ensure that  $a - dq \in \mathbb{N}$ , and hence  $S \neq \emptyset$ . Then by well-ordering principle of  $(\mathbb{N}, \leq)$ ,  $S$  has a least element, say  $r_0$ . Then  $0 \leq r_0 = a - dq_0$ , for some  $q_0 \in \mathbb{Z}$ . We claim that  $r_0 < d$ . If not, then  $r_0 \geq d$  and hence  $0 \leq r_0 - d = a - d(q_0 + 1)$  implies that  $r_0 - d \in S$ . Since  $d > 0$ , it contradicts the fact that  $r_0$  is the least element of  $S$ . Therefore, we must have  $r_0 < d$ . This completes the proof.  $\square$

**Definition 1.3.2.** The *absolute value* of  $n \in \mathbb{Z}$  is the integer  $|n|$  defined by

$$|n| := \begin{cases} n, & \text{if } n \geq 0, \\ -n, & \text{if } n < 0. \end{cases}$$

**Corollary 1.3.3.** *Given  $a, d \in \mathbb{Z}$  with  $d \neq 0$ , there exists unique  $q, r \in \mathbb{Z}$  with  $0 \leq r < |d|$  such that  $a = dq + r$ .*

*Proof.* If  $d > 0$ , this is precisely Theorem 1.3.1. If  $d < 0$ , then  $d' := -d > 0$ , and so by division algorithm (Theorem 1.3.1) we find unique integers  $q, r \in \mathbb{Z}$  with  $0 \leq r < d'$  such that  $a = d'q + r$ . Then the integers  $q' := -q$  and  $r$  satisfies  $0 \leq r < |d|$  with  $a = q'd + r$ .  $\square$

**Definition 1.3.4.** Given  $n, d \in \mathbb{Z}$ , with  $d \neq 0$ , we say that  $d$  divides  $n$ , written as  $d \mid n$ , if there is an element  $q \in \mathbb{Z}$  such that  $n = qd$ . Given finitely many integers  $a_1, \dots, a_n \in \mathbb{Z}$ , which are not all zero, we define their *greatest common divisor* to be a positive integer  $d \in \mathbb{Z}^+$  such that

- (i)  $d$  divides each of the numbers  $a_1, \dots, a_n$ , and
- (ii) if an integer  $r$  divides  $a_i$ , for all  $i = 1, \dots, n$ , then  $r$  divides  $d$ .

**Remark 1.3.5.** Given a finite number of integers  $a_1, \dots, a_n \in \mathbb{Z}$ , if  $d$  and  $d'$  are two greatest common divisors of  $a_1, \dots, a_n$ , then  $d \mid d'$  and  $d' \mid d$  implies  $d \in \{d', -d'\}$ . Since both  $d$  and  $d'$  are positive integers, we must have  $d = d'$ . Therefore, the greatest common divisor of  $a_1, \dots, a_n$  is unique, and we denote it by  $\gcd(a_1, \dots, a_n)$ . However, it is not yet clear if  $\gcd(a_1, \dots, a_n)$  exists in  $\mathbb{N}$ . This requires a proof.

**Lemma 1.3.6.** Given  $m, n \in \mathbb{Z}$ , not all zero, the greatest common divisor  $\gcd(m, n)$  exists in  $\mathbb{N}$ . Moreover, there exist  $a, b \in \mathbb{Z}$  such that  $\gcd(m, n) = am + bn$ .

*Proof.* Let  $S := \{am + bn : a, b \in \mathbb{Z}\}$ . Since at least one of  $m$  and  $n$  is non-zero, there is a non-zero element, say  $x$ , in  $S$ . Then  $x = am + bn$ , for some  $a, b \in \mathbb{Z}$ . If  $x < 0$ , then  $-x = (-a)m + (-b)n \in S \cap \mathbb{N}$ . Therefore,  $S \cap \mathbb{N}$  is a non-empty subset of  $\mathbb{N}$ . Then by well-ordering principle of  $\mathbb{N}$ , the non-empty subset  $S \cap \mathbb{N}$  has a least element, say  $d$ . Then  $d = a_0m + b_0n$ , for some  $a_0, b_0 \in \mathbb{Z}$ . We claim that  $d = \gcd(m, n)$ .

If  $r \mid m$  and  $r \mid n$ , then  $r \mid (a_0m + b_0n)$  and so  $r \mid d$ . Now we need to show that  $d \mid m$  and  $d \mid n$ . Let  $x \in S$  be arbitrary. Then  $x = am + bn \in S$ , for some  $a, b \in \mathbb{Z}$ . By division algorithm we can find  $q, r \in \mathbb{Z}$  with  $0 \leq r < d$  such that  $x = qd + r$ . Then  $am + bn = x = qd + r = q(a_0m + b_0n) + r$  implies  $r = (a - qa_0)m + (b - qb_0)n \in S$ . Since  $0 \leq r < d$  and  $d$  is the smallest positive integer in  $S \cap \mathbb{N}$ , we must have  $r = 0$ . Therefore,  $x = qd$  and hence  $d \mid x$ , for all  $x \in S$ . In particular, choosing  $(a, b) \in \{(1, 0), (0, 1)\}$ , we see that  $d \mid m$  and  $d \mid n$ . This completes the proof.  $\square$

**Definition 1.3.7.** Given  $m, n \in \mathbb{Z}$ , we say that  $m$  and  $n$  are *relatively prime* (or, *coprime*) if  $\gcd(m, n) = 1$ .

**Corollary 1.3.8.** Two integers  $m$  and  $n$  are coprime if and only if there exists  $a, b \in \mathbb{Z}$  such that  $am + bn = 1$ .

*Proof.* If  $\gcd(m, n) = 1$ , then by above Lemma 1.3.6, there exists  $a, b \in \mathbb{Z}$  such that  $am + bn = 1$ . Conversely, suppose that  $am + bn = 1$ , for some  $a, b \in \mathbb{Z}$ . If  $d = \gcd(m, n)$ , then  $d \mid m$  and  $d \mid n$  implies  $d \mid 1$ . Then  $d \in \{1, -1\}$ . Since  $d > 0$ , we have  $d = 1$ .  $\square$

**Exercise 1.3.9.** Given a finite number of integers  $a_1, \dots, a_n$ , not all zero, show that  $\gcd(a_1, \dots, a_n)$  exists in  $\mathbb{N}$ .

**Definition 1.3.10.** An integer  $p \in \mathbb{Z}$  is said to be a *prime number* if  $p > 1$  and its only divisors in  $\mathbb{Z}$  are  $\pm 1, \pm p$ .

**Exercise 1.3.11** (Principle of mathematical induction). Fix  $n_0 \in \mathbb{N}$ . Prove that the following are equivalent.

- (i) *Regular version:* Let  $S \subseteq \mathbb{N}$  be such that
  - (a)  $n_0 \in S$ , and
  - (b) for any  $n \in \mathbb{N}$  with  $n \geq n_0$ , if  $n \in S$  then  $n + 1 \in S$ .

Then  $S = \{n \in \mathbb{N} : n \geq n_0\}$ .

- (ii) *Strong version:* Let  $T \subseteq \mathbb{N}$  be such that
  - (a')  $n_0 \in T$ , and

(b') for any  $n \in \mathbb{N}$  with  $n \geq n_0$ , if  $\{k \in \mathbb{N} : n_0 \leq k \leq n\} \subseteq T$  then  $n + 1 \in T$ .

Then  $T = \{n \in \mathbb{N} : n \geq n_0\}$ .

Assuming well-ordering principle of  $(\mathbb{N}, \leq)$  show that the above two versions of induction holds true.

**Theorem 1.3.12** (Fundamental theorem of Arithmetic). *Given a positive integer  $n > 1$ , there exists a unique factorization of  $n$  as a product of positive integer powers of prime numbers. More precisely, there exist finite number of unique prime numbers  $p_1, \dots, p_k \in \mathbb{N}$  with  $p_1 > \dots > p_k$  and positive integers  $\alpha_1, \dots, \alpha_k \in \mathbb{N}$  such that  $n = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$ .*

## Chapter 2

# Group Theory

## 2.1 Group

A *binary operation* on a set  $A$  is a map  $*$  :  $A \times A \rightarrow A$ ; given  $(a, b) \in A \times A$  its image under the map  $*$  is denoted by  $a * b$ . We consider some examples of non-empty set together with a natural binary operation and study list down their common properties.

**Example 2.1.1.** The set of all integers

$$\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \pm 4, \pm 5, \dots\}$$

admits a binary operation, namely addition of integers:

$$+ : \mathbb{Z} \times \mathbb{Z} \longrightarrow \mathbb{Z}, \quad (a, b) \longmapsto a + b.$$

This binary operation has the following interesting properties:

- (i)  $a + (b + c) = (a + b) + c, \forall a, b, c \in \mathbb{Z}$ ,
- (ii) there is an element  $0 \in \mathbb{Z}$  such that  $a + 0 = 0 + a = a, \forall a \in \mathbb{Z}$ ,
- (iii) for each  $a \in \mathbb{Z}$ , there exists an element  $b \in \mathbb{Z}$  (depending on  $a$ ) such that  $a + b = b + a = 0$ ; the element  $b$  is denoted by  $-a$ .

**Example 2.1.2.** A *symmetry* on a non-empty set  $X$  is a bijective map from  $X$  onto itself. The set of all symmetries of  $X$  is denoted by  $S(X)$ . Note that  $S(X)$  admits a binary operation given by composition of maps:

$$\circ : S(X) \times S(X) \longrightarrow S(X), \quad (f, g) \longmapsto g \circ f.$$

Note that

- (i) given any  $f, g, h \in S(X)$ , we have  $(f \circ g) \circ h = f \circ (g \circ h)$ .
- (ii) there is a distinguished element, the identity map  $\text{Id}_X \in S(X)$  such that  $f \circ \text{Id}_X = f = \text{Id}_X \circ f$ , for all  $f \in S(X)$ .
- (iii) given any  $f \in S(X)$ , there is a element  $g := f^{-1} \in S(X)$  such that  $f \circ g = \text{Id}_X = g \circ f$ .

**Example 2.1.3.** Fix a natural number  $n \geq 1$ , and consider the set  $\text{GL}_n(\mathbb{R})$  of all invertible  $n \times n$  matrices with entries from  $\mathbb{R}$ . Note that  $\text{GL}_n(\mathbb{R})$  admits a natural binary operation given by matrix multiplication:

$$\cdot : \text{GL}_n(\mathbb{R}) \times \text{GL}_n(\mathbb{R}) \rightarrow \text{GL}_n(\mathbb{R}), \quad (A, B) \longmapsto AB.$$

Note that

- (i) given any  $A, B, C \in \text{GL}_n(\mathbb{R})$ , we have  $(AB)C = A(BC)$ .
- (ii) there is a distinguished element, the identity matrix  $I_n \in \text{GL}_n(\mathbb{R})$  such that  $AI_n = I_nA = A$ , for all  $A \in \text{GL}_n(\mathbb{R})$ .
- (iii) given any  $A \in \text{GL}_n(\mathbb{R})$ , there is a element  $B := A^{-1} \in \text{GL}_n(\mathbb{R})$  such that  $AB = BA = I_n$ .

A non-empty set together with a binary operation satisfying the three properties listed in the above examples is a mathematical model for many important mathematical and physical systems; such a mathematical model is called a group. Here is a formal definition.

**Definition 2.1.4.** A *group* is a pair  $(G, *)$  consisting of a non-empty set  $G$  together with a binary operation

$$* : G \times G \longrightarrow G, \quad (a, b) \longmapsto a * b,$$

satisfying the following conditions:

- (G1) *Associativity*:  $a * (b * c) = (a * b) * c$ , for all  $a, b, c \in G$ .
- (G2) *Existence of neutral element*:  $\exists$  an element  $e \in G$  such that  $a * e = e * a = a$ ,  $\forall a \in G$ .
- (G3) *Existence of inverse*: for each  $a \in G$ , there exists an element  $b \in G$ , depending on  $a$ , such that  $a * b = e = b * a$ .

A *semigroup* is a pair  $(G, *)$  consisting of a non-empty set  $G$  together with an associative binary operation  $* : G \times G \rightarrow G$  (i.e., the condition (G1) holds). A *monoid* is a semigroup  $(G, *)$  satisfying the condition (G2) as above. For example,  $(\mathbb{N}, +)$  is a semigroup but not a monoid, and  $(\mathbb{Z}_{\geq 0}, +)$  is a monoid but not a group. However, we shall not deal with these two notations in this text.

- Example 2.1.5.** (i) *Trivial group*: A singleton set  $\{e\}$  with the binary operation  $e * e := e$  is a group; such a group is called a *trivial group*.
- (ii) The set  $G := \{e, a\}$ , with the binary operation  $*$  given by  $a * e = e * a = a$  and  $a * a = e$ , is a group with two elements.
- (iii) Verify that  $G := \{e, a, b\}$  together with the binary operation  $*$  given by the following multiplication table, is a group (with three elements).

$*$	$e$	$a$	$b$
$e$	$e$	$a$	$b$
$a$	$a$	$b$	$e$
$b$	$b$	$e$	$a$

TABLE 2.1.5.1: A group with 3 elements

**Remark 2.1.6.** For a group consisting of small number of elements, it is convenient to write down the associated binary operation explicitly using a table as above, known as the *Cayley table*.

- (iv) The sets  $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$  and  $\mathbb{C}$  form groups with respect to usual addition.
- (v) The set  $\mathbb{Q}^* = \mathbb{Q} \setminus \{0\}$  forms a group with respect to usual multiplication.

**Exercise 2.1.7.** Let  $(G, *)$  be a group.

- (i) *Uniqueness of neutral element*: Show that the neutral element (also known as the *identity element*)  $e \in G$  is unique.

- (ii) *Uniqueness of inverse*: Show that, for each  $a \in G$ , there is a unique element  $b \in G$  such that  $a * b = b * a = e$ . The element  $b$  is called *the inverse* of  $a$ , and denoted by the symbol  $a^{-1}$ .
- (iii) *Cancellation Law*: If  $a * c = b * c$ , for some  $a, b, c \in G$ , show that  $a = b$ .
- (iv) Let  $a, b \in G$ . Show that  $\exists$  unique  $x, y \in G$  such that  $a * x = b$  and  $y * a = b$ .

Let  $(G, *)$  be a group. We say that  $G$  is *finite* or *infinite* according as its underlying set  $G$  is finite or infinite; the cardinality of  $G$  is called the *order* of the group  $(G, *)$ , and we denote it by the symbol  $|G|$ . For notational simplicity, we write  $ab$  to mean  $a * b$ , for all  $a, b \in G$ ; and for any integer  $n \geq 1$ , we denote by  $a^n$  the  $n$ -fold product of  $a$  with itself, i.e.,

$$a^n := \underbrace{a * \cdots * a}_{n\text{-fold product of } a}.$$

For a negative integer  $n$ , we define  $a^n := (a^{-1})^{-n}$ . When there is no confusion likely to arise, we simply denote a group  $(G, *)$  by  $G$  without specifying the binary operation.

**Exercise 2.1.8.** Let  $G$  be a group.

- (i) Show that  $(a^{-1})^{-1} = a$ , for all  $a \in G$ .
- (ii) Show that  $(ab)^{-1} = b^{-1}a^{-1}$ , for all  $a, b \in G$ .
- (iii) Show that  $a^m a^n = a^{m+n}$ , for all  $m, n \in \mathbb{Z}$  and  $a \in G$ .
- (iv) Show that  $(a^m)^n = a^{mn}$ , for all  $m, n \in \mathbb{Z}$  and  $a \in G$ .
- (v) Let  $a, b \in G$  be such that  $ab = ba$ . Show that  $(ab)^n = a^n b^n$ , for all  $n \in \mathbb{Z}$ .

**Example 2.1.9.** (i) The set  $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$  of non-zero complex numbers forms a group with respect to multiplication of complex numbers.

(ii) *Circle group*: The set

$$S^1 := \{z \in \mathbb{C} : |z| = 1\}$$

forms a group with respect to multiplication of complex numbers.

(iii) *Klein four-group*: Consider the set  $K_4 = \{e, a, b, c\}$  together with the binary operation

$$* : K_4 \times K_4 \longrightarrow K_4$$

defined by the Cayley table 2.1.9.1 below. Verify that  $K_4$  is a group.

*	e	a	b	c
e	e	a	b	c
a	a	e	c	b
b	b	c	e	a
c	c	b	a	e

TABLE 2.1.9.1: Klein four group

**Exercise 2.1.10.** Define a binary operation on  $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$  by

$$(x_1, y_1) + (x_2, y_2) := (x_1 + x_2, y_1 + y_2), \quad \forall (x_1, y_1), (x_2, y_2) \in \mathbb{R}^2.$$

Verify that  $(\mathbb{R}^2, +)$  is a commutative group. Similarly, for each  $n \in \mathbb{N}$ , show that the component-wise addition of real numbers:

$$(2.1.11) \quad (a_1, \dots, a_n) + (b_1, \dots, b_n) := (a_1 + b_1, \dots, a_n + b_n), \quad \forall a_j, b_j \in \mathbb{R},$$

defines a binary operation  $+$  on  $\mathbb{R}^n$  which makes the pair  $(\mathbb{R}^n, +)$  a commutative group.

**Definition 2.1.12.** A map  $f : A \rightarrow B$  is said to be

- (i) *injective* if given any  $a_1, a_2 \in A$  with  $f(a_1) = f(a_2)$ , we have  $a_1 = a_2$ ,
- (ii) *surjective* if given any  $b \in B$ , there is an element  $a \in A$  such that  $f(a) = b$ ,
- (iii) *bijective* if  $f$  is both injective and surjective.

**Exercise 2.1.13.** Let  $A, B$  and  $C$  be three sets. Given maps  $f : A \rightarrow B$  and  $g : B \rightarrow C$ , we define the *composition of  $g$  with  $f$* , also called “ $g$  composed  $f$ ”, to be the map  $g \circ f : A \rightarrow C$  defined by

$$(g \circ f)(a) = g(f(a)), \quad \forall a \in A.$$

Prove the following.

- (i) If both  $f$  and  $g$  are injective, so is  $g \circ f : A \rightarrow C$ .
- (ii) If both  $f$  and  $g$  are surjective, so is  $g \circ f : A \rightarrow C$ .
- (iii) If  $g \circ f$  is injective, show that  $f$  is injective.
- (iv) Give an example to show that  $g \circ f$  could be injective without  $g$  being injective.
- (v) If  $g \circ f$  is surjective, show that  $g$  is surjective.
- (vi) Give an example to show that  $g \circ f$  could be surjective without  $f$  being surjective.
- (vii) Given any set  $A$ , there is a map  $\text{Id}_A : A \rightarrow A$  defined by  $\text{Id}_A(a) = a, \forall a \in A$ , known as the *identity map* of  $A$ . Verify that  $\text{Id}_A$  is bijective.
- (viii) If  $f : A \rightarrow B$  is bijective, show that there is a bijective map  $\tilde{f} : B \rightarrow A$  such that  $\tilde{f} \circ f = \text{Id}_A$  and  $f \circ \tilde{f} = \text{Id}_B$ . The bijective map  $\tilde{f} : B \rightarrow A$ , defined above, is called the *inverse of  $f$* , and is usually denoted by  $f^{-1}$ .

**Definition 2.1.14.** A *permutation* on a set  $A$  is a bijective map from  $A$  onto itself.

For a non-empty set  $A$ , we denote by  $S_A$  the set of all permutations on  $A$ . Let  $A$  be a non-empty set. Define a binary operation on  $S_A$  by

$$\circ : S_A \times S_A \longrightarrow S_A, \quad (f, g) \longmapsto g \circ f.$$

Verify that  $(S_A, \circ)$  is a group. (Hint: Use Exercise 2.1.13).

**Example 2.1.15** (Symmetric group  $S_3$ ). Consider an equilateral triangle  $\triangle$  in a plane with its vertices labelled as 1, 2 and 3. Consider the symmetries of  $\triangle$  obtained by its rotations by angles  $2n\pi/3$ , for  $n \in \mathbb{Z}$ , around its centre, and reflections along a straight line passing through its top vertex and centre. Note that, we have only six possible symmetries of  $\triangle$  as follow:

$$\begin{aligned} \sigma_0 &= \begin{pmatrix} 1 \mapsto 1 \\ 2 \mapsto 2 \\ 3 \mapsto 3 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 1 \mapsto 2 \\ 2 \mapsto 3 \\ 3 \mapsto 1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 1 \mapsto 3 \\ 2 \mapsto 1 \\ 3 \mapsto 2 \end{pmatrix}, \\ \sigma_3 &= \begin{pmatrix} 1 \mapsto 1 \\ 2 \mapsto 3 \\ 3 \mapsto 2 \end{pmatrix}, \quad \sigma_4 = \begin{pmatrix} 1 \mapsto 3 \\ 2 \mapsto 2 \\ 3 \mapsto 1 \end{pmatrix}, \quad \sigma_5 = \begin{pmatrix} 1 \mapsto 2 \\ 2 \mapsto 1 \\ 3 \mapsto 3 \end{pmatrix}. \end{aligned}$$

Let  $S_3 := \{\sigma_0, \sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5\}$ . Note that, each of symmetries are bijective maps from the set  $J_3 := \{1, 2, 3\}$  onto itself, and any bijective map from  $J_3$  onto itself is one of the symmetries in  $S_3$ . Since composition of bijective maps is bijective (see Exercise 2.1.13), we get a binary operation

$$S_3 \times S_3 \longrightarrow S_3, \quad (\sigma_i, \sigma_j) \longmapsto \sigma_i \circ \sigma_j.$$



**Exercise 2.1.16.** Write down the Cayley table for this binary operation on  $S_3$  defined by composition of maps, and show that  $S_3$  together with this binary operation is a group. Find  $\sigma_1, \sigma_2 \in S_3$  such that  $\sigma_1 \circ \sigma_2 \neq \sigma_2 \circ \sigma_1$ .

**Definition 2.1.17.** The *order* of a group  $G$  is the cardinality of its underlying set  $G$ . We denote this by  $|G|$ . In particular, if  $G$  is a finite set, then  $|G|$  is the number of elements of the set  $G$ .

**Example 2.1.18.** Let  $S_4$  be the set of all bijective maps from  $J_4 := \{1, 2, 3, 4\}$  onto itself. Given any two elements  $\sigma, \tau \in S_4$ , note that their composition  $\sigma \circ \tau \in S_4$ . Thus we have a binary operation on  $S_4$  given by sending  $(\sigma, \tau) \in S_4 \times S_4$  to  $\sigma \circ \tau \in S_4$ . Show that the set  $S_4$  together with this binary operation (composition of bijective maps) is a non-commutative group of order  $4! = 24$ .

**Definition 2.1.19.** Let  $A \subseteq \mathbb{R}$ . A map  $f : A \rightarrow \mathbb{R}$  is said to be *continuous* at  $a \in A$  if given any real number  $\epsilon > 0$ , there is a real number  $\delta > 0$  (depending on both  $\epsilon$  and  $a$ ) such that for each  $x \in A$  satisfying  $|a - x| < \delta$ , we have  $|f(a) - f(x)| < \epsilon$ . If  $f$  is continuous at each point of  $A$ , we say that  $f$  is *continuous on  $A$* .

**Exercise 2.1.20.** Let  $A \subseteq \mathbb{R}$ , and let  $C(A) := \{f : A \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ . Verify that  $C(A)$  is a group with respect to the binary operation defined for all  $f, g \in C(A)$  by the formula

$$(f + g)(x) := f(x) + g(x), \quad \forall x \in A.$$

*Solution.* Let  $f_1, f_2 \in C(A)$ . Let  $a \in A$  be arbitrary but fixed after choice. Since both  $f_1$  and  $f_2$  are continuous at  $a$ , given a real number  $\epsilon > 0$ , there exist real numbers  $\delta_1, \delta_2 > 0$  such that for each  $x \in A$  satisfying  $|a - x| < \delta_j$  we have  $|f_j(a) - f_j(x)| < \epsilon/2$ , for all  $j = 1, 2$ . Let  $\delta := \min\{\delta_1, \delta_2\}$ . Then  $\delta > 0$ , and for any  $x \in A$  satisfying  $|a - x| < \delta$ , we have  $|f_j(a) - f_j(x)| < \epsilon/2$ , for all  $j = 1, 2$ . Then we have,

$$\begin{aligned} |(f_1 + f_2)(a) - (f_1 + f_2)(x)| &= |f_1(a) + f_2(a) - f_1(x) - f_2(x)| \\ &= |(f_1(a) - f_1(x)) + (f_2(a) - f_2(x))| \\ &\leq |f_1(a) - f_1(x)| + |f_2(a) - f_2(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Therefore,  $f_1 + f_2$  is continuous at  $a \in A$ . Since  $a \in A$  is arbitrary,  $f_1 + f_2$  is continuous at every points of  $A$ , and hence  $f_1 + f_2 \in C(A)$ . Since for given  $f_1, f_2, f_3 \in C(A)$  and any  $x \in A$ , we have

$$\begin{aligned} ((f_1 + f_2) + f_3)(x) &= (f_1 + f_2)(x) + f_3(x) \\ &= (f_1(x) + f_2(x)) + f_3(x) \\ &= f_1(x) + (f_2(x) + f_3(x)) \\ &= f_1(x) + (f_2 + f_3)(x) \\ &= (f_1 + (f_2 + f_3))(x), \end{aligned}$$

we have  $(f_1 + f_2) + f_3 = f_1 + (f_2 + f_3)$ . Note that, the constant function

$$0 : A \rightarrow \mathbb{R}$$

defined by sending all points of  $A$  to  $0 \in \mathbb{R}$ , given by  $0(a) = 0, \quad \forall a \in A$ , is continuous (*Hint*: given  $\epsilon > 0$ , take any  $\delta > 0$ ), and satisfies  $f + 0 = f = 0 + f$ , for all  $f \in A$ . Given  $f \in C(A)$ , note that the function  $-f$  defined by  $(-f)(a) = -f(a)$ , for all  $a \in A$ , is continuous on  $A$  (*Hint*: given  $\epsilon > 0$ , take the same  $\delta > 0$  which works for  $f$ ), and satisfies  $f + (-f) = (-f) + f = 0$ . Therefore,  $(C(A), +)$  satisfies all axioms of a group, and hence is a group.  $\square$

**Example 2.1.21** (Matrix groups). (i) Fix two integers  $m, n \geq 1$ , and let  $M_{m \times n}(\mathbb{R})$  be the set of all  $m \times n$  matrices with entries from  $\mathbb{R}$ . Given  $A, B \in M_{m \times n}(\mathbb{R})$ , we define their *addition*

to be the matrix  $A + B \in M_{m \times n}(\mathbb{R})$  whose  $(i, j)$ -th entry is given by  $a_{ij} + b_{ij}$ , where  $a_{ij}$  and  $b_{ij}$  are the  $(i, j)$ -th entries of  $A$  and  $B$ , respectively. Then we have a binary operation

$$+ : M_{m \times n}(\mathbb{R}) \times M_{m \times n}(\mathbb{R}) \longrightarrow M_{m \times n}(\mathbb{R}), \quad (A, B) \longmapsto A + B.$$

Clearly, the set  $M_{m \times n}(\mathbb{R})$  is non-empty, and the pair  $(M_{m \times n}(\mathbb{R}), +)$  satisfies the properties (G1)–(G3) in Definition 2.1.4.

- (ii) **Matrix multiplication:** Fix positive integers  $m, n, p$ , and let  $A \in M_{m \times n}(\mathbb{R})$  and  $B \in M_{n \times p}(\mathbb{R})$ . Define the *product of  $A$  and  $B$*  to be the  $m \times p$  matrix  $AB \in M_{m \times p}(\mathbb{R})$ , whose  $(i, j)$ -th entry is

$$(2.1.22) \quad c_{ij} = \sum_{k=1}^n a_{ik} b_{kj},$$

where  $a_{ik}$  is the  $(i, k)$ -th entry of  $A$ , and  $b_{kj}$  is the  $(k, j)$ -th entry of  $B$ .

Let  $A \in M_{n \times n}(\mathbb{R})$ . A matrix  $B \in M_{n \times n}(\mathbb{R})$  is said to be the *left inverse* (resp., *right inverse*) of  $A$  if  $BA = I_n$  (resp.,  $AB = I_n$ ), where  $I_n \in M_{n \times n}(\mathbb{R})$  whose  $(i, j)$ -th entry is

$$\delta_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}$$

**Exercise 2.1.23.** Show that the left inverse and the right inverse of  $A \in M_{n \times n}(\mathbb{R})$ , when they exist, are the same. In other words, if  $AB = I_n$  and  $CA = I_n$ , for some  $B, C \in M_{n \times n}(\mathbb{R})$ , show that  $B = C$ .

A matrix  $A \in M_{n \times n}(\mathbb{R})$  is said to be *invertible* if there is a matrix  $B \in M_{n \times n}(\mathbb{R})$  such that  $AB = BA = I_n$ .

**General linear group:** Let

$$\text{GL}_n(\mathbb{R}) = \{A \in M_{n \times n}(\mathbb{R}) : A \text{ is invertible}\}$$

be the set of all invertible  $n \times n$  matrices with real entries.

- (a) Show that  $\text{GL}_n(\mathbb{R})$  is a group with respect to matrix multiplication.
- (b) Give examples of  $A, B \in \text{GL}_n(\mathbb{R})$  such that  $A + B \notin \text{GL}_n(\mathbb{R})$ .
- (c) Give an example of  $A \in M_{n \times n}(\mathbb{R})$  such that  $AB \neq I_n, \forall B \in M_{n \times n}(\mathbb{R})$ .
- (d) Assuming  $n \geq 2$  give examples of  $A, B \in \text{GL}_n(\mathbb{R})$  such that  $AB \neq BA$ .

The group  $\text{GL}_n(\mathbb{R})$  is called the *general linear group* (of degree  $n$ ).

As we see in Example 2.1.21 that the relation  $ab = ba$  need not hold for all  $a, b \in G$ , in general. We shall see later that the symmetric group  $S_3$  in Example 2.1.9 (2.1.15) is the smallest such group; in this case, we have  $\sigma_3 \circ \sigma_1 = \sigma_4$  while  $\sigma_1 \circ \sigma_3 = \sigma_5$ .

**Definition 2.1.24.** A group  $G$  is said to be *commutative* (or, *abelian*) if  $ab = ba$ , for all  $a, b \in G$ . A group which is not commutative (or, abelian) is called a *non-commutative* (or, *non-abelian*) group.

**Exercise 2.1.25.** (i) Verify that  $\{e\}, \mathbb{Z}, \mathbb{C}^*, S^1, K_4$  are abelian groups.

- (ii) Show that  $S_3$  and  $\text{GL}_2(\mathbb{R})$  are non-abelian groups.

**Exercise 2.1.26.** Show that  $\text{GL}_n(\mathbb{R})$  is not abelian, for all  $n \geq 2$ .

**Definition 2.1.27.** A *relation* on a non-empty set  $A$  is a non-empty subset  $\rho \subseteq A \times A$ . If  $(a, b) \in \rho$ , sometimes we may express it as  $a \rho b$ , and call  $a$  is  $\rho$ -related to  $b$  in  $A$ . A relation  $\rho$  on  $A$  is said to be

- (i) *reflexive* if  $(a, a) \in \rho, \forall a \in A$ ;
- (ii) *symmetric* if  $(a, b) \in \rho$  implies  $(b, a) \in \rho$ ;
- (iii) *anti-symmetric* if  $(a, b) \in \rho$  and  $(b, a) \in \rho$  implies  $a = b$ ;
- (iv) *transitive* if  $(a, b) \in \rho$  and  $(b, c) \in \rho$  implies  $(a, c) \in \rho$ ;
- (v) *equivalence* if  $\rho$  is reflexive, symmetric and transitive; and
- (vi) *partial order* if  $\rho$  is reflexive, anti-symmetric and transitive.

Let  $A$  be a non-empty set, and let  $\rho$  be an equivalence relation on  $A$ . The  $\rho$ -equivalence class of an element  $a \in A$  is the subset

$$[a]_\rho := \{b \in A : (b, a) \in \rho\} \subseteq A.$$

**Proposition 2.1.28.** *With the above notations, given any  $a, b \in A$ ,  $[a]_\rho = [b]_\rho$  if and only if  $(a, b) \in \rho$ .*

*Proof.* Suppose that  $(a, b) \in \rho$ . Then for any  $c \in [a]_\rho$ , we have  $(c, a) \in \rho$ . Since  $\rho$  is transitive, from  $(c, a), (a, b) \in \rho$  we have  $(c, b) \in \rho$ , and so  $c \in [b]_\rho$ . Therefore,  $[a]_\rho \subseteq [b]_\rho$ . Since  $\rho$  is symmetric,  $(a, b) \in \rho$  implies  $(b, a) \in \rho$ . Then following above arguments, we conclude that  $[b] \subseteq [a]$ . Therefore,  $[a]_\rho = [b]_\rho$ .

Conversely, suppose that  $[a]_\rho = [b]_\rho$ . Since  $\rho$  is reflexive,  $a \in [a]_\rho$ . Then  $[a]_\rho = [b]_\rho$  implies that  $a \in [b]_\rho$ , and so  $(a, b) \in \rho$ . This completes the proof.  $\square$

**Proposition 2.1.29.** *With the above notations, given  $a, b \in A$ , either  $[a]_\rho \cap [b]_\rho = \emptyset$  or  $[a]_\rho = [b]_\rho$ .*

*Proof.* It is enough to show that if  $[a]_\rho \cap [b]_\rho \neq \emptyset$ , then  $[a]_\rho = [b]_\rho$ . Let  $c \in [a]_\rho \cap [b]_\rho$ . Then  $(c, a), (c, b) \in \rho$ . Since  $\rho$  is symmetric,  $(c, a) \in \rho$  implies  $(a, c) \in \rho$ . Then  $(a, c) \in \rho$  and  $(c, b) \in \rho$  together implies  $(a, b) \in \rho$ , since  $\rho$  is transitive. Then by Proposition 2.1.28 we have  $[a]_\rho = [b]_\rho$ .  $\square$

**Definition 2.1.30.** Let  $A$  be a non-empty set. A *partition* on  $A$  is a non-empty collection  $\mathcal{P} := \{A_\alpha : \alpha \in \Lambda\}$ , where

- (i)  $A_\alpha \subseteq A$ , for all  $\alpha \in \Lambda$ ,
- (ii)  $A_\alpha \cap A_\beta = \emptyset$ , for  $\alpha \neq \beta$  in  $\Lambda$ , and
- (iii)  $A = \bigcup_{\alpha \in \Lambda} A_\alpha$ .

**Proposition 2.1.31.** *To give an equivalence relation on a non-empty set is equivalent to give a partition on it.*

*Proof.* Suppose that we have given an equivalence relation  $\rho$  on  $A$ . Since  $\rho$  is reflexive,  $a \in [a]_\rho$ , for all  $a \in A$ , and hence  $A = \bigcup_{a \in A} [a]_\rho$ . Since  $\rho$ -equivalence classes of elements of  $A$  are either disjoint or equal (see Proposition 2.1.29), the collection  $\mathcal{P}$  consisting of all distinct  $\rho$ -equivalence classes of elements of  $A$  is a partition of  $A$ .

Conversely, suppose that  $\mathcal{P} = \{A_\alpha : \alpha \in \Lambda\}$  be a partition of  $A$ . Define

$$\rho = \{(a, b) \in A \times A : a, b \in A_\alpha, \text{ for some } \alpha \in \Lambda\}.$$

Note that  $(a, a) \in \rho$ , for all  $a \in A$ . If  $(a, b) \in \rho$ , then both  $a$  and  $b$  are in the same  $A_\alpha$ , for some  $\alpha \in \Lambda$ , and so  $(b, a) \in \rho$ . So  $\rho$  is symmetric. If  $(a, b), (b, c) \in \rho$ , then  $a, b \in A_\alpha$  and  $b, c \in A_\beta$ , for some  $\alpha, \beta \in \Lambda$ . Since  $b \in A_\alpha \cap A_\beta$ , so we must have  $A_\alpha = A_\beta$ . Therefore,  $(a, c) \in \rho$ . Thus  $\rho$  is transitive. Therefore,  $\rho$  is an equivalence relation on  $A$ . One should note that the elements of  $\mathcal{P}$  are precisely the  $\rho$ -equivalence classes in  $A$  (verify!).  $\square$

**Example 2.1.32** (The groups  $\mathbb{Z}_n$  and  $U_n$ ). Fix an integer  $n \geq 2$ . Define a relation  $\equiv_n$  on  $\mathbb{Z}$  by setting

$$a \equiv_n b, \text{ if } a - b = nk, \text{ for some } k \in \mathbb{Z}.$$

If  $a \equiv_n b$  sometimes we also express it as  $a \equiv b \pmod{n}$ , and say that  $a$  is congruent to  $b$  modulo  $n$ . Verify that  $\equiv_n$  is an equivalence relation on  $\mathbb{Z}$ . Given any  $a \in \mathbb{Z}$ , let

$$[a] := \{b \in \mathbb{Z} : b \equiv_n a\} \subseteq \mathbb{Z}$$

be the  $\equiv_n$ -equivalence class of  $a$  in  $\mathbb{Z}$ . Let

$$\mathbb{Z}_n := \{[a] : a \in \mathbb{Z}\}$$

be the set of all  $\equiv_n$ -equivalence classes of elements of  $\mathbb{Z}$ . Let  $a, b \in \mathbb{Z}$ . If  $c \in [a] \cap [b]$ , then  $c = a + nk_1$  and  $c = b + nk_2$ , for some  $k_1, k_2 \in \mathbb{Z}$ . Then  $a - b = n(k_1 - k_2)$ , and hence  $a \equiv_n b$ . Then  $[a] = [b]$  in  $\mathbb{Z}_n$ . Therefore, the  $\equiv_n$ -equivalence classes are either disjoint or identical (c.f. Proposition 2.1.29). Use division algorithm (Theorem 1.3.1) to show that  $\equiv_n$ -equivalence classes  $[0], [1], \dots, [n-1]$  are all distinct, and

$$\mathbb{Z}_n = \{[k] : 0 \leq k \leq n-1\}.$$

In particular,  $\mathbb{Z}_n$  is a finite set containing  $n$  elements.

We now define two binary operations on  $\mathbb{Z}_n$ . Suppose that  $[a] = [a']$  and  $[b] = [b']$  in  $\mathbb{Z}_n$ , for some  $a, a', b, b' \in \mathbb{Z}$ . Then we have

$$\begin{aligned} a - a' &= nk_1, \\ \text{and } b - b' &= nk_2, \end{aligned}$$

for some  $k_1, k_2 \in \mathbb{Z}$ . Therefore,

$$(a + b) - (a' + b') = n(k_1 + k_2),$$

and hence  $[a + b] = [a' + b']$  in  $\mathbb{Z}_n$ . Therefore, we have a well-defined binary operation on  $\mathbb{Z}_n$  (called *addition of integers modulo  $n$* ) given by

$$[a] + [b] := [a + b], \quad \forall [a], [b] \in \mathbb{Z}_n.$$

Now it is easy to see that,

- (i)  $([a] + [b]) + [c] = [a] + ([b] + [c])$ , for all  $[a], [b], [c] \in \mathbb{Z}_n$ .
- (ii)  $[a] + [0] = [a] = [0] + [a]$ , for all  $[a] \in \mathbb{Z}_n$ .
- (iii)  $[a] + [-a] = [0]$ , for all  $[a] \in \mathbb{Z}_n$ .

Therefore,  $(\mathbb{Z}_n, +)$  is a group. Note that, for all  $[a], [b] \in \mathbb{Z}_n$  we have

$$\begin{aligned} [a] + [b] &= [a + b] = [b + a], \text{ since addition in } \mathbb{Z} \text{ is commutative,} \\ &= [b] + [a]. \end{aligned}$$

Therefore,  $(\mathbb{Z}_n, +)$  is an abelian group.

Now we define *multiplication operation on  $\mathbb{Z}_n$* . Suppose that  $[a] = [a']$  and  $[b] = [b']$ . Then  $a - a' = nk_1$  and  $b - b' = nk_2$ , for some  $k_1, k_2 \in \mathbb{Z}$ . Then

$$\begin{aligned} ab - a'b' &= (a - a')b + a'(b - b') \\ &= nk_1b + a'nk_2 \\ &= n(k_1b + a'k_2), \end{aligned}$$

implies that  $[ab] = [a'b']$ . Thus we have a well-defined binary operations on  $\mathbb{Z}_n$  (called the *multiplication of integers modulo  $n$* ) defined by

$$[a] \cdot [b] := [ab], \quad \forall [a], [b] \in \mathbb{Z}_n.$$

Clearly the multiplication modulo  $n$  operation on  $\mathbb{Z}_n$  is both associative and commutative. Note that,

$$[1] \cdot [a] = [a] = [a] \cdot [1], \quad \forall [a] \in \mathbb{Z}_n.$$

Therefore,  $[1] \in \mathbb{Z}_n$  is the multiplicative identity in  $\mathbb{Z}_n$ . Moreover, the multiplication distributes over addition from left and right on  $\mathbb{Z}_n$ . Indeed, we have

$$\begin{aligned} [a] \cdot ([b] + [c]) &= [a] \cdot [b] + [a] \cdot [c], \\ \text{and } ([a] + [b]) \cdot [c] &= [a] \cdot [c] + [b] \cdot [c]. \end{aligned}$$

Such a triple  $(\mathbb{Z}_n, +, \cdot)$  is called a *ring*. Since  $n \geq 2$  by assumption,  $n$  does not divide 1 in  $\mathbb{Z}_n$ . So  $[0] \neq [1]$  in  $\mathbb{Z}_n$  by Proposition 2.1.28. Since for any  $[a] \in \mathbb{Z}_n$ , we have  $[0] \cdot [a] = [0 \cdot a] = [0] \neq [1]$ , we see that  $[0] \in \mathbb{Z}_n$  has no multiplicative inverse in  $\mathbb{Z}_n$ . Therefore,  $(\mathbb{Z}_n, \cdot)$  is just a commutative monoid, but not a group.

We now find out elements of  $\mathbb{Z}_n$  that have multiplicative inverse in  $\mathbb{Z}_n$ , and use them to construct a subset of  $\mathbb{Z}_n$  which forms a group with respect to the multiplication modulo  $n$  operation. Recall that given  $n, k \in \mathbb{Z}$ , we have  $\gcd(n, k) = 1$  if and only if there exists  $a, b \in \mathbb{Z}$  such that  $an + bk = 1$  (see Corollary 1.3.8). Use this to verify that if  $[k] = [k']$  in  $\mathbb{Z}_n$ , then  $\gcd(n, k) = 1$  if and only if  $\gcd(n, k') = 1$ . Thus we get a well-defined subset

$$U_n := \{[k] \in \mathbb{Z}_n : \gcd(k, n) = 1\} \subset \mathbb{Z}_n.$$

Note that,  $[0] \notin U_n$ . If  $[k_1], [k_2] \in U_n$ , then  $\gcd(k_1, n) = 1 = \gcd(k_2, n)$ . Then there exists  $a_1, b_1, a_2, b_2 \in \mathbb{Z}$  such that

$$\begin{aligned} a_1 k_1 + b_1 n &= 1 \\ \text{and } a_2 k_2 + b_2 n &= 1. \end{aligned}$$

Multiplying these two equations, we have

$$(a_1 a_2)(k_1 k_2) + (a_1 k_1 b_2 + a_2 k_2 b_1 + b_1 b_2)n = 1.$$

Then we have  $\gcd(k_1 k_2, n) = 1$ . Therefore,

$$[k_1] \cdot [k_2] = [k_1 k_2] \in U_n, \quad \forall [k_1], [k_2] \in U_n.$$

Verify that  $(U_n, \cdot)$  is an abelian group. If  $p > 1$  is a prime number (see Definition 1.3.10), show that  $U_p = \mathbb{Z}_p \setminus \{[0]\}$ , as sets.

**Exercise 2.1.33.** Let  $X$  be a non-empty set. Let  $\mathcal{P}(X)$  be the set of all subsets of  $X$ ; called the *power set of  $X$* . Given any two elements  $A, B \in \mathcal{P}(X)$ , define

$$A \triangle B := (A \setminus B) \cup (B \setminus A).$$

The set  $A \triangle B$  is known as the *symmetric difference* of  $A$  and  $B$ . Show that  $(\mathcal{P}(X), \triangle)$  is a commutative group. (*Hint:* The empty subset  $\emptyset \subset X$  acts as the neutral element in  $\mathcal{P}(X)$ , and every element of  $\mathcal{P}(X)$  is inverse of itself).

**Exercise 2.1.34** (Direct product of two groups). Let  $(A, *)$  and  $(B, \star)$  be two groups. Show that the Cartesian product  $G_1 \times G_2$  is a group with respect to the binary operation on it defined by

$$(a_1, b_1)(a_2, b_2) := (a_1 * a_2, b_1 \star b_2), \quad \forall (a_1, b_1), (a_2, b_2) \in A \times B.$$

The group  $A \times B$  defined above is called the *direct product* of  $A$  with  $B$ .

## 2.2 Subgroup

**Definition 2.2.1** (Subgroup). Let  $G$  be a group. A *subgroup* of  $G$  is a subset  $H \subseteq G$  such that  $H$  is a group with respect to the binary operation induced from  $G$ . A subgroup  $H$  of  $G$  is said to be *proper* if  $H \neq G$ . A subgroup whose underlying set is singleton is called a *trivial* subgroup. If  $H$  is a subgroup of  $G$ , we express it symbolically by  $H \leq G$ .

For example,  $\mathbb{Z}$  is a subgroup of  $\mathbb{Q}$ ;  $S^1$  is a subgroup of  $\mathbb{C}^*$  etc.

**Exercise 2.2.2.** For each integer  $n$ , let  $n\mathbb{Z} := \{nk : k \in \mathbb{Z}\}$ .

- (i) Show that  $n\mathbb{Z}$  is a proper subgroup of  $\mathbb{Z}$ , for all  $n \in \mathbb{Z} \setminus \{1, -1\}$ .
- (ii) Show that any subgroup of  $\mathbb{Z}$  is of the form  $n\mathbb{Z}$ , for some  $n \in \mathbb{Z}$ .

**Exercise 2.2.3** (Group of  $n^{\text{th}}$  roots of unity). Fix an integer  $n \geq 1$ , and let

$$\mu_n := \{\zeta \in \mathbb{C} \mid \zeta^n = 1\}.$$

Show that  $\mu_n$  is a subgroup of the circle group  $S^1$ .

**Exercise 2.2.4.** Show that a finite subgroup of  $\mathbb{C}^*$  of order  $n$  is  $\mu_n$ .

**Exercise 2.2.5.** Show that  $\{1, -1, i, -i\}$  is a subgroup of  $\mathbb{C}^*$ , where  $i = \sqrt{-1}$ .

**Exercise 2.2.6.** For each integer  $n \geq 1$ , show that there is a commutative group of order  $n$ .

**Remark 2.2.7.** It is easy to see that any subgroup of an abelian group is abelian. However, the converse is not true, in general. For example, one can easily check that  $S_3$  is a non-abelian group whose all proper subgroups are abelian.

**Lemma 2.2.8.** Let  $G$  be a group. A non-empty subset  $H \subseteq G$  forms a subgroup of  $G$  if and only if  $ab^{-1} \in H$ , for all  $a, b \in H$ .

*Proof.* Since  $H \neq \emptyset$ , there is an element  $a \in H$ . Then  $e = aa^{-1} \in H$ . In particular, for any  $b \in H$ , its inverse  $b^{-1} = eb^{-1} \in H$ . Then for any  $a, b \in H$ , their product  $ab = a(b^{-1})^{-1} \in H$ . Thus  $H$  is closed under the binary operation induced from  $G$ . Associativity is obvious. Thus,  $H$  is a subgroup of  $G$ .  $\square$

**Exercise 2.2.9.** Let  $G$  be a group. Show that a non-empty subset  $H \subseteq G$  forms a subgroup of  $G$  if and only if  $a^{-1}b \in H$ , for all  $a, b \in H$ .

**Exercise 2.2.10.** Let  $G$  be a group. Let  $H$  be a finite non-empty subset of  $G$ . Show that  $H$  forms a subgroup of  $G$  if and only if  $ab \in H$ , for all  $a, b \in H$ . Show by an example that this fails if  $H$  is infinite.

**Exercise 2.2.11** (Special linear group). Fix an integer  $n \geq 1$ , and let

$$\text{SL}_n(\mathbb{R}) = \{A \in \text{GL}_n(\mathbb{R}) : \det(A) = 1\},$$

where  $\det(A)$  denotes the determinant of the matrix  $A$ . Show that  $\text{SL}_n(\mathbb{R})$  is a non-trivial proper subgroup of  $\text{GL}_n(\mathbb{R})$ . Also show that  $\text{SL}_n(\mathbb{R})$  is non-commutative for  $n \geq 2$ .

**Exercise 2.2.12** (Orthogonal and special orthogonal groups). Fix an integer  $n \geq 1$ , and let

$$O_n(\mathbb{R}) := \{A \in M_{n \times n}(\mathbb{R}) : A^t A = I_n = {}^t A A\},$$

where  ${}^t A$  denotes the *transpose* of  $A$  (i.e., the  $n \times n$  matrix whose  $(i, j)$ -th entry is equal to the  $(j, i)$ -th entry of  $A$ , for all  $i, j \in \{1, \dots, n\}$ ).

- (i) Show that  $O_n(\mathbb{R})$  is a subgroup of  $GL_n(\mathbb{R})$ .
- (ii) Show that  $SO_n(\mathbb{R}) := \{A \in O_n(\mathbb{R}) : \det(A) = 1\}$  is a subgroup of both  $O_n(\mathbb{R})$  and  $SL_n(\mathbb{R})$ .

The groups  $O_n(\mathbb{R})$  and  $SO_n(\mathbb{R})$  are called the *orthogonal group* and the *special orthogonal group* over  $\mathbb{R}$ , respectively.

**Exercise 2.2.13** (Unitary and special unitary groups). Fix an integer  $n \geq 1$ , and let

$$U_n(\mathbb{C}) := \{A \in M_{n \times n}(\mathbb{C}) : AA^* = I_n = A^*A\},$$

where  $A^* = \overline{A}^t$  is the  $n \times n$  matrix over  $\mathbb{C}$  whose  $(i, j)$ -th entry is equal to the complex conjugate of the  $(j, i)$ -th entry of  $A$ , for all  $i, j \in \{1, \dots, n\}$ .

- (i) Show that  $U_n(\mathbb{C})$  is a subgroup of  $GL_n(\mathbb{C})$ .
- (ii) Show that  $U_1(\mathbb{C}) = S^1$ .
- (iii) Show that  $SU_n(\mathbb{C}) := \{A \in U_n(\mathbb{C}) : \det(A) = 1\}$  is a subgroup of both  $U_n(\mathbb{C})$  and  $SL_n(\mathbb{C})$ .

The groups  $U_n(\mathbb{C})$  and  $SU_n(\mathbb{C})$  are called the *unitary group* and the *special unitary group* over  $\mathbb{C}$ , respectively.

**Proposition 2.2.14** (Center of a group). *Let  $G$  be a group. Then*

$$Z(G) := \{a \in G : ab = ba, \forall b \in G\}$$

*is a commutative subgroup of  $G$ , called the center of  $G$ .*

*Proof.* Clearly  $e \in Z(G)$ . Let  $a \in Z(G)$ . Then for any  $c \in G$  we have

$$ac = ca \Rightarrow c = a^{-1}ca \Rightarrow ca^{-1} = a^{-1}caa^{-1} = a^{-1}c,$$

and hence  $a^{-1} \in Z(G)$ . Then for any  $a, b \in Z(G)$ , we have  $c(ab^{-1})c^{-1} = cac^{-1}cb^{-1}c^{-1} = ab^{-1}$ , for all  $c \in G$ , and hence  $ab^{-1} \in Z(G)$ . Therefore,  $Z(G)$  is a subgroup of  $G$ . Clearly  $Z(G)$  is commutative.  $\square$

**Exercise 2.2.15.** Show that a group  $G$  is commutative if and only if  $Z(G) = G$ .

**Exercise 2.2.16.** Find the centers of  $S_3$ ,  $GL_n(\mathbb{R})$  and  $SL_n(\mathbb{R})$ , where  $n \in \mathbb{N}$ .

**Exercise 2.2.17** (Centralizer). Let  $G$  be a group. Given an element  $a \in G$  show that the subset

$$C_G(a) := \{b \in G : ab = ba\}$$

is a subgroup of  $G$ , called the *centralizer of  $a$  in  $G$* . Show that  $Z(G) = \bigcap_{a \in G} C_G(a)$ .

**Lemma 2.2.18.** *Let  $G$  be a group, and let  $\{H_\alpha\}_{\alpha \in \Lambda}$  be a non-empty collection of subgroups of  $G$ . Then  $\bigcap_{\alpha \in \Lambda} H_\alpha$  is a subgroup of  $G$ .*

*Proof.* Since  $e \in H_\alpha$ , for all  $\alpha \in \Lambda$ , we have  $e \in \bigcap_{\alpha \in \Lambda} H_\alpha$ . Let  $a, b \in \bigcap_{\alpha \in \Lambda} H_\alpha$  be arbitrary. Since  $a, b \in H_\alpha$ , for all  $\alpha \in \Lambda$ , we have  $ab^{-1} \in H_\alpha$ , for all  $\alpha \in \Lambda$ , and hence  $ab^{-1} \in \bigcap_{\alpha \in \Lambda} H_\alpha$ . Thus  $\bigcap_{\alpha \in \Lambda} H_\alpha$  is a subgroup of  $G$ .  $\square$

**Corollary 2.2.19.** *Let  $G$  be a group and  $S$  a subset of  $G$ . Let  $\mathcal{C}_S$  be the collection of all subgroups of  $G$  that contains  $S$ . Then  $\langle S \rangle := \bigcap_{H \in \mathcal{C}_S} H$  is the smallest subgroup of  $G$  containing  $S$ .*

*Proof.* By Lemma 2.2.18,  $\langle S \rangle := \bigcap_{H \in \mathcal{C}_S} H$  is a subgroup of  $G$  containing  $S$ . If  $H'$  is any subgroup of  $G$  containing  $S$ , then  $H' \in \mathcal{C}_S$ , and hence  $\langle S \rangle := \bigcap_{H \in \mathcal{C}_S} H \subseteq H'$ .  $\square$

**Exercise 2.2.20.** Recall Exercise 2.2.2, and find the subgroup  $2\mathbb{Z} \cap 3\mathbb{Z}$  of  $\mathbb{Z}$ .

**Exercise 2.2.21.** Is  $2\mathbb{Z} \cup 3\mathbb{Z}$  a subgroup of  $\mathbb{Z}$ ? Justify your answer.

**Exercise 2.2.22.** Show that a group cannot be written as a union of its two proper subgroups.

**Definition 2.2.23.** Let  $G$  be a group and  $S \subseteq G$ . The group  $\langle S \rangle := \bigcap_{H \in \mathcal{C}_S} H$  is called the *subgroup of  $G$  generated by  $S$* . If  $S$  is a singleton subset  $S = \{a\}$  of  $G$ , we denote by  $\langle a \rangle$ .

**Exercise 2.2.24.** Let  $G$  be a group. Find the subgroup of  $G$  generated by the empty subset of  $G$ .

**Proposition 2.2.25.** Let  $G$  be a group, and let  $S$  be a non-empty subset of  $G$ . Then

$$\langle S \rangle = \{a_1^{e_1} \cdots a_n^{e_n} \mid n \in \mathbb{N}, \text{ and } a_i \in S, e_i \in \{1, -1\}, \forall i \in \{1, 2, \dots, n\}\}.$$

*Proof.* Let

$$K := \{a_1^{e_1} \cdots a_n^{e_n} \mid n \in \mathbb{N}, \text{ and } a_i \in S, e_i \in \{1, -1\}, \forall i \in \{1, 2, \dots, n\}\}.$$

Clearly  $S \subset K \subseteq G$ . Taking  $n = 2$ ,  $a_1 = a_2 = a \in S$ ,  $e_1 = 1$  and  $e_2 = -1$ , we have  $e = a a^{-1} \in K$ . Let  $a, b \in K$ . Then  $a = a_1^{e_1} \cdots a_n^{e_n}$  and  $b = b_1^{f_1} \cdots b_m^{f_m}$ , for some  $a_i, b_j \in S$ ,  $e_i, f_j \in \{1, -1\}$ ,  $1 \leq i \leq n$ ,  $1 \leq j \leq m$ , and  $m, n \in \mathbb{N}$ . Then  $ab^{-1} = a_1^{e_1} \cdots a_n^{e_n} \cdot (b_1^{f_1} \cdots b_m^{f_m})^{-1} = a_1^{e_1} \cdots a_n^{e_n} \cdot b_m^{-f_m} \cdots b_1^{-f_1} \in K$ . Therefore,  $K$  is a subgroup of  $G$  containing  $S$ . Then by Proposition 2.2.19, we have  $\langle S \rangle \subseteq K$ . To see the reverse inclusion, note that if  $S \subseteq H$ , for some subgroup  $H$  of  $G$ , then all the elements of  $K$  lies inside  $H$ . Therefore,  $K \subseteq \bigcap_{H \in \mathcal{C}_S} H = \langle S \rangle$ .  $\square$

**Definition 2.2.26.** A group  $G$  is said to be *finitely generated* if there exists a finite subset  $S \subseteq G$  such that the subgroup generated by  $S$  is equal to  $G$ , i.e.,  $\langle G \rangle = G$ .

**Example 2.2.27.** (i) Any finite group is finitely generated.

(ii) The additive group  $(\mathbb{Z}, +)$  is finitely generated.

**Exercise 2.2.28.** Let  $G$  and  $H$  be finitely generated groups. Verify if the direct product  $G \times H$  of  $G$  and  $H$ , as defined in Exercise 2.1.34, is finitely generated.

**Example 2.2.29.** Let  $G$  be a group. Given an element  $a \in G$ , the subgroup of  $G$  generated by  $a$  can be written as

$$\langle a \rangle = \{a^n : n \in \mathbb{Z}\};$$

and is called the *cyclic subgroup* of  $G$  generated by  $a$ .

**Definition 2.2.30.** Let  $G$  be a group. The *order* of an element  $a \in G$  is the smallest positive integer  $n$ , if exists, such that  $a^n = e$ . If no such positive integer  $n$  exists, we say that the order of  $a$  is infinite. We denote by  $\text{ord}(a)$  the order of  $a \in G$ . In other words, if we set  $S_a := \{n \in \mathbb{Z} : n \geq 1 \text{ and } a^n = e\}$ , then

$$\text{ord}(a) := \begin{cases} \inf S_a, & \text{if } S_a \neq \emptyset, \text{ and} \\ \infty, & \text{if } S_a = \emptyset. \end{cases}$$

**Exercise 2.2.31.** Let  $G$  be a group and  $a, b \in G$  be such that  $ab = ba$ . Show that  $(ab)^n = a^n b^n$ , for all  $n \in \mathbb{N}$ .

**Exercise 2.2.32.** Let  $G$  be a group. Let  $a, b \in G$  be elements of finite orders.

(i) If  $a^m = e$ , for some  $m \in \mathbb{N}$ , then show that  $\text{ord}(a) \mid m$ .



- (ii) Show that  $\text{ord}(a^n) = \frac{\text{ord}(a)}{\gcd(n, \text{ord}(a))}$ , for all  $n \in \mathbb{N}$ .
- (iii) Show that both  $a$  and  $a^{-1}$  have the same order in  $G$ .
- (iv) Show that both  $ab$  and  $ba$  have the same finite order in  $G$ .

**Exercise 2.2.33.** Let  $G$  be a group, and let  $a$  and  $b$  two elements of  $G$  of finite orders with  $ab = ba$ .

- (i) Show that  $\text{ord}(ab)$  divides  $\text{lcm}(\text{ord}(a), \text{ord}(b))$ .
- (ii) If  $\gcd(\text{ord}(a), \text{ord}(b)) = 1$ , show that  $\text{ord}(ab) = \text{ord}(a) \text{ord}(b)$ .

**Remark 2.2.34.** If we remove the assumption that  $ab = ba$  from the above Exercise 2.2.33 we can say absolutely nothing about the order of the product  $ab$ . In fact, given any integers  $m, n, r > 1$ , there exists a finite group  $G$  with elements  $a, b \in G$  such that  $\text{ord}(a) = m$ ,  $\text{ord}(b) = n$  and  $\text{ord}(ab) = r$ . The proof of this surprising fact requires some advanced techniques, and may appear at the end of this course.

**Exercise 2.2.35.** Consider the matrices

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & 2 \\ 1/2 & 0 \end{pmatrix}$$

in  $\text{GL}_2(\mathbb{R})$ . Show that  $\text{ord}(A) = \text{ord}(B) = 2$  while  $\text{ord}(AB) = \infty$ . Consequently, the subgroup  $\langle A, B \rangle \leq \text{GL}_2(\mathbb{R})$  generated by two order 2 elements of  $\text{GL}_2(\mathbb{R})$  is infinite.

**Exercise 2.2.36.** Let  $G$  be an abelian group. Let  $H := \{a \in G : \text{ord}(a) \text{ is finite}\}$ . Show that  $H$  is a subgroup of  $G$ .

**Exercise 2.2.37.** Show that any finite group of even order contains an element of order 2.

**Exercise 2.2.38.** Let  $G$  be a group such that any non-identity element of  $G$  has order 2. Show that  $G$  is abelian.

**Exercise 2.2.39.** Find two elements  $\sigma$  and  $\tau$  of  $S_3$  such that  $\langle \sigma, \tau \rangle = S_3$ .

**Exercise 2.2.40 (Derived subgroup).** Let  $G$  be a group. The *commutator* of two elements  $a, b \in G$  is the element  $[a, b] := aba^{-1}b^{-1} \in G$ . Given  $a, b \in G$ , show that

- (i)  $[a, b] = e$  if and only if  $ab = ba$ ;
- (ii)  $[a, b]^{-1} = [b, a]$ ; and
- (iii)  $g[a, b]g^{-1} = [gag^{-1}, gbg^{-1}]$ , for all  $g \in G$ .

The subgroup  $[G, G] := \langle [a, b] : a, b \in G \rangle$  of  $G$  generated by all commutators of elements of  $G$  is called the *derived subgroup* or the *commutator subgroup* of  $G$ . Show that  $[G, G]$  is a trivial subgroup of  $G$  if and only if  $G$  is abelian.

## 2.3 Cyclic group

Let  $G$  be a group. For any element  $a \in G$ , we consider the subset

$$\langle a \rangle := \{a^n : n \in \mathbb{Z}\} \subseteq G.$$

Clearly  $e \in \langle a \rangle$ , and for any two elements  $a^n, a^m \in \langle a \rangle$ , we have  $a^n \cdot (a^m)^{-1} = a^{n-m} \in \langle a \rangle$ . Therefore,  $\langle a \rangle$  is a subgroup of  $G$ , called the *cyclic subgroup* of  $G$  generated by  $a$ . If  $H$  is any subgroup of  $G$  with  $a \in H$ , then  $a^{-1} \in H$ , and hence  $a^n \in H$ , for all  $n \in \mathbb{Z}$ . Therefore,  $\langle a \rangle \subseteq H$ . Therefore,  $\langle a \rangle$  is the smallest subgroup of  $G$  containing  $a$ .

**Definition 2.3.1.** A group  $G$  is said to be *cyclic* if there is an element  $a \in G$  such that  $G = \langle a \rangle$ . The element  $a$  is called the *generator* of  $\langle a \rangle$ .

**Remark 2.3.2.** If  $G$  is a cyclic group generated by  $a \in G$ , then  $\langle a^{-1} \rangle = G$ . Therefore, if  $a^2 \neq e$ , the cyclic group  $\langle a \rangle$  has at least two distinct generators, namely  $a$  and  $a^{-1}$ . We shall see later that if a cyclic group  $\langle a \rangle$  has at least two distinct generators, then we must have  $a^2 \neq e$ .

For example, the additive group  $\mathbb{Z}$  is a cyclic group generated by 1 or  $-1$ . It is clear that a cyclic group may have more than one generators. For example,  $\mathbb{Z}_3$  is a cyclic group that can be generated by  $[1]$  or  $[2]$ .

**Example 2.3.3.**  $\mathbb{Z}_n$  is a finite cyclic group generated by  $[1] \in \mathbb{Z}_n$ . To see this, note that for any  $[m] \in \mathbb{Z}_n$ , we have  $[m] = [m \cdot 1] = m[1] \in \langle [1] \rangle \subseteq \mathbb{Z}_n$ . Therefore,  $\mathbb{Z}_n \subseteq \langle [1] \rangle$ , and hence  $\mathbb{Z}_n = \langle [1] \rangle$ .

**Proposition 2.3.4.** Fix an integer  $n \geq 2$ . Then  $[a] \in \mathbb{Z}_n$  is a generator of the group  $\mathbb{Z}_n$  if and only if  $\gcd(a, n) = 1$ .

*Proof.* Suppose that  $\langle [a] \rangle = \mathbb{Z}_n$ . Then there exists  $m \in \mathbb{Z}$  such that  $[1] = m[a] = [ma]$ . Then  $n \mid (ma - 1)$  and so  $ma - 1 = nd$ , for some  $d \in \mathbb{Z}$ . Therefore,  $ma + n(-d) = 1$ , and hence by Corollary 1.3.8 we have  $\gcd(a, n) = 1$ . Conversely, if  $\gcd(a, n) = 1$ , then there exists  $m, q \in \mathbb{Z}$  such that  $am + nq = 1$ . Then  $n \mid (1 - am)$  and hence  $[a] = [1]$  in  $\mathbb{Z}_n$ . Hence the result follows.  $\square$

**Corollary 2.3.5.** For a prime number  $p > 0$ ,  $\mathbb{Z}_p$  has  $p - 1$  distinct generators.

Clearly any cyclic group is abelian. However, the converse is not true in general. For example, the Klein four-group  $K_4$  in Example 2.1.9 (iii) is abelian but not cyclic (verify).

**Exercise 2.3.6.** Give an example of an infinite abelian group which is not cyclic.

**Proposition 2.3.7.** Subgroup of a cyclic group is cyclic.

*Proof.* Let  $G = \langle a \rangle$  be a cyclic group generated by  $a \in G$ . Let  $H \subseteq G$  be a subgroup of  $G$ . If  $H = \{e\}$  is the trivial subgroup of  $G$ , then  $H = \langle e \rangle$ . Suppose that  $H \neq \{e\}$ . Then there exists  $b \in G$  such that  $b \neq e$  and  $b \in H$ . Since  $G = \langle a \rangle$ , we have  $b = a^n$ , for some  $n \in \mathbb{Z}$ . Since  $H$  is a group and  $a^n = b \in H$ , we have  $a^{-n} = b^{-1} \in H$ . Therefore,

$$S := \{k \in \mathbb{N} : a^k \in H\} \subseteq \mathbb{N}$$

is a non-empty subset of  $\mathbb{N}$ . Then by well-ordering principle of  $(\mathbb{N}, \leq)$  (see Theorem 1.1.25)  $S$  has a least element, say  $m \in S$ . We claim that  $H = \langle a^m \rangle$ . Clearly  $\langle a^m \rangle \subseteq H$ . Let  $h \in H$  be arbitrary. Since  $H \subseteq G = \langle a \rangle$ , we have  $h = a^n$ , for some  $n \in \mathbb{Z}$ . Then by division algorithm (see Theorem 1.3.1) there exists  $q, r \in \mathbb{Z}$  with  $0 \leq r < m$  such that  $n = mq + r$ . Then  $a^r = a^{n-mq} = a^n(a^m)^{-q} = h(a^m)^{-q} \in H$ . Since  $m$  is the least element of  $S$ , we must have  $r = 0$ . Then  $n = mq$ , and so we have  $h = a^n = a^{mq} \in \langle a^m \rangle$ . Therefore,  $H \subseteq \langle a^m \rangle$ , and hence  $H = \langle a^m \rangle$ .  $\square$

**Exercise 2.3.8.** Show that any subgroup of  $\mathbb{Z}$  is of the form  $n\mathbb{Z} := \{nk : k \in \mathbb{Z}\}$ , for some  $n \in \mathbb{Z}$ .

**Lemma 2.3.9.** Let  $G = \langle a \rangle$  be an infinite cyclic group. Then for all  $m, n \in \mathbb{Z}$  with  $m \neq n$ , we have  $a^n \neq a^m$ .

*Proof.* Suppose not, then there exists  $m, n \in \mathbb{Z}$  with  $m > n$  such that  $a^m = a^n$ . Then  $a^{m-n} = a^m(a^n)^{-1} = e$ . Since  $m - n$  is a positive integer, the subset

$$S := \{k \in \mathbb{N} : a^k = e\} \subseteq \mathbb{N}$$

is non-empty. Then by well-ordering principle  $S$  has a least element, say  $d$ . We claim that  $G = \{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq d - 1\}$ . Clearly  $\{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq d - 1\} \subseteq G$ .

Let  $b \in G$  be arbitrary. Then  $b = a^n$ , for some  $n \in \mathbb{Z}$ . Then by division algorithm (Theorem 1.3.1), there exists  $q, r \in \mathbb{Z}$  with  $0 \leq r < d$  such that  $n = dq + r$ . Since  $d \in S$ , we have  $a^d = e$ . Then  $b = a^n = a^{dq+r} = (a^d)^q a^r = a^r \in \{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq d-1\}$  implies  $G \subseteq \{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq d-1\}$ , and hence  $G = \{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq d-1\}$ . This is not possible since  $G$  is infinite by our assumption. Hence the result follows.  $\square$

**Corollary 2.3.10.** *Let  $G = \langle a \rangle$  be a cyclic group generated by  $a \in G$ . Then  $G$  is infinite if and only if  $\text{ord}(a)$  is infinite.*

*Proof.* If  $G = \langle a \rangle$  is infinite, then for any non-zero integer  $n$ , we have  $a^n \neq a^0 = e$  by Lemma 2.3.9. Therefore,  $\text{ord}(a)$  is infinite. Conversely, if  $\text{ord}(a)$  is infinite, then  $a^n \neq e$ , for all  $n \in \mathbb{Z} \setminus \{0\}$ . Since  $a^n = a^m$  implies  $a^{m-n} = e$ , the map  $f : \mathbb{Z} \rightarrow G$  given by  $f(n) = a^n$ ,  $\forall n \in \mathbb{Z}$ , is injective. Therefore, since  $\mathbb{Z}$  is infinite,  $G$  must be infinite.  $\square$

**Corollary 2.3.11.** *Let  $G$  be a finite cyclic group generated by  $a$ . Then  $|G| = \text{ord}(a)$ .*

*Proof.* Since  $G$  is finite,  $\text{ord}(a)$  must be finite by Corollary 2.3.10. Suppose that  $\text{ord}(a) = n \in \mathbb{N}$ . Then for any two integers  $r, s \in \{k \in \mathbb{Z} : 0 \leq k \leq n-1\}$ ,  $a^r = a^s$  implies  $a^{r-s} = e$ , and hence  $r = s$ , because  $|r-s| < n = \text{ord}(a)$ . Then all the elements in the collection  $\mathcal{C} := \{a^k : k \in \mathbb{Z} \text{ with } 0 \leq k \leq n-1\}$  are distinct, and that  $\mathcal{C}$  has  $n$  elements. Clearly  $\mathcal{C} \subseteq G$ . Given any  $b \in G = \langle a \rangle$ ,  $b = a^m$ , for some  $m \in \mathbb{Z}$ . Then by division algorithm (Theorem 1.3.1) there exists  $q, r \in \mathbb{Z}$  with  $0 \leq r < n$  such that  $m = nq + r$ . Then  $b = a^m = a^{nq+r} = (a^n)^q a^r = a^r \in \mathcal{C}$ , since  $a^n = e$ . Therefore,  $G \subseteq \mathcal{C}$ , and hence  $G = \mathcal{C}$ . Thus,  $|G| = \text{ord}(a)$ .  $\square$

**Corollary 2.3.12.** *Let  $G$  be a finite group of order  $n$ . Then  $G$  is cyclic if and only if it contains an element of order  $n$ .*

*Proof.* If  $G$  is cyclic, then the result follows from Corollary 2.3.11. Conversely, if  $G$  contains an element  $a$  of order  $n$ , then it follows from the proof of Corollary 2.3.11 that the cyclic subgroup  $\langle a \rangle$  of  $G$  has  $n$  elements, and hence  $\langle a \rangle = G$ .  $\square$

**Corollary 2.3.13.** *Any non-trivial subgroup of an infinite cyclic group is infinite and cyclic.*

*Proof.* Let  $G$  be an infinite cyclic group generated by  $a \in G$ . Let  $H$  be a non-trivial subgroup of  $G$ . Since  $H$  is cyclic by Proposition 2.3.7, we have  $H = \langle b \rangle$ , where  $b = a^r$  for some  $r \in \mathbb{Z} \setminus \{0\}$ . Since  $G$  is an infinite cyclic group, by above Lemma 2.3.9, we have  $b^m = a^{mr} \neq a^{nr} = b^n$  for  $m \neq n$  in  $\mathbb{Z}$ . Therefore,  $H = \langle b \rangle = \{b^k : k \in \mathbb{Z}\}$  is infinite.  $\square$

**Proposition 2.3.14.** *Let  $G$  be a finite cyclic group of order  $n$ . Then for each positive integer  $d$  such that  $d \mid n$ , there is a unique subgroup  $H$  of  $G$  of order  $d$ .*

*Proof.* Let  $G = \langle a \rangle$  be a finite cyclic group of order  $n$ . Then  $\text{ord}(a) = n$  by Corollary 2.3.11. Since  $d \mid n$ , there exists  $q \in \mathbb{Z}$  such that

$$n = dq.$$

Let  $H := \langle a^q \rangle$  be the cyclic subgroup of  $G$  generated by  $a^q$ . Since  $G$  is finite, so is  $H$ . Since  $\text{ord}(a) = n$ , we see that  $d$  is the least positive integer such that  $(a^q)^d = a^{qd} = a^n = e$ . Therefore,  $\text{ord}(a^q) = d$ , and hence  $|H| = d$  by Corollary 2.3.11.

We now show uniqueness of  $H$  in  $G$ . If  $d = 1$ , then the trivial subgroup  $\{e\} \subseteq G$  is the only subgroup of  $G$  of order  $d = 1$ . Suppose that  $d > 1$ . Let  $H$  and  $K$  be two subgroups of  $G$  of order  $d$ , where  $d \mid n$ . Then by Proposition 2.3.7 we have  $H = \langle a^n \rangle$  and  $K = \langle a^m \rangle$ , for some  $m, n \in \mathbb{N}$ . Since subgroup of a finite group is finite, by Corollary 2.3.10 we have  $\text{ord}(a^n) = d = \text{ord}(a^m)$ . By division algorithm (Theorem 1.3.1) there exists unique integers  $k, r$  with  $0 \leq r < q$  such that  $m = kq + r$ . Then  $dm = kdq + dr = kn + dr$  gives

$$e = (a^m)^d = a^{dm} = (a^n)^k a^{dr} = a^{dr}.$$

Since  $0 \leq r < q$ , we have  $0 \leq dr < dq = n$ . If  $r \neq 0$ , this contradicts the fact that  $\text{ord}(a) = n$ . Therefore, we must have  $r = 0$ , and hence  $a^m = a^{kq+r} = (a^k)^q \in \langle a^k \rangle = H$ . Therefore,  $K \subseteq H$ . Since  $|H| = |K| = d$ , we have  $H = K$ .  $\square$

**Proposition 2.3.15.** *An infinite cyclic group has exactly two generators.*

*Proof.* Let  $G = \langle a \rangle = \{a^n : n \in \mathbb{Z}\}$  be an infinite cyclic group. Let  $b \in G$  be any generator of  $G$ . Then  $b = a^n$ , for some  $n \in \mathbb{Z}$ . Similarly, since  $a \in G = \langle b \rangle$ , we have  $a = b^m$ , for some  $m \in \mathbb{Z}$ . Then we have  $a = b^m = (a^n)^m = a^{mn}$ . Then by Lemma 2.3.9 we have  $mn = 1$ . Since both  $m$  and  $n$  are integers, we must have  $m, n \in \{1, -1\}$ . Therefore,  $b \in \{a, a^{-1}\}$ .  $\square$

**Exercise 2.3.16.** Let  $G = \langle a \rangle$  be a finite cyclic group of order  $n$ . Given any  $k \in \mathbb{N}$  with  $1 \leq k \leq n-1$ , show that  $\langle a^k \rangle = G$  if and only if  $\gcd(n, k) = 1$ . Conclude that  $G$  has exactly  $\phi(n)$  number of generators, where  $\phi(n)$  is the number of elements in the set  $\{k \in \mathbb{N} : \gcd(n, k) = 1\}$ . (Hint: Use the idea of the proof of Proposition 2.3.4.)

**Remark 2.3.17.** The map  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  given by sending  $n \in \mathbb{N}$  to the cardinality of the set

$$\{k \in \mathbb{N} : 1 \leq k \leq n \text{ and } \gcd(n, k) = 1\},$$

is called the *Euler phi function*.

**Exercise 2.3.18.** Give an example of a non-abelian group  $G$  such that all of its proper subgroups are cyclic.

**Exercise 2.3.19.** Show that a non-commutative group always has a non-trivial proper subgroup.

**Exercise 2.3.20.** Show that a group having at most two non-trivial subgroups is cyclic.

**Exercise 2.3.21.** Let  $G$  be a finite group having exactly one non-trivial subgroup. Show that  $|G| = p^2$ , for some prime number  $p$ .

**Exercise 2.3.22.** Give examples of infinite abelian groups having

- (i) exactly one element of finite order;
- (ii) all of its non-trivial elements have order 2.

**Exercise 2.3.23.** (i) Show that  $(\mathbb{Q}, +)$  is not cyclic.

- (ii) Show that any finitely generated subgroup of  $(\mathbb{Q}, +)$  is cyclic.
- (iii) Conclude that  $(\mathbb{Q}, +)$  is not finitely generated.
- (iv) Give an example of a proper subgroup of  $(\mathbb{Q}, +)$  that is not cyclic.

## 2.4 Product of subgroups

**Definition 2.4.1.** Let  $G$  be a group. For any two non-empty subsets  $H$  and  $K$  of  $G$ , we define their product  $HK := \{hk : h \in H, k \in K\}$ .

**Exercise 2.4.2.** Show by example that  $HK$  need not be a group in general even if both  $H$  and  $K$  are subgroups of a group.

**Theorem 2.4.3.** *Let  $H$  and  $K$  be two subgroups of  $G$ . Then  $HK$  is a group if and only if  $HK = KH$ .*

*Proof.* Note that, for any  $h \in H$  and  $k \in K$  we have  $h = h \cdot e \in HK$  and  $k = e \cdot k \in HK$ . Therefore,  $H \subseteq HK$  and  $K \subseteq HK$ .

Suppose that  $HK$  is a group. Then  $kh \in HK$ , for all  $h \in H \subseteq HK$  and  $k \in K \subseteq HK$ , and hence  $KH \subseteq HK$ . Let  $h \in H$  and  $k \in K$ . Since  $HK$  is a group,  $hk \in HK$  implies  $(hk)^{-1} \in HK$ , and so  $(hk)^{-1} = h_1k_1$ , for some  $h_1 \in H$  and  $k_1 \in K$ . Then  $hk = ((hk)^{-1})^{-1} = k_1^{-1}h_1^{-1} \in KH$ . Therefore,  $HK \subseteq KH$ , and hence  $HK = KH$ .

Conversely suppose that  $HK = KH$ . Let  $h_1k_1, h_2k_2 \in HK$  with  $h_1, h_2 \in H$  and  $k_1, k_2 \in K$ . Since  $k_2^{-1}h_2^{-1} \in KH = HK$ , there exists  $h_3 \in H$  and  $k_3 \in K$  such that  $k_2^{-1}h_2^{-1} = h_3k_3$ . Again  $k_1h_3 \in KH = HK$  implies there exists  $h_4 \in H$  and  $k_4 \in K$  such that  $k_1h_3 = h_4k_4$ . Now

$$\begin{aligned} (h_1k_1)(h_2k_2)^{-1} &= h_1k_1k_2^{-1}h_2^{-1} \\ &= h_1k_1h_3k_3 \\ &= h_1h_4k_4k_3 \in HK. \end{aligned}$$

Therefore,  $HK$  is a subgroup of  $G$ . □

**Corollary 2.4.4.** *If  $H$  and  $K$  are subgroups of a commutative group, then  $HK$  is a group.*

**Notation:** For a finite set  $S$ , we denote by  $|S|$  the number of elements of  $S$ .

**Remark 2.4.5.** The phrase “number of elements of  $S$ ” is ambiguous when  $S$  is not a finite set. For example, both  $\mathbb{Z}$  and  $\mathbb{R}$  are infinite sets, but there are some considerable differences between “the number of elements” of them;  $\mathbb{Z}$  is a countable set, while  $\mathbb{R}$  is an uncountable set. So the “number of elements” (whatever that means) for  $\mathbb{Z}$  and  $\mathbb{R}$  should not be the same. For this reason, we need an appropriate concept of “number of elements” for an infinite set  $S$ , known as the *cardinality* of  $S$ , also denoted by  $|S|$ . When  $S$  is a finite set, the cardinality of  $S$  is determined by the number of elements of  $S$ . The cardinality of  $\mathbb{Z}$  is denoted by  $\aleph_0$  (aleph-naught) and the cardinality of  $\mathbb{R}$  is  $2^{\aleph_0}$ , which is also denoted by  $\aleph_1$  or  $\mathfrak{c}$ .

**Definition 2.4.6.** The *order* of a group  $G$  is the cardinality  $|G|$  of its underlying set  $G$ . For a finite group, its order is precisely the number of elements in it.

For example, the order of  $S_3$  is 6, while the order of  $\mathbb{Z}$  is  $\aleph_0$ .

**Lemma 2.4.7.** *If  $H$  and  $K$  are finite subgroups of a group  $G$ , then*

$$|HK| = \frac{|H| \cdot |K|}{|H \cap K|}.$$

*Proof.* For each positive integer  $n$ , let  $J_n := \{k \in \mathbb{N} : k \leq n\}$ . Let  $H = \{h_i : i \in J_n\}$  and  $K = \{k_j : j \in J_m\}$ . Then  $HK = \{h_ik_j : i \in J_n, j \in J_m\}$ . To find the number of elements of  $HK$ , for each pair  $(i, j) \in J_n \times J_m$ , we need to count the number of times  $h_ik_j$  repeats in the collection  $\mathcal{C} := \{h_ik_j : (i, j) \in J_n \times J_m\}$ . Fix  $(i, j) \in J_n \times J_m$ . If  $h_ik_j = h_pk_q$ , for some  $(p, q) \in J_n \times J_m$ , then  $t := h_p^{-1}h_i = k_qk_j^{-1} \in H \cap K$ . So any element  $h_pk_q \in \mathcal{C}$ , which coincides with  $h_ik_j$  is of the form  $(h_it^{-1})(tk_j)$ , for some  $t \in H \cap K$ . Conversely, for any  $t \in H \cap K$ , we have  $(h_it^{-1})(tk_j) = h_i(t^{-1}t)k_j = h_iek_j = h_ik_j$ . Therefore, the element  $h_ik_j$  appears exactly  $|H \cap K|$ -times in the collection  $\mathcal{C}$ , and hence we have

$$|HK| = \frac{|H| \cdot |K|}{|H \cap K|}.$$

This completes the proof. □

**Proposition 2.4.8.** *Let  $H$  and  $K$  be subgroups of  $G$ . Then  $HK$  is a subgroup of  $G$  if and only if  $HK = \langle H \cup K \rangle$ .*

*Proof.* Suppose that  $HK$  is a subgroup of  $G$ . Since  $H \subseteq HK$  and  $K \subseteq HK$ , we have  $H \cup K \subseteq HK$ , and hence  $\langle H \cup K \rangle \subseteq HK$ . Since  $\langle H \cup K \rangle$  is a group containing  $H \cup K$ , for any  $h \in H$  and  $k \in K$  we have  $hk \in \langle H \cup K \rangle$ . Therefore,  $HK \subseteq \langle H \cup K \rangle$ , and hence  $HK = \langle H \cup K \rangle$ . Converse is obvious since  $\langle H \cup K \rangle$  is a group and  $HK = \langle H \cup K \rangle$  by assumption.  $\square$

### 2.4.1 Lattice diagram

**Definition 2.4.9.** A relation “ $\leq$ ” on a non-empty set  $S$  is said to be a *partial order relation* if it is reflexive, anti-symmetric and transitive (see Definition 2.1.27). A *partially ordered set* (or, in short a *poset*) is a pair  $(S, \leq)$ , where  $S$  is a non-empty set together with a partial order relation “ $\leq$ ” on it.

Let  $(S, \leq)$  be a poset. Given a collection of elements  $\{a_\lambda : \lambda \in \Lambda\}$  from  $S$ , an element  $c \in S$  is said to be an

- (i) *upper bound* of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$  if  $a_\lambda \leq c, \forall \lambda \in \Lambda$ .
- (ii) *lower bound* of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$  if  $c \leq a_\lambda, \forall \lambda \in \Lambda$ .

An element  $c_0 \in S$  is said to be

- (i) a *least upper bound* of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$  if
  - $c_0$  is an upper bound of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ , and
  - if  $d$  is an upper bound of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ , then  $c_0 \leq d$ .
- (ii) a *greatest lower bound* of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$  if
  - $c_0$  is a lower bound of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ , and
  - if  $d$  is any lower bound of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ , then  $d \leq c_0$ .

**Lemma 2.4.10.** Let  $(S, \leq)$  be a poset. Let  $\{a_\lambda : \lambda \in \Lambda\}$  be a non-empty collection of elements of  $S$ . If least upper bound (resp., greatest lower bound) of  $\{a_\lambda : \lambda \in \Lambda\}$  exists in  $(S, \leq)$ , then it must be unique.

*Proof.* Suppose that  $c_0$  and  $d_0$  be least upper bounds of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ . Then  $c_0 \leq d_0$  and  $d_0 \leq c_0$ . Since “ $\leq$ ” is anti-symmetric, we have  $c_0 = d_0$ . The same argument shows that, the greatest upper bound of  $\{a_\lambda : \lambda \in \Lambda\}$  in  $(S, \leq)$ , if exists, is unique.  $\square$

**Definition 2.4.11 (Lattice).** A partially ordered set  $(S, \leq)$  is said to be a *lattice* if the least upper bound and the greatest lower bound of any two elements of  $S$  exist in  $S$ .

**Proposition 2.4.12.** Let  $G$  be a group, and let  $S$  be the set of all subgroups of  $G$ . Define a relation  $\leq$  on  $S$  by setting

$$H \leq K \text{ if } H \subseteq K.$$

Then  $(S, \leq)$  is a lattice.


*Proof.* Clearly “ $\leq$ ” is a partial ordering relation on  $S$  (verify). Let  $H$  and  $K$  be any two subgroups of  $G$ . As we have noticed before,  $\langle H \cup K \rangle$  is the smallest subgroup of  $G$  containing  $H$  and  $K$ , it is the least upper bound of  $\{H, K\}$  inside  $(S, \leq)$ . Since  $H \cap K \leq H$  and  $H \cap K \leq K$ , and for any subgroup  $J$  of  $G$  with  $J \subseteq H$  and  $J \subseteq K$ , we have  $J \subseteq H \cap K$ , we see that  $H \cap K$  is the greatest lower bound of  $\{H, K\}$  in  $(S, \leq)$ .  $\square$

**Definition 2.4.13.** Let  $G$  be a group. Given any two subgroups  $H$  and  $K$  of  $G$ , if  $H \leq K$ , we place  $H$  below  $K$  and draw a vertical line segment between them to indicate that  $H$  is sitting inside  $K$ . This process generates a diagram, known as the *lattice diagram of subgroups of  $G$* .

**Example 2.4.14.** Consider the group  $\mu_4 = \{\zeta \in \mathbb{C}^* : \zeta^4 = 1\} = \{1, -1, \sqrt{-1}, -\sqrt{-1}\}$  of 4<sup>th</sup> roots of unity. Note that  $\mu_2 := \{1, -1\}$  and  $\mu_1 := \{1\}$  are only subgroups of  $G = \mu_4$ , and that  $\mu_1 \leq \mu_2 \leq \mu_4$ . Then the lattice structure of  $\mu_4$  can be written as

$$\begin{array}{c} \mu_4 = \{1, -1, \sqrt{-1}, -\sqrt{-1}\} \\ | \\ \mu_2 = \{1, -1\} \\ | \\ \mu_1 = \{1\} \end{array}$$

**Exercise 2.4.15.** Write down all subgroups of the symmetric group  $S_3$  and the associated lattice structure. The subgroup of  $S_3$  generated by a 2-cycle  $\sigma \in S_3$  consist of  $\sigma$  and the neutral element only. There are three such subgroups. There are only two 3-cycles in  $S_3$ , namely  $(1\ 2\ 3)$ ,  $(1\ 3\ 2)$ , and they satisfies  $(1\ 2\ 3)^2 = (1\ 3\ 2)$  and  $(1\ 2\ 3)^3 = e$ . So,  $\langle (1\ 2\ 3) \rangle = \{e, (1\ 2\ 3), (1\ 3\ 2)\}$ . Thus the lattice structure of  $(S_3, \leq)$  can be written as follows.



## 2.5 Permutation Groups

Let  $X$  be a non-empty set. A *permutation* on  $X$  is a bijective map  $\sigma : X \rightarrow X$ . We denote by  $S_X$  the set of all permutations on  $X$ . For notational simplicity, when  $|X| = n$ , fixing a bijection of  $X$  with the subset  $J_n := \{1, 2, 3, \dots, n\} \subset \mathbb{N}$  we may identify  $S_X$  with  $S_n$ . An element  $\sigma \in S_n$  can be described by a *two-column notation* as follow.

$$(2.5.1) \quad \sigma = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ \sigma(1) & \sigma(2) & \sigma(3) & \cdots & \sigma(n) \end{pmatrix} \quad \text{or,} \quad \sigma = \begin{cases} 1 \mapsto \sigma(1) \\ 2 \mapsto \sigma(2) \\ \vdots \\ n \mapsto \sigma(n) \end{cases}.$$

Since elements of  $S_n$  are bijective maps of  $J_n$  onto itself, composition of two elements of  $S_n$  is again an element of  $S_n$ . Thus we have a binary operation

$$\circ : S_n \times S_n \longrightarrow S_n, \quad (\sigma, \tau) \longmapsto \tau \circ \sigma.$$

For example, consider the elements  $\sigma, \tau \in S_4$  defined by

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 1 & 3 \end{pmatrix}, \quad \tau = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 2 & 4 \end{pmatrix}.$$



Then their composition  $\tau \circ \sigma$  is the permutation

$$\tau \circ \sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix}$$

Clearly composition of functions  $J_n \rightarrow J_n$  is associative, and for any  $\sigma \in S_n$  its pre-composition and post-composition with the identity map of  $I_n$  is  $\sigma$  itself. Also inverse of a bijective map is again bijective. Thus for all integer  $n \geq 1$ ,  $(S_n, \circ)$  is a group, called the *Symmetric group* (or, the *permutation group*) on  $J_n$ .

**Remark 2.5.2.** For each integer  $n \geq 0$ , the symmetric group  $S_{n+1}$  can be understood as the group of symmetries of a regular  $n$ -simplex inside  $\mathbb{R}^{n+1}$ . The *standard  $n$ -simplex*

$$\Delta^n := \{(t_0, \dots, t_n) \in \mathbb{R}^{n+1} : \sum_{j=0}^n t_j = 1, t_j \geq 0, \forall j = 0, 1, \dots, n\} \subset \mathbb{R}^{n+1}$$

is an example of a regular  $n$ -simplex. This has vertices the unit vectors  $\{e_0, e_1, \dots, e_n\}$  in  $\mathbb{R}^{n+1}$ , where

$$\begin{aligned} e_0 &= (1, 0, 0, \dots, 0, 0), \\ e_1 &= (0, 1, 0, \dots, 0, 0), \\ &\vdots \\ e_n &= (0, 0, 0, \dots, 0, 1). \end{aligned}$$

For example,

- $\Delta^0$  is a point,
- $\Delta^1$  is the straight line segment  $[-1, 1] \subset \mathbb{R} \subset \mathbb{R}^2$ ,
- $\Delta^2$  is an equilateral triangle in the plane  $\mathbb{R}^2$ ,
- $\Delta^3$  is a regular tetrahedron in  $\mathbb{R}^3$ , and so on.

**Exercise 2.5.3.** Show that  $S_1$  is a trivial group, and  $S_2$  is an abelian group with two elements.

**Lemma 2.5.4.** For all integer  $n \geq 3$ , the group  $S_n$  is non-commutative.

*Proof.* Let  $\sigma, \tau \in S_n$  be defined by

$$\sigma(k) = \begin{cases} 2, & \text{if } k = 1 \\ 1, & \text{if } k = 2 \\ k, & \text{if } k \in I_n \setminus \{1, 2\} \end{cases}, \quad \text{and } \tau(k) = \begin{cases} 3, & \text{if } k = 1 \\ 1, & \text{if } k = 3 \\ k, & \text{if } k \in I_n \setminus \{1, 3\} \end{cases}.$$

Since  $\tau \circ \sigma(1) = 2$  and  $\sigma \circ \tau(1) = 3$ , we have  $\sigma \circ \tau \neq \tau \circ \sigma$ . Therefore,  $S_n$  is non-commutative.  $\square$

Let  $\sigma \in S_n$  be given. Consider its two-column notation as in (2.5.1).

(R1) If  $\sigma(k) = k$ , for some  $k \in J_n$ , we may drop the corresponding column from its two-column notation, and rearrange its columns, if required, to get an expression of the form

$$\sigma = \begin{pmatrix} k_1 & k_2 & \cdots & k_{r-1} & k_r \\ \sigma(k_1) & \sigma(k_2) & \cdots & \sigma(k_{r-1}) & \sigma(k_r) \end{pmatrix},$$

where  $k_1, \dots, k_r$  are all distinct.

By re-indexing, if required, we can find a partition of  $\{k_1, \dots, k_r\}$  into *disjoint subsets*, say

$$\{k_1, \dots, k_r\} = \bigcup_{i=1}^m \{k_{i,1}, \dots, k_{i,r_i}\}$$



with  $m \geq 1$ ,  $2 \leq r_i \leq r$ , for all  $i \in \{1, \dots, m\}$ , and  $r_1 + \dots + r_m = r$ , such that for all  $i \in \{1, \dots, m\}$  we have

$$(2.5.5) \quad \sigma(k_{i,j}) = \begin{cases} k_{i,j+1}, & \text{if } j \in \{1, \dots, r_i - 1\}, \\ k_{i,1}, & \text{if } j = r_i, \text{ and} \\ k_{ij}, & \text{if } k_{ij} \in J_n \setminus \{k_1, \dots, k_r\}. \end{cases}$$

Then  $\sigma$  can be expressed as

$$(2.5.6) \quad \sigma = \begin{pmatrix} k_{1,1} & \cdots & k_{1,r_1-1} & k_{1,r_1} & \cdots & k_{m,1} & \cdots & k_{m,r_m} & k_{m,r_m-1} \\ k_{1,2} & \cdots & k_{1,r_1} & k_{1,1} & \cdots & k_{m,2} & \cdots & k_{m,r_m} & k_{m,1} \end{pmatrix}.$$

When  $m = 1$  in the above notation,  $\sigma$  can be expressed as

$$(2.5.7) \quad \sigma = \begin{pmatrix} k_1 & k_2 & \cdots & k_{r-1} & k_r \\ k_2 & k_3 & \cdots & k_r & k_1 \end{pmatrix}.$$

Such a permutation is called a cycle.

**Definition 2.5.8 (Cycle).** An element  $\sigma \in S_n$  is called a  $r$ -cycle or a cycle of length  $r$  if there exists distinct  $r$  elements, say  $k_1, \dots, k_r \in J_n := \{1, \dots, n\}$  such that  $\sigma(k) = k$ , for all  $k \in J_n \setminus \{k_1, \dots, k_r\}$  and

$$\sigma(k_i) = \begin{cases} k_{i+1} & \text{if } i \in \{1, \dots, r-1\}, \\ k_1 & \text{if } i = r. \end{cases}$$

In this case,  $\sigma$  is expressed as  $\sigma = (k_1 \ k_2 \ \cdots \ k_r)$ . A 2-cycle is called a *transposition*.

**Remark 2.5.9.** Note that according to our definition 2.5.8, a cycle in  $S_n$  always have length at least 2. So we don't talk about 1-cycle as used in some of the standard text books.

With the notation above, the permutation  $\sigma$  in (2.5.6) can be written as a product of cycles

$$\begin{aligned} \sigma &= \begin{pmatrix} k_{1,1} & \cdots & k_{1,r_1-1} & k_{1,r_1} \\ k_{1,2} & \cdots & k_{1,r_1} & k_{1,1} \end{pmatrix} \circ \cdots \circ \begin{pmatrix} k_{m,1} & \cdots & k_{m,r_m-1} & k_{m,r_m} \\ k_{m,2} & \cdots & k_{m,r_m} & k_{m,1} \end{pmatrix} \\ &= (k_{1,1} \ \cdots \ k_{1,r_1-1} \ k_{1,r_1}) \circ \cdots \circ (k_{m,1} \ \cdots \ k_{m,r_m-1} \ k_{m,r_m}) \end{aligned}$$

**Remark 2.5.10.** Transpositions are of particular interests. We shall see later that any  $\sigma \in S_n$  can be written as product of either even number of transpositions or odd number of transpositions, and accordingly we call  $\sigma \in S_n$  an even permutation or an odd permutation.

**Example 2.5.11.** Using cycle notation, the group  $S_3$  can be written as

$$S_3 = \{e, (1 \ 2), (1 \ 3), (2 \ 3), (1 \ 2 \ 3), (1 \ 3 \ 2)\},$$

where  $(1 \ 2)$ ,  $(1 \ 3)$  and  $(2 \ 3)$  are transpositions. However, we can write 3-cycles as product of 2-cycles as  $(1 \ 2 \ 3) = (2 \ 3) \circ (1 \ 3)$  and  $(1 \ 3 \ 2) = (2 \ 3) \circ (1 \ 2)$ . Also, the identity element  $e$  can be written as  $e = (1 \ 2) \circ (1 \ 2)$  or  $e = (1 \ 3) \circ (1 \ 3)$  etc. So the decomposition of  $\sigma \in S_n$  as a product of transpositions is not unique.

**Proposition 2.5.12.** Let  $\sigma = (k_1 \ k_2 \ \cdots \ k_r) \in S_n$  be a  $r$ -cycle. Then for any  $\tau \in S_n$  we have

$$\tau \sigma \tau^{-1} = (\tau(k_1) \ \tau(k_2) \ \cdots \ \tau(k_r)).$$

*Proof.* Note that we have

$$\begin{aligned} (\tau \sigma \tau^{-1})(\tau(k_i)) &= \tau(\sigma(k_i)) = \tau(k_{i+1}), \quad \forall i \in \{1, \dots, r-1\}, \\ \text{and } (\tau \sigma \tau^{-1})(\tau(k_r)) &= \tau(\sigma(k_r)) = \tau(k_1). \end{aligned}$$

It remains to show that  $(\tau\sigma\tau^{-1})(k) = k$ ,  $\forall k \in J_n \setminus \{\tau(k_1), \dots, \tau(k_r)\}$ . For this, note that  $\tau^{-1}(k) \in J_n \setminus \{k_1, \dots, k_r\}$ , and so  $\sigma(\tau^{-1}(k)) = \tau^{-1}(k)$ . Therefore, we have  $(\tau\sigma\tau^{-1})(k) = \tau(\sigma(\tau^{-1}(k))) = \tau(\tau^{-1}(k)) = k$ . This completes the proof.  $\square$

**Corollary 2.5.13.** Let  $\sigma \in S_n$  is a product of pairwise disjoint cycles  $\sigma_1, \dots, \sigma_r$  in  $S_n$ . Suppose that  $\sigma_i = (k_{i1} \dots k_{i\ell_i}) \in S_n$ , for all  $i \in \{1, \dots, r\}$ . Then for any  $\tau \in S_n$  we have  $\tau\sigma\tau^{-1} = (\tau(k_{11}) \dots \tau(k_{1\ell_1})) \circ \dots \circ (\tau(k_{r1}) \dots \tau(k_{r\ell_r}))$ . In particular, both  $\sigma$  and  $\tau\sigma\tau^{-1}$  have the same cycle type.

*Proof.* Since  $\tau\sigma\tau^{-1} = (\tau\sigma_1\tau^{-1}) \circ \dots \circ (\tau\sigma_r\tau^{-1})$ , the result follows from Proposition 2.5.12.  $\square$

**Proposition 2.5.14.** Let  $\sigma \in S_n$  be a cycle. Then  $\sigma$  is a  $r$  cycle if and only if  $\text{ord}(\sigma) = r$ .

*Proof.* Let  $\sigma = (k_1 \ k_2 \ \dots \ k_r)$ , for some distinct elements  $k_1, \dots, k_r \in J_n$ . Then for any  $k \in J_n \setminus \{k_1, \dots, k_r\}$  we have  $\sigma(k) = k$ . It follows from the definition of the cyclic expression of  $\sigma$  given in (2.5.5) that  $\sigma^i(k_1) = k_{i+1}$ , for all  $i \in \{1, \dots, r-1\}$  and  $\sigma^r(k_1) = k_1$ . In general, for any  $k_i$  with  $1 \leq i \leq r$  we have  $\sigma^{r-i}(k_i) = k_r$  and so  $\sigma^{r-i+1}(k_i) = k_1$ . Therefore,  $\sigma^{r-i+\ell}(k_i) = k_\ell$  for all  $\ell \in \{1, \dots, r-1\}$ , and hence  $\sigma^r(k_i) = k_i$ , for all  $i \in \{1, \dots, r\}$ . Combining all these, we have  $\sigma^r(k) = k$ , for all  $k \in J_n$ . In other words,  $\sigma^r = e$ , where  $e$  is the identity element in  $S_n$ . Since  $\sigma^s(k_1) = k_{s+1}$ , for all  $s \in \{1, \dots, r-1\}$  (see (2.5.5)), we conclude that  $r$  is the smallest positive integer such that  $\sigma^r = e$  in  $S_n$ . Therefore,  $\text{ord}(\sigma) = r$ . Conversely, suppose that  $\sigma$  is a  $t$  cycle with  $\text{ord}(\sigma) = r$ . But then as shown above  $\text{ord}(\sigma) = t$ , and hence  $t = r$ .  $\square$

**Exercise 2.5.15.** Show that the number of distinct  $r$  cycles in  $S_n$  is  $\frac{n!}{r(n-r)!}$ .

*Solution:* Note that, we can choose a  $r$  cycle from  $S_n$  in

$${}^nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

ways. Fix a  $r$ -cycle  $\sigma = (k_1 \ k_2 \ \dots \ k_r) \in S_n$ . Note that, the cycles

$$(k_1 \ k_2 \ \dots \ k_r) \text{ and } (k_2 \ k_3 \ \dots \ k_r \ k_1)$$

represents the same element  $\sigma \in S_n$ . Note that, given any two permutations (bijective maps)

$$\phi, \psi : \{2, 3, \dots, r\} \rightarrow \{2, 3, \dots, r\},$$

two  $r$  cycles (note that  $k_1$  is fixed!)

$$(k_1 \ k_{\phi(2)} \ \dots \ k_{\phi(r)}) \text{ and } (k_1 \ k_{\psi(2)} \ \dots \ k_{\psi(r)})$$

represents the same element of  $S_n$  if and only if  $\phi = \psi$ . Since there are  $(r-1)!$  number of distinct bijective maps  $\{2, 3, \dots, r\} \rightarrow \{2, 3, \dots, r\}$  (verify!), fixing  $k_1$  in one choice of  $r$  cycle  $(k_1 \ k_2 \ \dots \ k_r)$  in  $S_n$ , considering all permutations of the remaining  $(r-1)$  entries  $k_2, \dots, k_r$ , we get  $(r-1)!$  number of distinct  $r$  cycles in  $S_n$ . Therefore, the total number of distinct  $r$  cycles in  $S_n$  is precisely

$$(r-1)! \cdot \frac{n!}{r!(n-r)!} = \frac{n!}{r(n-r)!}.$$

This completes the proof.  $\square$

**Definition 2.5.16.** Two cycles  $\sigma = (i_1 \ i_2 \ \dots \ i_r)$  and  $\tau = (j_1 \ j_2 \ \dots \ j_s)$  in  $S_n$  are said to be *disjoint* if  $\{i_1, i_2, \dots, i_r\} \cap \{j_1, j_2, \dots, j_s\} = \emptyset$ .

**Proposition 2.5.17.** If  $\sigma$  and  $\tau$  are disjoint cycles in  $S_n$ , show that  $\sigma \circ \tau = \tau \circ \sigma$ .

*Proof.* Let  $\sigma = (i_1 \ i_2 \ \cdots \ i_r)$  and  $\tau = (j_1 \ j_2 \ \cdots \ j_s)$  be two disjoint cycles in  $S_n$ . Let  $k \in J_n$  be arbitrary. If  $k \notin \{i_1, \dots, i_r\} \cup \{j_1, \dots, j_s\}$ , then  $\sigma(k) = k = \tau(k)$  and hence  $(\sigma\tau)(k) = (\tau\sigma)(k)$  in this case. Suppose that  $k \in \{i_1, \dots, i_r\}$ . Then  $\sigma(k) \in \{i_1, \dots, i_r\}$  and  $k \notin \{j_1, \dots, j_s\}$  together gives  $\tau\sigma(k) = \sigma(k) = \sigma\tau(k)$ . Interchanging the roles of  $\sigma$  and  $\tau$  we see that  $\tau\sigma(k) = \sigma(k) = \sigma\tau(k)$  holds for the case  $k \in \{j_1, \dots, j_s\}$ . Therefore,  $\sigma\tau = \tau\sigma$ .  $\square$

**Lemma 2.5.18.** *For  $n \geq 2$ , any non-identity element of  $S_n$  can be uniquely written as a product of disjoint cycles of length at least 2. This expression is unique up to ordering of factors.*

*Proof.* For  $n = 2$ ,  $S_2$  has only one non-identity element, which is a 2-cycle  $(1 \ 2)$ . Assume that  $n \geq 3$  and the result is true for any non-identity element of  $S_r$  for  $2 \leq r < n$ . Let  $\sigma \in S_n$  be a non-identity element. Since  $\{\sigma^i(1) : i \in \mathbb{N}\} \subseteq J_n$  and  $J_n$  is a finite set, there exists distinct integers  $i, j \in \mathbb{N}$  such that  $\sigma^i(1) = \sigma^j(1)$ . Without loss of generality we may assume that  $i - j \geq 1$ . Then  $\sigma^{i-j}(1) = 1$ . Then

$$\{i \in \mathbb{N} : \sigma^i(1) = 1\}$$

is a non-empty subset of  $\mathbb{N}$ , and hence it has a least element, say  $r$ . Then all the elements in

$$A := \{1, \sigma(1), \sigma^2(1), \dots, \sigma^{r-1}(1)\}$$

are all distinct, and defines an  $r$ -cycle

$$\tau := (1 \ \sigma(1) \ \sigma^2(1) \ \cdots \ \sigma^{r-1}(1))$$

in  $S_n$ . Let  $B := J_n \setminus A$ . In cases  $\sigma|_B$  is the identity map of  $B$  onto itself or  $B = \emptyset$ , we have  $\tau = \sigma$  and so  $\sigma$  is a cycle in  $S_n$ . Assume that  $B \neq \emptyset$  and  $\pi := \sigma|_B$  is not the identity map. Then  $\pi$  is a non-identity element of  $S_k$ , where  $2 \leq k := |B| < n$ . Then by induction hypothesis  $\pi = \pi_1 \cdots \pi_\ell$  is a finite product of disjoint cycles  $\pi_1, \dots, \pi_\ell$  of lengths at least 2 in  $S_k$ . Then for each  $i \in \{1, \dots, \ell\}$  we define  $\sigma_i \in S_n$  by setting

$$\sigma_i(a) = \begin{cases} \pi_i(a), & \text{if } a \in B, \\ a, & \text{if } a \in J_n \setminus B. \end{cases}$$

Then  $\sigma_1, \dots, \sigma_\ell, \tau$  are pairwise disjoint cycles in  $S_n$  and that  $\sigma = \sigma_1 \cdots \sigma_\ell \tau$ .

For the uniqueness part, let  $\sigma = \sigma_1 \cdots \sigma_r = \tau_1 \cdots \tau_s$  be two decomposition of  $\sigma$  into product of disjoint cycles of lengths  $\geq 2$  in  $S_n$ . We need to show that  $r = s$ , and there is a permutation  $\delta \in S_r$  such that  $\sigma_i = \tau_{\delta(i)}$ , for all  $i \in \{1, \dots, r\}$ . Suppose that  $\sigma_i = (k_1 \ k_2 \ \cdots \ k_t)$  with  $t \geq 2$ . Then  $\sigma(k_1) \neq k_1$ . Since  $\tau_1, \dots, \tau_r$  are pairwise disjoint cycles of lengths  $\geq 2$  in  $S_n$ , there is a unique element, say  $\delta(i) \in \{1, \dots, r\}$  such that  $\tau_{\delta(i)}(k_1) \neq k_1$ . By reordering, if required, we may write  $\tau_{\delta(i)} = (k_1 \ v_2 \ \cdots \ v_u)$ . Then we have

$$\begin{array}{ccccccccc} k_2 & = & \sigma_i(k_1) & = & \sigma(k_1) & = & \tau_{\delta(i)}(k_1) & = & v_2, \\ k_3 & = & \sigma_i(k_2) & = & \sigma(k_2) & = & \tau_{\delta(i)}(v_2) & = & v_3, \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ k_t & = & \sigma_i(k_{r-1}) & = & \sigma(k_{r-1}) & = & \tau_{\delta(i)}(v_{r-1}) & = & v_t. \end{array}$$

If  $t < u$ , then  $k_1 = \sigma_i(k_t) = \sigma(k_t) = \tau_{\delta(i)}(v_t) = v_{t+1}$ , which is a contradiction. Therefore,  $t = u$  and hence  $\sigma_i = \tau_{\delta(i)}$ . Hence the result follows by induction on  $r$ .  $\square$

**Definition 2.5.19 (Cycle type).** Given  $\sigma \in S_n$ , by Lemma 2.5.18 there exists a unique finite set of pairwise disjoint cycles  $\{\sigma_1, \dots, \sigma_r\}$  in  $S_n$  such that  $\sigma = \sigma_1 \circ \cdots \circ \sigma_r$ . Since disjoint cycles commutes by Proposition 2.5.17, by reindexing  $\sigma_j$ 's, if required, we may assume that  $n_1 \geq \dots \geq n_r$ , where  $n_j = \text{length}(\sigma_j)$ , for all  $j \in \{1, \dots, r\}$ . Since  $\sigma_1, \dots, \sigma_r$  are pairwise disjoint cycles in  $S_n$ , we have  $\ell + \sum_{j=1}^r n_j = n$ , for some non-negative integer  $\ell$ . If  $\ell = 0$ , then the sequence

$(n_1, \dots, n_r)$  is called the *cycle type* of  $\sigma$ , and if  $\ell > 0$ , then the sequence  $(n_1, \dots, n_r, f_1, \dots, f_\ell)$ , where  $f_1 = \dots = f_\ell = 1$ , is called the cycle type of  $\sigma$ .

**Example 2.5.20.** (i) The cycle type of  $\sigma := (1\ 2) \circ (3\ 6) \circ (4\ 5\ 7) \in S_7$  is  $(3, 2, 2)$ .

(ii) The cycle type of  $\tau := (1\ 4\ 3) \circ (2\ 5) \in S_7$  is  $(3, 2, 1, 1)$ .

(iii) The cycle type of  $\delta := (1\ 3\ 5) \circ (2\ 4\ 7) \in S_6$  is  $(3, 3, 1)$ .

**Definition 2.5.21.** Two permutations  $\sigma$  and  $\tau$  in  $S_n$  are said to be *conjugate* in  $S_n$  if there exists  $\delta \in S_n$  such that  $\tau = \delta \circ \sigma \circ \delta^{-1}$ .

**Theorem 2.5.22.** Two elements  $\sigma, \tau \in S_n$  are conjugate if and only if they have the same cycle type.

*Proof.* Conjugate permutations in  $S_n$  have the same cycle type by Corollary 2.5.13. Conversely suppose that  $\sigma, \tau \in S_n$  have the same cycle type, say  $(n_1, \dots, n_r, f_1, \dots, f_\ell)$ , where  $n_1 \geq \dots \geq n_r \geq 2$  and  $f_1 = \dots = f_\ell = 1$ ,  $\ell \geq 0$  and that  $\sum_{j=1}^r n_j + \ell = n$ . Let  $\sigma = \sigma_1 \circ \dots \circ \sigma_r$  and  $\tau = \tau_1 \circ \dots \circ \tau_r$ , where  $\sigma_i, \tau_j$  are cycles in  $S_n$  of lengths  $n_i$  and  $n_j$ , respectively. Suppose that  $\sigma_i = (a_{i1} \ \dots \ a_{in_i})$  and  $\tau_j = (b_{j1} \ \dots \ b_{jn_j})$ . If  $\ell > 0$ , then we write the subset  $I_n \setminus \{a_{ij} : 1 \leq i \leq r, 1 \leq j \leq n_i\}$  as  $\{a_1, \dots, a_\ell\}$ . Then  $I_n$  is a disjoint union of the subsets  $\{a_{11}, \dots, a_{1n_1}\}, \dots, \{a_{r1}, \dots, a_{rn_r}\}, \{a_1, \dots, a_\ell\}$ . Similarly if we write the subset  $I_n \setminus \{b_{ij} : 1 \leq i \leq r, 1 \leq j \leq n_i\}$  as  $\{b_1, \dots, b_\ell\}$ , then  $I_n$  is a disjoint union of the subsets  $\{b_{11}, \dots, b_{1n_1}\}, \dots, \{b_{r1}, \dots, b_{rn_r}\}, \{b_1, \dots, b_\ell\}$ . Then we define a map  $\delta : I_n \rightarrow I_n$  by sending  $a_{ij}$  to  $b_{ij}$ , for all  $(i, j) \in \{1, \dots, r\} \times \{1, \dots, n_i\}$ , and by sending  $a_k$  to  $b_k$ , for all  $k \in \{1, \dots, \ell\}$ , if  $\ell > 0$ . Clearly  $\delta$  is a bijective map, and hence is an element of  $S_n$ . Then Proposition 2.5.12 ensures that  $\delta \sigma_i \delta^{-1} = \tau_i$ , for all  $i \in \{1, \dots, r\}$ . Then we have

$$\begin{aligned} \delta \sigma \delta^{-1} &= \delta(\sigma_1 \dots \sigma_r) \delta^{-1} \\ &= (\delta \sigma_1 \delta^{-1}) \dots (\delta \sigma_r \delta^{-1}) \\ &= \tau_1 \dots \tau_r \\ &= \tau. \end{aligned}$$

This completes the proof.  $\square$

**Exercise 2.5.23.** Find the number of elements of order 2 and 3 in  $S_4$ . Show that  $S_4$  has no element of order 4.

**Corollary 2.5.24.** For  $n \geq 2$ , every element of  $S_n$  can be written as a finite product of transpositions.

*Proof.* In view of above Lemma 2.5.18 it suffices to show that every cycle of  $S_n$  is a product of transpositions. Clearly the identity element  $e \in S_n$  can be written as  $e = (1\ 2)(1\ 2)$ . If  $\sigma = (k_1\ k_2\ \dots\ k_r)$  is an  $r$ -cycle,  $r \geq 2$ , in  $S_n$ , then we can rewrite it as

$$\sigma = (k_1\ k_2\ \dots\ k_r) = (k_1\ k_r)(k_1\ k_{r-1}) \dots (k_1\ k_2).$$

Hence the result follows.  $\square$

Note that decompositions of  $\sigma \in S_n$  into a finite product of transpositions is not unique. For example, when  $n \geq 3$  we have  $e = (1\ 2)(1\ 2) = (1\ 3)(1\ 3)$ . However, we shall see shortly that the number of transpositions appearing in such a product expression for  $\sigma \in S_n$  is either odd or even, but cannot be both in two such decompositions.

**Lemma 2.5.25.** Fix an integer  $n \geq 2$ , and consider the action of a permutation  $\sigma \in S_n$  on the formal product  $\chi := \prod_{1 \leq i < j \leq n} (x_i - x_j)$  given by

$$\sigma(\chi) := \prod_{1 \leq i < j \leq n} (x_{\sigma(i)} - x_{\sigma(j)}).$$

If  $\sigma \in S_n$  is a 2-cycle (transposition), then  $\sigma(\chi) = -\chi$ .

*Proof.* Since  $\sigma \in S_n$  is a 2-cycle, there exists a unique subset  $\{p, q\} \subseteq J_n$  with  $p < q$  such that  $\sigma = (p \ q)$ . Then  $\sigma(k) = k$ ,  $\forall k \in J_n \setminus \{p, q\}$ . Consider the factor  $(x_i - x_j)$  of  $\chi$  with  $1 \leq i < j \leq n$ . We have the following situations:

- (a) If  $\{i, j\} = \{p, q\}$ , then  $\sigma(x_i - x_j) = x_{\sigma(i)} - x_{\sigma(j)} = -(x_i - x_j)$ .
- (b) If  $\{i, j\} \cap \{p, q\} = \emptyset$ , then  $\sigma(x_i - x_j) = x_{\sigma(i)} - x_{\sigma(j)} = (x_i - x_j)$ .
- (c) If  $\{i, j\} \cap \{p, q\}$  is singleton set, then we have the following subcases.
  - I. If  $t < p < q$ , then  $\sigma((x_t - x_p)(x_t - x_q)) = (x_{\sigma(t)} - x_{\sigma(p)})(x_{\sigma(t)} - x_{\sigma(q)}) = (x_t - x_q)(x_t - x_p)$ .
  - II. If  $p < t < q$ , then  $\sigma((x_p - x_t)(x_t - x_q)) = (x_{\sigma(p)} - x_{\sigma(t)})(x_{\sigma(t)} - x_{\sigma(q)}) = (x_q - x_t)(x_p - x_t)$ .
  - III. If  $p < q < t$ , then  $\sigma((x_p - x_t)(x_q - x_t)) = (x_{\sigma(p)} - x_{\sigma(t)})(x_{\sigma(q)} - x_{\sigma(t)}) = (x_q - x_t)(x_p - x_t)$ .

Therefore, in the above three subcases the product  $(x_t - x_p)(x_t - x_q)$  remains fixed under the action of  $\sigma$ .

From these it immediately follows that  $\sigma(\chi) = -\chi$ , for all 2-cycle  $\sigma \in S_n$ .  $\square$

**Corollary 2.5.26.** Fix an integer  $n \geq 2$ , and let  $\sigma \in S_n$ . If  $\sigma = \sigma_1 \cdots \sigma_r = \tau_1 \cdots \tau_s$ , where  $\sigma_i, \tau_j$  are all transpositions in  $S_n$ , then both  $r$  and  $s$  are either even or odd.

*Proof.* Consider the formal product  $\chi := \prod_{1 \leq i < j \leq n} (x_i - x_j)$ . Then  $\sigma(\chi) = (\sigma_1 \circ \cdots \circ \sigma_r)(\chi) = (-1)^r \chi$  and  $\sigma(\chi) = (\tau_1 \circ \cdots \circ \tau_s)(\chi) = (-1)^s \chi$  together implies that  $(-1)^r = (-1)^s$ , and hence both  $r$  and  $s$  are either even or odd.  $\square$

**Definition 2.5.27.** A permutation  $\sigma \in S_n$  is called *even* (respectively, *odd*) if  $\sigma$  can be written as a product of even (respectively, odd) number of transpositions in  $S_n$ .

Note that given a permutation  $\sigma \in S_n$ , if  $\sigma = \sigma_1 \circ \cdots \circ \sigma_r$ , where  $\sigma_1, \dots, \sigma_r$  are 2-cycles in  $S_n$ , then by Corollary 2.5.26 we see that  $\sigma$  is even if and only if  $(-1)^r = 1$ . Thus we have a well-defined map  $\text{sgn} : S_n \rightarrow \{1, -1\}$  given by sending  $\sigma \in S_n$  to  $(-1)^r$ , where  $r$  is a number of 2-cycles appearing in the decomposition of  $\sigma$  into a product of 2-cycles in  $S_n$ . In other words,

$$(2.5.28) \quad \text{sgn}(\sigma) = \begin{cases} -1, & \text{if } \sigma \text{ is odd,} \\ 1, & \text{if } \sigma \text{ is even,} \end{cases}$$

The number  $\text{sgn}(\sigma)$  is called the *signature* of the permutation  $\sigma \in S_n$ .

**Proposition 2.5.29.** An  $r$ -cycle  $\sigma \in S_n$  is even if and only if  $r$  is odd.

*Proof.* Let  $\sigma = (k_1 \ k_2 \ \cdots \ k_r)$  be an  $r$ -cycle in  $S_n$ . Then we can write it as a product  $\sigma = (k_1 \ k_2 \ \cdots \ k_r) = (k_1 \ k_r)(k_1 \ k_{r-1}) \cdots (k_1 \ k_2)$  of  $r - 1$  number of transpositions in  $S_n$ . Hence the result follows.  $\square$

**Exercise 2.5.30.** Express the following permutations as product of disjoint cycles, and then express them as a product of transpositions. Determine if they are even or odd permutations.

$$(i) \ \sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 3 & 8 & 5 & 6 & 4 & 7 & 1 \end{pmatrix} \in S_8.$$

*Answer:* Note that,

$$\begin{aligned} \sigma &= (1 \ 2 \ 3 \ 8) \circ (4 \ 5 \ 6) \\ &= (1 \ 8) \circ (1 \ 3) \circ (1 \ 2) \circ (4 \ 6) \circ (4 \ 5). \end{aligned}$$

Since  $\sigma$  is a product of 5 transpositions in  $S_8$ , we conclude that  $\sigma$  is odd.

$$(ii) \tau = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 1 & 4 & 2 & 3 & 6 \end{pmatrix} \in S_6.$$

$$(iii) \pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 2 & 4 & 6 & 1 & 3 & 7 & 5 \end{pmatrix} \in S_7.$$

**Exercise 2.5.31.** If  $\sigma \in S_5$  has order 3, show that  $\sigma$  is a 3-cycle. More generally, if  $\sigma \in S_n$  has order  $p > 0$ , a prime number, such that  $n < 2p$ , show that  $\sigma$  is a  $p$ -cycle in  $S_n$ .

**Proposition 2.5.32.** Let  $A_n = \{\sigma \in S_n : \sigma \text{ is even}\}$  be the set of all even permutations in  $S_n$ . Then  $A_n$  is a subgroup of  $S_n$ , known as the alternating group on  $J_n$ .

*Proof.* Since  $e = (1 \ 2) \circ (1 \ 2)$ , we see that  $e \in A_n$ . Thus  $A_n$  is a non-empty subset of  $S_n$ . Let  $\sigma, \tau \in A_n$  be arbitrary. Suppose that  $\tau = \tau_1 \circ \cdots \circ \tau_{2r}$ , where  $\tau_1, \dots, \tau_{2r}$  are transpositions in  $S_n$ . Since transpositions are elements of order 2 (see Proposition 2.5.14), they are self inverse in  $S_n$ . Now it follows from Exercise 2.1.8 (ii) that

$$\tau^{-1} = \tau_{2r} \circ \cdots \circ \tau_1.$$

Therefore,  $\tau^{-1}$  is also an even permutation. Since  $\sigma$  and  $\tau^{-1}$  are even, their product  $\sigma \circ \tau^{-1} \in A_n$ . Therefore,  $A_n$  is a subgroup of  $S_n$  by Lemma 2.2.8.  $\square$

**Remark 2.5.33.** Assume that  $n \geq 3$ . Note that, any transposition  $(i \ j) \in S_n$ , with  $i \neq 1$  and  $j \neq 1$ , can be written as

$$(i \ j) = (1 \ i) \circ (1 \ j) \circ (1 \ i).$$

Again  $(1 \ i) \circ (1 \ j) = (1 \ j \ i)$ . Since each element of  $A_n$  are product of even number of transpositions, using above two observations, one can write each element of  $A_n$  as product of 3 cycles in  $S_n$ .

**Exercise 2.5.34.** For all  $n \geq 3$ , show that  $A_n$  is generated by 3-cycles.

*Solution:* Note that any 3-cycle is an even permutation by Proposition 2.5.29, and hence is in  $A_n$ . Therefore, the subgroup of  $S_n$  generated by all 3-cycles is a subgroup of  $A_n$ . For the converse part, we show that any even permutation can be written as product of 3-cycles. Note that any element of  $A_n$  is a product of even number of 2-cycles in  $S_n$ . Let  $\sigma = (i \ j)$  and  $\tau = (k \ \ell)$  be two 2-cycles in  $S_n$ . If  $\sigma$  and  $\tau$  are not disjoint, then we may assume that  $j = k$ . Then  $\sigma \circ \tau = (i \ j)(j \ \ell) = (i \ j \ \ell)$  is a 3-cycle. If  $\sigma$  and  $\tau$  are disjoint, then

$$\begin{aligned} \sigma \circ \tau &= (i \ j)(k \ \ell) \\ &= (i \ j)(j \ k)(j \ k)(k \ \ell) \\ &= (i \ j \ k)(j \ k \ \ell), \end{aligned}$$

where the last equality is due to the first case. Hence the result follows.  $\square$

**Exercise 2.5.35.** Show that  $|A_n| = n!/2$ .

*Solution:* Let  $\{\sigma_1, \dots, \sigma_r\}$  and  $\{\tau_1, \dots, \tau_s\}$  be the set of all even permutations and the set of all odd permutations in  $S_n$ , respectively. Since  $r + s = n!$ , it suffices to show that  $r = s$ . Fix a transposition  $\pi \in S_n$ . Then  $\pi\sigma_1, \dots, \pi\sigma_r$  are all distinct (verify) odd permutations in  $S_n$ , and hence  $r \leq s$ . Similarly  $s \leq r$ , and hence  $r = s$ , as required.  $\square$

**Exercise 2.5.36.** Determine the groups  $A_3$  and  $A_4$ .

**Exercise 2.5.37.** Given  $\sigma, \tau \in S_n$ , show that  $[\sigma, \tau] := \sigma \circ \tau \circ \sigma^{-1} \circ \tau^{-1} \in A_n$ . The element  $[\sigma, \tau]$  is called the *commutator* of  $\sigma$  and  $\tau$  in  $S_n$ . Deduce that  $A_n$  is generated by  $\{[\sigma, \tau] : \sigma, \tau \in S_n\}$ , for all  $n \geq 3$ .

**Exercise 2.5.38.** Show that  $S_n$  is generated by  $\{(1 \ 2), (1 \ 2 \ \cdots \ n)\}$ , for all  $n \geq 3$ .

**Example 2.5.39** (Dihedral group  $D_n$ ). Consider a regular  $n$ -gon in the plane  $\mathbb{R}^2$  whose vertices are labelled as  $1, 2, 3, \dots, n$  in clockwise order. Let  $D_n$  be the set of all symmetries of this regular  $n$ -gon given by the following operations and their finite compositions:

$a :=$  The rotations about its centre through the angles  $2\pi/n$ , and

$b :=$  The reflections along the vertical straight line passing through the centre of the regular  $n$ -gon.

Note that  $\text{ord}(a) = n$ ,  $\text{ord}(b) = 2$  and that  $a^{n-1}b = ba$ . Therefore, the group generated by all such symmetries of the regular  $n$ -gon can be expressed in terms of generators and relations as

$$D_n := \langle a, b \mid \text{ord}(a) = n, \text{ord}(b) = 2, \text{ and } a^{n-1}b = ba \rangle.$$

This group is called the *dihedral group* of degree  $n$ . Note that  $D_n$  is a non-commutative finite group of order  $2n$  and its elements can be expressed as

$$D_n = \{e, a, a^2, a^3, \dots, a^{n-1}, b, ba, ba^2, ba^3, \dots, ba^{n-1}\}.$$

Note that each element of  $D_n$  is given by a bijection of the set  $J_n := \{1, 2, \dots, n\}$  onto itself, and hence is a permutation on  $J_n$ . However, not all permutations of the set  $J_n$  corresponds to a symmetry of a regular  $n$ -gon as described above (see Exercise 2.5.40 below). We can define a binary operation on  $D_n$  by composition of bijective maps. Then it is easy to check using Lemma 2.2.8 that  $D_n$  is a subgroup of  $S_n$ . The group  $D_n$  is called the *Dihedral group* of degree  $n$ . It is a finite group of order  $2n$  which is non-commutative for  $n \geq 3$ .

**Exercise 2.5.40.** Show that  $D_3 = S_3$ , and  $D_n$  is a proper subgroup of  $S_n$ , for all  $n \geq 4$ .

**Exercise 2.5.41.** Let  $G$  be the subgroup of  $S_4$  generated by the cycles

$$a := (1 \ 2 \ 3 \ 4) \text{ and } b := (2 \ 4)$$

in  $S_4$ . Show that  $G$  is a dihedral group of degree 4.

## 2.6 Group homomorphism

A group homomorphism is a map from a group  $G$  into another group  $H$  that respects the binary operations on them. Here is a formal definition.

**Definition 2.6.1.** Let  $G$  and  $H$  be two groups. A *group homomorphism* from  $(G, *)$  into  $(H, \star)$  is a map  $f : G \rightarrow H$  satisfying  $f(a * b) = f(a) \star f(b)$ , for all  $a, b \in G$ .

**Example 2.6.2.** (i) For any group  $G$ , the constant map  $c_e : G \rightarrow G$ , which sends all points of  $G$  to the neutral element  $e \in G$ , is a group homomorphism, called the *trivial group homomorphism* of  $G$ .

(ii) Let  $H$  be a subgroup of a group  $G$ . Then the set theoretic inclusion map  $H \hookrightarrow G$  is a group homomorphism. In particular, for any group  $G$ , the identity map

$$\text{Id}_G : G \rightarrow G, \ a \mapsto a$$

is a group homomorphism.

(iii) Fix an integer  $m$ , and define a function

$$\varphi_m : \mathbb{Z} \longrightarrow \mathbb{Z}, \ n \longmapsto mn, \ \forall n \in \mathbb{Z}.$$

Then  $\varphi_m(n_1 + n_2) = m(n_1 + n_2) = mn_1 + mn_2 = \varphi_m(n_1) + \varphi_m(n_2)$ , for all  $n_1, n_2 \in \mathbb{Z}$ . Therefore,  $\varphi_m$  is a group homomorphism. Note that,  $\varphi_m$  is always injective, and it is surjective only for  $m \in \{1, -1\}$ .

(iv) Let  $\mathbb{R}^* := \mathbb{R} \setminus \{0\}$ , and consider the exponential map

$$f : \mathbb{R} \longrightarrow \mathbb{R}^*, \quad x \longmapsto e^x, \quad \forall x \in \mathbb{R}.$$

Since  $f(a + b) = e^{a+b} = e^a \cdot e^b = f(a) \cdot f(b)$ , for all  $a, b \in \mathbb{R}$ , the map  $f$  is a group homomorphism from  $(\mathbb{R}, +)$  into  $(\mathbb{R}^*, \cdot)$ . Verify that  $f$  is injective.

(v) The map  $f : \mathbb{R} \rightarrow S^1 := \{z \in \mathbb{C}^* : |z| = 1\}$  defined by  $f(t) = e^{2\pi it}$ ,  $\forall t \in \mathbb{R}$  is a surjective group homomorphism. Is it injective?

(vi) Let

$$\phi : \mathbb{R} \longrightarrow \mathrm{SL}_2(\mathbb{R}), \quad a \longmapsto \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}, \quad \forall a \in \mathbb{R}.$$

Verify that  $\phi$  is an injective group homomorphism from the additive group  $\mathbb{R}$  into the multiplicative group  $\mathrm{SL}_2(\mathbb{R})$ .

(vii) Fix an integer  $n \geq 2$ , and consider the map

$$\psi : \mathbb{Z} \longrightarrow \mathbb{Z}_n, \quad a \longmapsto [a], \quad \forall a \in \mathbb{Z}.$$

Verify that  $\psi$  is a surjective group homomorphism.

(viii) Fix a prime number  $p > 0$ , and let  $\mathbf{F} : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$  be the map defined by  $\mathbf{F}(a) = a^p$ , for all  $a \in \mathbb{Z}_p$ . Since any multiple of  $p$  is 0 in  $\mathbb{Z}_p$ , using binomial expansion we have

$$\mathbf{F}(a + b) = (a + b)^p = \sum_{j=0}^p \binom{p}{j} a^{p-j} b^j = a^p + b^p.$$

Therefore,  $\mathbf{F}$  is a group homomorphism, known as the *Frobenius endomorphism*.

(ix) Fix an integer  $n \geq 1$ , and let  $f : \mathrm{GL}_n(\mathbb{R}) \rightarrow \mathbb{R}^*$  be the map defined by

$$f(A) = \det(A), \quad \forall A \in \mathrm{GL}_n(\mathbb{R}).$$

Verify that  $f$  is a group homomorphism.

(x) Let  $m, n > 1$  be integers such that  $n \mid m$  in  $\mathbb{Z}$ . Verify that the map  $\varphi : \mathbb{Z}_m \rightarrow \mathbb{Z}_n$  defined by sending  $[a] \in \mathbb{Z}_m$  to  $[a] \in \mathbb{Z}_n$  is a well-defined map that is a group homomorphism.

(xi) Let  $G$  be a group. For each  $a \in G$ , the map  $\varphi_a : G \rightarrow G$  defined by  $\varphi_a(b) = aba^{-1}$ ,  $\forall b \in G$ , is a group homomorphism.

**Exercise 2.6.3.** For each integer  $n \geq 1$ , let  $J_n := \{k \in \mathbb{Z} : 1 \leq k \leq n\}$ . For each  $\sigma \in S_n$ , consider the map  $\tilde{\sigma} : J_{n+1} \rightarrow J_{n+1}$  defined by

$$\tilde{\sigma}(k) = \begin{cases} \sigma(k), & \text{if } 1 \leq k \leq n, \\ n+1, & \text{if } k = n+1. \end{cases}$$

Note that,  $\tilde{\sigma}$  is a bijective map, and hence is an element of  $S_{n+1}$ . Show that the map

$$f : S_n \rightarrow S_{n+1}, \quad \sigma \mapsto \tilde{\sigma},$$

is an injective group homomorphism. Thus, we can identify  $S_n$  as a subgroup of  $S_{n+1}$ .



**Lemma 2.6.4.** Let  $n \geq 2$  be an integer. Then the map  $\text{sgn} : S_n \rightarrow \{1, -1\}$  defined by sending  $\sigma \in S_n$  to

$$\text{sgn}(\sigma) = \begin{cases} -1, & \text{if } \sigma \text{ is odd,} \\ 1, & \text{if } \sigma \text{ is even,} \end{cases}$$

is a group homomorphism, called the *signature homomorphism for  $S_n$* .

*Proof.* Let  $\sigma, \tau \in S_n$  be arbitrary. Let  $\sigma = \sigma_1 \circ \cdots \circ \sigma_r$  and  $\tau = \tau_1 \circ \cdots \circ \tau_s$ , where  $\sigma_i, \tau_j$  are all 2-cycles in  $S_n$ . Then  $\sigma \circ \tau = \sigma_1 \circ \cdots \circ \sigma_r \circ \tau_1 \circ \cdots \circ \tau_s$ , and hence  $\text{sgn}(\sigma \circ \tau) = (-1)^{r+s} = (-1)^r (-1)^s = \text{sgn}(\sigma) \text{sgn}(\tau)$ .  $\square$

**Proposition 2.6.5.** Let  $f : G \rightarrow H$  be a group homomorphism. Let  $e_G \in G$  and  $e_H \in H$  be the neutral elements of  $G$  and  $H$ , respectively. Then we have the following.

- (i)  $f(e_G) = e_H$ .
- (ii)  $f(a^{-1}) = (f(a))^{-1}$ , for all  $a \in G$ .
- (iii) If  $\text{ord}(a) < \infty$ , then  $\text{ord}(f(a)) \mid \text{ord}(a)$ .

*Proof.* (i) Since  $f(e_G)f(e_G) = f(e_G \cdot e_G) = f(e_G) = f(e_G) \cdot e_H$ , applying cancellation law we have  $f(e_G) = e_H$ . The second statement follows immediately.

(ii) Since  $f$  is a group homomorphism, for any  $a \in G$ , we have

$$\begin{aligned} f(a)f(a^{-1}) &= f(a \cdot a^{-1}) = f(e_G) = e_H \\ \text{and } f(a^{-1})f(a) &= f(a^{-1} \cdot a) = f(e_G) = e_H, \end{aligned}$$

and hence  $f(a^{-1}) = (f(a))^{-1}$ .

- (iii) Let  $n := \text{ord}(a) < \infty$ . Since  $f(a)^n = f(a^n) = f(e_G) = e_H$ , it follows from Exercise 2.2.32 (i) that  $\text{ord}(f(a)) \mid n$ .

$\square$

**Exercise 2.6.6.** Let  $G$  and  $H$  be two groups. Show that there is a unique constant group homomorphism from  $G$  to  $H$ .

**Proposition 2.6.7.** Let  $f : G \rightarrow H$  be a group homomorphism.

- (i) For any subgroup  $G'$  of  $G$ , its image  $f(G') := \{f(a) : a \in G'\}$  is a subgroup of  $H$ . Moreover, if  $G'$  is commutative, so is  $f(G')$ .
- (ii) For any subgroup  $H'$  of  $H$ , its inverse image  $f^{-1}(H') := \{a \in G : f(a) \in H'\}$  is a subgroup of  $G$ .

*Proof.* (i) Clearly,  $f(G') \neq \emptyset$  as  $e \in G'$ . For  $h_1, h_2 \in f(G')$ , we have  $h_1 = f(a_1)$  and  $h_2 = f(a_2)$ , for some  $a_1, a_2 \in G'$ . Since  $a_1 a_2^{-1} \in G'$ , we have  $h_1 h_2^{-1} = f(a_1) f(a_2)^{-1} = f(a_1 a_2^{-1}) \in f(G')$ . If  $G'$  is commutative, we have  $f(a)f(b) = f(ab) = f(ba) = f(b)f(a)$ , for all  $a, b \in G'$ . Hence the result follows.

- (ii) Let  $e_G \in G$  and  $e_H \in H$  be the neutral elements of  $G$  and  $H$ , respectively. Since  $f(e_G) = e_H$  by Proposition 2.6.5 (i), we have  $e_G \in f^{-1}(H')$ . Since  $H'$  is a subgroup of  $H$ , for any  $a, b \in f^{-1}(H')$  we have  $f(ab^{-1}) = f(a)f(b)^{-1} \in H'$ , and hence  $ab^{-1} \in f^{-1}(H')$ . Thus  $f^{-1}(H')$  is a subgroup of  $G$ .

$\square$

**Proposition 2.6.8.** Composition of group homomorphisms is a group homomorphism.

*Proof.* Let  $f : G_1 \rightarrow G_2$  and  $g : G_2 \rightarrow G_3$  be two group homomorphisms. Since  $(g \circ f)(ab) = g(f(ab)) = g(f(a)f(b)) = g(f(a))g(f(b)) = (g \circ f)(a)(g \circ f)(b)$ , for all  $a, b \in G_1$ , the result follows.  $\square$

Given any two groups  $G$  and  $H$ , we denote by  $\text{Hom}(G, H)$  the set of all group homomorphisms from  $G$  into  $H$ .

**Exercise 2.6.9.** Let  $G$  and  $H$  be two groups. Show that the projection maps  $\pi_G : G \times H \rightarrow G$  and  $\pi_H : G \times H \rightarrow H$  defined by

$$\pi_G(a, b) = a \quad \text{and} \quad \pi_H(a, b) = b, \quad \forall (a, b) \in G \times H,$$

are surjective group homomorphisms.

**Proposition 2.6.10.** Let  $G, H$  and  $K$  be groups. Then there is a natural bijective map from  $\text{Hom}(G, H \times K)$  onto  $\text{Hom}(G, H) \times \text{Hom}(G, K)$ .

*Proof.* Let  $\pi_H : H \times K \rightarrow H$  and  $\pi_K : H \times K \rightarrow K$  be the projection maps onto the first and the second factors, respectively (see Exercise 2.6.9). Since both  $\pi_H$  and  $\pi_K$  are group homomorphisms, given any group homomorphism  $f : G \rightarrow H \times K$ , we have  $\pi_H \circ f \in \text{Hom}(G, H)$  and  $\pi_K \circ f \in \text{Hom}(G, K)$  by Proposition 2.9.20. Thus we get a map  $\Phi : \text{Hom}(G, H \times K) \rightarrow \text{Hom}(G, H) \times \text{Hom}(G, K)$  defined by

$$\Phi(f) = (\pi_H \circ f, \pi_K \circ f), \quad \forall f \in \text{Hom}(G, H \times K).$$

To show that  $\Phi$  is surjective, given  $f \in \text{Hom}(G, H)$  and  $g \in \text{Hom}(G, K)$ , let  $h : G \rightarrow H \times K$  be the map defined by

$$h(a) = (f(a), g(a)), \quad \forall a \in G.$$

Since for given any  $a, b \in G$ , we have

$$\begin{aligned} h(ab) &= (f(ab), g(ab)) = (f(a)f(b), g(a)g(b)) \\ &= (f(a), g(a))(f(b), g(b)) \\ &= h(a)h(b), \end{aligned}$$

we see that  $h \in \text{Hom}(G, H \times K)$ . Clearly  $\Phi(h) = (\pi_H \circ h, \pi_K \circ h) = (f, g)$ . Therefore,  $\Phi$  is surjective. To show that  $\Phi$  is injective, note that given any  $f \in \text{Hom}(G, H \times K)$ , we have

$$f(a) = ((\pi_H \circ f)(a), (\pi_K \circ f)(a)), \quad \forall a \in G.$$

Therefore, if  $\Phi(f) = \Phi(g)$  for some  $f, g \in \text{Hom}(G, H \times K)$ , then the conditions  $\pi_H \circ f = \pi_H \circ g$  and  $\pi_K \circ f = \pi_K \circ g$  together forces that  $f = g$ . This completes the proof.  $\square$

**Definition 2.6.11.** Let  $f : G \rightarrow H$  be a group homomorphism. We say that

- (i)  $f$  is *trivial* if  $f(a) = e_H$ , for all  $a \in G$ .
- (ii)  $f$  is a *monomorphism* if  $f$  is injective (c.f. Proposition 2.6.24),
- (iii)  $f$  is an *epimorphism* if  $f$  is surjective (c.f. Proposition 2.6.25), and
- (iv)  $f$  is an *isomorphism* if  $f$  is bijective. In that case, we say that  $G$  is *isomorphic to*  $H$ , and express it as  $G \cong H$ .

**Lemma 2.6.12.** Being isomorphic groups is an equivalence relation.

*Proof.* Given any group  $G$ , the identity map  $\text{Id}_G : G \rightarrow G$  given by  $\text{Id}_G(a) = a$ , for all  $a \in G$ , is an isomorphism of groups. Therefore, being isomorphic is a reflexive relation. If  $f : G \rightarrow H$  is

an isomorphism of groups, then its inverse map  $f^{-1} : H \rightarrow G$  is also a group homomorphism, and hence is an isomorphism because it is bijective. Therefore, being isomorphic groups is a symmetric relation. If  $f : G \rightarrow H$  and  $g : H \rightarrow K$  be isomorphism of groups. Then the composite map  $g \circ f : G \rightarrow K$  is a group homomorphism, which is an isomorphism of groups. Therefore, being isomorphic groups is a transitive relation. Hence the result follows.  $\square$

**Proposition 2.6.13.** *Given a group  $G$ , the set  $\text{Aut}(G)$  consisting of all group isomorphisms from  $G$  onto itself is a group with respect to the binary operation given by composition of maps; the group  $\text{Aut}(G)$  is known as the **automorphism group of  $G$** .*

*Proof.* Since composition of two bijective group homomorphisms is bijective and a group homomorphism, we see that the map

$$G \times G \rightarrow G, (f, g) \mapsto f \circ g,$$

is a binary operation on  $\text{Aut}(G)$ . Clearly composition of maps is associative. The identity map  $\text{Id}_G : G \rightarrow G$  plays the role of a neutral element in a group. Given  $f \in \text{Aut}(G)$ , its inverse  $f^{-1} : G \rightarrow G$  is again a group homomorphism. Indeed, given  $a, b \in G$  there exists unique  $x, y \in G$  such that  $f(x) = a$  and  $f(y) = b$ . Then we have  $f^{-1}(ab) = f^{-1}(f(x)f(y)) = f^{-1}(f(xy)) = xy = f^{-1}(a)f^{-1}(b)$ , and hence  $f^{-1} \in \text{Aut}(G)$ . This proves that  $\text{Aut}(G)$  is a group.  $\square$

**Example 2.6.14.** The complex conjugation map  $z \mapsto \bar{z}$  from the additive group  $\mathbb{C}$  into itself is an automorphism of  $\mathbb{C}$ .

**Exercise 2.6.15.** Show that  $\text{Aut}(K_4)$  is isomorphic to  $S_3$ . (Hint: Note that  $K_4 = \{e, a, b, c\}$ , where  $a^2 = b^2 = c^2 = e$  and  $ab = ba = c, bc = cb = a, ac = ca = b$ . If  $f \in \text{Aut}(K_4)$ , then  $f(e) = e$  and hence  $f|_{\{a, b, c\}}$  is a bijection of the subset  $\{a, b, c\} \subset K_4$  onto itself, producing an element of  $S_3$ . Thus we get a map  $\varphi : \text{Aut}(K_4) \rightarrow S_3$ . Verify that  $\varphi$  is a group isomorphism.)

**Definition 2.6.16.** The *kernel* of a group homomorphism  $f : G \rightarrow H$  is the subset

$$\text{Ker}(f) := \{a \in G : f(a) = e_H\} \subseteq G.$$

Since  $f(e_G) = e_H$  by Proposition 2.6.5 (i), we have  $e_G \in \text{Ker}(f)$ . Therefore,  $\text{Ker}(f)$  is a non-empty subset of  $G$ . Given any two elements  $a, b \in \text{Ker}(f)$  we have  $f(ab^{-1}) = f(a)f(b^{-1}) = f(a)f(b)^{-1} = e_H \cdot e_H^{-1} = e_H$ . Therefore,  $\text{Ker}(f)$  is a subgroup of  $G$ .

**Example 2.6.17.** (i) Fix an integer  $n$  and consider the homomorphism

$$f : \mathbb{Z} \rightarrow \mathbb{Z}_n, a \mapsto [a].$$

Then  $\text{Ker}(f) = \{a \in \mathbb{Z} : n \text{ divides } a\} = n\mathbb{Z}$ .

(ii) Let  $S^1 := \{z \in \mathbb{C} : |z| = 1\}$ . Consider the homomorphism

$$f : \mathbb{R} \rightarrow S^1, t \mapsto e^{2\pi\sqrt{-1}t}.$$

Then  $\text{Ker}(f) = \{t \in \mathbb{R} : e^{2\pi\sqrt{-1}t} = 1\} = \mathbb{Z}$ .

The following lemma shows that the kernel of a group homomorphism can be uniquely determined purely using its universal property. Interesting fact to note is that this description of kernel of a group homomorphism use only arrows and not any points.

**Proposition 2.6.18** (Universal Property of Kernel). *Let  $f : G \rightarrow H$  be a group homomorphism. Then there is a unique subgroup  $K$  of  $G$  satisfying the following properties.*

(K1)  $f \circ \iota_K$  is trivial, where  $\iota_K : K \hookrightarrow G$  is the inclusion map, and

(K2) given any group homomorphism  $\phi : G' \rightarrow G$  with  $f \circ \phi$  trivial, there is a unique group homomorphism  $\psi : G' \rightarrow K$  such that  $\iota_K \circ \psi = \phi$ .

$$(2.6.19) \quad \begin{array}{ccccc} & & G' & & \\ & \swarrow \exists! \psi & \downarrow \phi & \searrow f \circ \phi = e_H & \\ K & \xrightarrow{\iota_K} & G & \xrightarrow{f} & H \end{array}$$

*Proof.* We first show the uniqueness of  $K$ . Let  $\iota_{K'} : K' \hookrightarrow G$  be any subgroup of  $G$  satisfying (K1) and (K2). Since the homomorphism  $f \circ \iota_{K'}$  is trivial, applying (K2) for  $K$  we have a unique group homomorphism  $\eta : K' \rightarrow K$  such that  $\iota_{K'} = \iota_K \circ \eta$ . Similarly replacing  $(K, \iota_K)$  with  $(K', \iota_{K'})$ , and  $(G', \phi)$  with  $(K, \iota_K)$  in the above diagram (2.6.19), we get a unique group homomorphism  $\eta' : K \rightarrow K'$  such that  $\iota_K = \iota_{K'} \circ \eta'$ . Now replace  $(G', \phi)$  with  $(K, \iota_K)$  in the above diagram 2.6.19. Since both the group homomorphisms  $\text{Id}_K : K \rightarrow K$  and  $\eta \circ \eta' : K \rightarrow K$  satisfies  $\iota_K \circ (\eta \circ \eta') = \iota_K$  and  $\iota_K \circ \text{Id}_K = \iota_K$ , by uniqueness assumption in (K2), we have  $\eta \circ \eta' = \text{Id}_K$ . Similarly, we have  $\eta' \circ \eta = \text{Id}_{K'}$ . Therefore, both  $\eta' : K \rightarrow K'$  and  $\eta : K' \rightarrow K$  are isomorphisms. Since both  $\iota_K : K \hookrightarrow G$  and  $\iota_{K'} : K' \hookrightarrow G$  are inclusion maps, and  $\iota_K \circ \eta' = \iota_{K'}$ , we must have  $\eta'$  is an inclusion map, and hence  $K \subseteq K'$ . Similarly, we have  $K' \subseteq K$ , and hence  $K = K'$ .

To prove existence, take  $K = \text{Ker}(f)$  and  $\iota_K : K \hookrightarrow G$  the inclusion map. Clearly,  $f \circ \iota_K$  is trivial. For any group homomorphism  $\phi : G' \rightarrow G$  with  $f \circ \phi$  trivial, we have  $\phi(a) \in K$ , for all  $a \in G'$ . Thus the image of  $\phi$  lands inside  $K$  and hence we have a group homomorphism

$$\psi : G' \rightarrow K, \quad a \mapsto \phi(a)$$

such that  $\iota_K \circ \psi = \phi$  as required.  $\square$

**Proposition 2.6.20.** A group homomorphism  $f : G \rightarrow H$  is injective if and only if  $\text{Ker}(f)$  is trivial.

*Proof.* If  $\text{Ker}(f) \neq \{e\}$ , clearly  $f$  is not injective. Conversely, suppose that  $\text{Ker}(f) = \{e\}$ . If  $f(a) = f(b)$ , for some  $a, b \in G$  with  $a \neq b$ , then  $ab^{-1} \neq e$  and  $f(ab^{-1}) = f(a)f(b^{-1}) = f(a)f(b)^{-1} = e_H$ , which contradicts our assumption that  $\text{Ker}(f) = \{e\}$ . This completes the proof.  $\square$

**Proposition 2.6.21.** Any infinite cyclic group is isomorphic to  $\mathbb{Z}$ .

*Proof.* Let  $G = \langle a \rangle$  be an infinite cyclic group. Define a map  $f : \mathbb{Z} \rightarrow G$  by  $f(n) = a^n$ , for all  $n \in \mathbb{Z}$ . Since

$$f(n+m) = a^{n+m} = a^n a^m = f(n)f(m), \quad \forall m, n \in \mathbb{Z},$$

the map  $f$  is a group homomorphism. Since  $G$  is infinite, we have  $a^n \neq e$ ,  $\forall n \in \mathbb{Z} \setminus \{0\}$ . Therefore,  $\text{Ker}(f) = \{e\}$ , and so  $f$  is injective. Clearly  $f$  is surjective, and hence is an isomorphism.  $\square$

**Proposition 2.6.22.** Let  $G$  be a cyclic group generated by  $a \in G$ . A homomorphism  $f : G \rightarrow G$  is an automorphism of  $G$  if and only if  $f(a)$  is a generator of  $G$ .

*Proof.* Let  $f : G \rightarrow G$  be an automorphism of  $G$ . Let  $b = f(a)$ . Let  $x \in G$  be arbitrary. Since  $f$  is surjective, there exists  $y \in G$  such that  $f(y) = x$ . Since  $G = \langle a \rangle$ , we have  $y = a^n$ , for some  $n \in \mathbb{Z}$ . Then  $x = f(y) = f(a^n) = [f(a)]^n = b^n \in \langle b \rangle$ . This shows that  $G = \langle b \rangle$ , and hence  $b$  is a generator of  $G$ . Conversely if  $f : G \rightarrow G$  is a homomorphism such that  $f(a)$  generates  $G$ , then  $f$  is surjective. If  $|G|$  is finite, we must have  $f$  is bijective. If  $G$  is not finite, then  $G$  has only two generators, namely  $a$  and  $a^{-1}$  by Proposition 2.3.15, and hence  $f$  must be either  $\text{Id}_G$  or the map given by sending  $b \in G$  to  $b^{-1}$ . In both cases,  $f$  is injective, and hence is in  $\text{Aut}(G)$ .  $\square$

**Theorem 2.6.23 (Cayley).** Every group is a subgroup of a symmetric group.

*Proof.* Let  $G$  be a group. Let  $S(G)$  be the symmetric group on  $G$ ; its elements are all bijective maps from  $G$  onto itself and the group operation is given by composition of bijective maps. Define a map

$$\varphi : G \longrightarrow S(G)$$

by sending an element  $a \in G$  to the map

$$\varphi_a : G \rightarrow G, \quad g \mapsto ag,$$

which is bijective (verify!), and hence is an element of  $S(G)$ . Then given any  $g \in G$  we have

$$\begin{aligned} \varphi(ab)(g) &= \varphi_{ab}(g) \\ &= (ab)g = a(bg) \\ &= (\varphi_a \circ \varphi_b)(g) \\ &= (\varphi(a) \circ \varphi(b))(g), \end{aligned}$$

and hence  $\varphi$  is a group homomorphism. Note that  $\varphi_a = \text{Id}_G$  if and only if  $a = e$  in  $G$  (verify!). Therefore,  $\varphi$  is an injective group homomorphism, and hence we can identify  $G$  with the subgroup  $\varphi(G)$  of the symmetric group  $S(G)$ .  $\square$

We end this section with the following two results which justify the terminologies introduced in Definition 2.6.11 in the light of category theory.

**Proposition 2.6.24.** *Let  $f : G \rightarrow H$  be a group homomorphism. Then the following are equivalent.*

- (i)  $f$  is injective.
- (ii) Given a group  $T$  and group homomorphisms  $\phi, \psi : T \rightarrow G$  with  $f \circ \phi = f \circ \psi$ , we have  $\phi = \psi$ .  
In other words,  $f$  is a **monomorphism in the category of groups**.
- (iii) Given a group  $T$  and a group homomorphism  $\phi : T \rightarrow G$  with  $f \circ \phi$  trivial, we have  $\phi$  is trivial.

*Proof.* (i)  $\Rightarrow$  (ii) is Clear. To show (ii)  $\Rightarrow$  (iii), take  $\psi : T \rightarrow G$  to be the trivial group homomorphism. Then both  $f \circ \phi$  and  $f \circ \psi$  are trivial, and hence  $\phi$  is trivial by (ii). To show (iii)  $\Rightarrow$  (i), take  $T = \text{Ker}(f)$  and  $\phi : T \rightarrow G$  the inclusion map of  $\text{Ker}(f)$  into  $G$ . Then  $f \circ \phi$  is trivial, and hence the inclusion map  $\phi : \text{Ker}(f) \hookrightarrow G$  is a trivial group homomorphism by (iii). This forces  $\text{Ker}(f) = \{e\}$ , and hence  $f$  is injective.  $\square$

**Proposition 2.6.25.** *Let  $f : G \rightarrow H$  be a group homomorphism. Then the following are equivalent.*

- (i)  $f$  is surjective.
- (ii) Given a group  $T$  and group homomorphisms  $\phi, \psi : H \rightarrow T$  with  $\phi \circ f = \psi \circ f$ , we have  $\phi = \psi$ .  
In other words,  $f$  is an **epimorphism in the category of groups**.
- (iii) Given a group  $T$  and a group homomorphism  $\phi : H \rightarrow T$  with  $\phi \circ f$  trivial, we have  $\phi$  is trivial.

*Proof.* (i)  $\Rightarrow$  (ii): Let  $\phi, \psi : H \rightarrow T$  be group homomorphisms with  $\phi \circ f = \psi \circ f$ . Since  $f$  is surjective, given  $h \in H$  there exists  $g \in G$  such that  $f(g) = h$ . Then  $(\phi \circ f)(g) = (\psi \circ f)(g)$  gives  $\phi(h) = \psi(h)$ . Since  $h \in H$  is arbitrary, we have  $\phi = \psi$ .

(ii)  $\Rightarrow$  (iii): Take  $\psi : H \rightarrow T$  to be the trivial group homomorphism.

(iii)  $\Rightarrow$  (i): We use the notion of coset of a subgroup. See Proposition 2.7.19 for a proof.  $\square$

## 2.7 Notion of Quotient & Cosets

Let  $G$  be a group, and  $H$  a subgroup of  $G$ . In this section we introduce the notion of a quotient group of  $G$  by  $H$  and prove its uniqueness. In the process of construction of quotient, we identify a class of subsets of  $G$ , known as *cosets* of  $H$  in  $G$ , and discuss their basic properties with some applications. An explicit construction of quotient group will appear in the next section.

**Definition 2.7.1 (Quotient Group).** Let  $H$  be a subgroup of a group  $G$ . The *quotient* of  $G$  by  $H$  is a pair  $(Q, \pi)$ , where  $Q$  is a group and  $\pi : G \rightarrow Q$  is a surjective group homomorphism such that

(QG1)  $\pi(h) = e_Q$ , the neutral element of  $Q$ , for all  $h \in H$ , and

(QG2) **Universal property of quotient:** given a group  $T$  and a group homomorphism  $t : G \rightarrow T$  satisfying  $H \subseteq \text{Ker}(t)$ , there exists a **unique** group homomorphism  $\tilde{t} : Q \rightarrow T$  such that  $\tilde{t} \circ \pi = t$ ; i.e., the following diagram commutes.

$$(2.7.2) \quad \begin{array}{ccc} G & \xrightarrow{t} & T \\ \pi \downarrow & \searrow \tilde{t} & \\ Q & & \end{array}$$

Interesting point is that, without knowing existence of such a pair  $(Q, \pi)$ , it follows immediately from the properties (QG1) and (QG2) that such a pair  $(Q, \pi)$ , if it exists, must be unique up to a unique isomorphism of groups in the following sense.

**Proposition 2.7.3 (Uniqueness of Quotient).** With the above notations, if  $(Q, \pi)$  and  $(Q', \pi')$  are two quotients of  $G$  by  $H$ , then there exists a **unique** group isomorphism  $\varphi : Q \rightarrow Q'$  such that  $\varphi \circ \pi = \pi'$ .

*Proof.* Taking  $(T, t) = (Q', \pi')$  by universal property of quotient  $(Q, \pi)$  we have a unique group homomorphism  $\tilde{\pi}' : Q \rightarrow Q'$  such that  $\tilde{\pi}' \circ \pi = \pi'$ . Similarly, taking  $(T, t) = (Q, \pi)$  by universal property of quotient  $(Q', \pi')$  we have a unique group homomorphism  $\tilde{\pi} : Q' \rightarrow Q$  such that  $\tilde{\pi} \circ \pi' = \pi$ . Since both  $\tilde{\pi} \circ \tilde{\pi}'$  and  $\text{Id}_Q$  are group homomorphisms from  $Q$  into itself making the following diagram commutative,

$$\begin{array}{ccccc} & & G & & \\ & \swarrow \pi & \downarrow \pi' & \searrow \pi & \\ Q & \xrightarrow{\tilde{\pi}'} & Q' & \xrightarrow{\tilde{\pi}} & Q \\ & \searrow & \text{Id}_Q & \swarrow & \end{array}$$

it follows that  $\tilde{\pi} \circ \tilde{\pi}' = \text{Id}_Q$ . Similarly  $\tilde{\pi}' \circ \tilde{\pi} = \text{Id}_{Q'}$ . Therefore,  $\tilde{\pi}' : Q \rightarrow Q'$  is the unique group isomorphism such that  $\tilde{\pi}' \circ \pi = \pi'$ . This completes the proof.  $\square$

Now question is about existence of quotient. We shall see shortly that we need to impose an additional hypothesis on  $H$  (namely  $H$  should be a normal subgroup of  $G$ ) for existence of quotient. The condition (QG1) says that  $\pi(H) = \{e_Q\}$ . Since  $\pi : G \rightarrow Q$  is a group homomorphism by assumption, given any two elements  $a, b \in G$  with  $a^{-1}b \in H$  we have  $\pi(a^{-1}b) = e_Q$ , and hence  $\pi(a) = \pi(b)$ . In other words, two elements  $a, b \in G$  are in the same fiber of the map  $\pi : G \rightarrow Q$  if  $a^{-1}b \in H$ . Since the set of all fibers of any set map  $f : G \rightarrow Q$  gives a partition of  $G$ , and hence an equivalence relation on  $G$ , the condition (QG1) suggests us to define a relation  $\rho_L$  on  $G$  by setting

$$(a, b) \in \rho_L \quad \text{if} \quad a^{-1}b \in H.$$

It is easy to check that  $\rho_L$  is an equivalence relation on  $G$  (verify!). The  $\rho_L$ -equivalence class of an element  $a \in G$  is the subset

$$[a]_{\rho_L} := \{b \in G : a^{-1}b \in H\} = \{ah : h \in H\},$$

which we denote by  $aH$ ; the subset  $aH$  is called the **left coset** of  $H$  in  $G$  represented by  $a$ . Note that (verify!), given  $a, b \in G$ ,

- (i) either  $aH \cap bH = \emptyset$  or  $aH = bH$ ,
- (ii)  $aH = bH$  if and only if  $a^{-1}b \in H$ , and
- (iii)  $G = \bigcup_{a \in G} aH$ .

**Proposition 2.7.4.** *For each  $a \in G$ , the map  $\varphi_a : H \rightarrow aH$  defined by  $\varphi_a(h) = ah$ , for all  $h \in H$ , is bijective. Consequently,  $|aH| = |bH|$ , for all  $a, b \in H$ .*

*Proof.* Since every element of  $aH$  is of the form  $ah$ , for some  $h \in H$ , we see that  $\varphi_a(h) = ah$ , and hence  $\varphi_a$  is surjective. Since  $ah = ah'$  implies that  $h = (a^{-1}a)h = a^{-1}(ah) = a^{-1}(ah') = (a^{-1}a)h' = h'$ , we see that  $\varphi_a$  is injective. Therefore,  $\varphi_a$  is bijective. Thus, both  $H$  and  $aH$  have the same cardinality.  $\square$

Let  $G/H = \{aH : a \in G\}$  be the set of all distinct left cosets of  $H$  in  $G$ .

**Theorem 2.7.5 (Lagrange's Theorem).** *Let  $G$  be a finite group, and  $H$  a subgroup of  $G$ . Then  $|H|$  divides  $|G|$ .*

*Proof.* Since  $\rho_L$  is an equivalence relation on  $G$ , it follows from Proposition 2.1.31 that  $G$  is a disjoint union of distinct left cosets of  $H$  in  $G$ . Since  $G$  is finite, there can be at most finitely many distinct left cosets of  $H$  in  $G$ . Since  $|aH| = |bH|$ , for all  $a, b \in G$  (see Proposition 2.7.4), it follows that

$$|G| = |G/H| \cdot |H|,$$

where  $|G/H|$  is the cardinality of the set  $G/H$ , i.e., the number of distinct left cosets of  $H$  in  $G$ . This completes the proof.  $\square$

**Exercise 2.7.6.** Let  $G$  be a finite group of order  $mn$  having subgroups  $H$  and  $K$  of orders  $m$  and  $n$ , respectively. If  $\gcd(m, n) = 1$  show that  $HK := \{hk \in G : h \in H, k \in K\}$  is a group.

**Corollary 2.7.7.** *Let  $G$  be a finite group of order  $n$ . Then for any  $a \in G$ ,  $\text{ord}(a)$  divides  $n$ . In particular,  $a^n = e$ ,  $\forall a \in G$ .*

*Proof.* Let  $H$  be the cyclic subgroup of  $G$  generated by  $a$ . Since  $G$  is a finite group, so is  $H$ . Then by Lagrange's theorem 2.7.5,  $|H|$  divides  $|G| = n$ . Since  $|H| = \text{ord}(a)$ , the result follows. To see the second part, note that if  $\text{ord}(a) = k$ , then  $n = km$ , for some  $m \in \mathbb{N}$ , and so  $a^n = (a^k)^m = e^m = e$ .  $\square$

**Exercise 2.7.8.** Let  $G$  be a finite group of order  $n$ . Let  $k \in \mathbb{N}$  be such that  $\gcd(n, k) = 1$ . Show that the map  $f : G \rightarrow G$  defined by  $f(a) = a^k$ ,  $\forall a \in G$ , is injective, and hence is bijective.

**Corollary 2.7.9.** *Any group of prime order is cyclic.*

*Proof.* Let  $G$  be a finite group of order  $p$ , where  $p$  is a prime number. If  $p = 2$ , then clearly  $G$  is cyclic. Suppose that  $p > 2$ . Then there is an element  $a \in G$  such that  $a \neq e$ . Since the cyclic subgroup  $\langle a \rangle = H_a := \{a^n : n \in \mathbb{Z}\}$  contains both  $a$  and  $e$ , we have  $|H_a| \geq 2$ . Since  $|H_a|$  divides  $|G| = p$  by Lagrange's theorem, we must have  $|H_a| = p$ , because  $p$  is prime. Then we must have  $G = H_a$ , and hence  $G$  is cyclic.  $\square$



**Corollary 2.7.10** (Euler's Theorem). *Let  $n \geq 2$  be an integer. Then for any positive integer  $a$  with  $\gcd(a, n) = 1$ , we have  $a^{\phi(n)} \equiv 1 \pmod{n}$ , where  $\phi(n)$  is the number of elements in the set  $\{k \in \mathbb{N} : 1 \leq k < n \text{ and } \gcd(k, n) = 1\}$ .*

*Proof.* Note that,  $U_n := \{[a] \in \mathbb{Z}_n : \gcd(a, n) = 1\}$  is a finite subset of  $\mathbb{Z}_n$  containing  $\phi(n)$  elements. Since  $U_n$  is a group with respect to the multiplication operation modulo  $n$ , for any  $[a] \in U_n$  we have  $[a]^{\phi(n)} = [1]$ . In other words,  $a^{\phi(n)} \equiv 1 \pmod{n}$ .  $\square$

**Corollary 2.7.11** (Fermat's little theorem). *If  $p > 0$  is a prime number, then for any positive integer  $a$  with  $\gcd(a, p) = 1$ , we have  $a^{p-1} \equiv 1 \pmod{p}$ .*

*Proof.* Since  $\phi(p) = |U_p| = p - 1$ , the result follows from the Corollary 2.7.10.  $\square$

**Exercise 2.7.12.** Show that  $2^{6000} - 1$  is divisible by 7.

*Solution.* Since  $\gcd(2, 7) = 1$ , by Fermat's little theorem we have  $2^{7-1} \equiv 1 \pmod{7}$ . So  $[2^6] = [1]$  in  $\mathbb{Z}_7$ . Then  $[2^6]^{1000} = [1]^{1000} = [1^{1000}] = [1]$  in  $\mathbb{Z}_7$ . Therefore,  $2^{6000} \equiv 1 \pmod{7}$ , and hence  $2^{6000} - 1$  is divisible by 7.  $\square$

**Exercise 2.7.13.** Show that  $15^{1000} - 1$  and  $105^{1200} - 1$  are divisible by 8.

**Exercise 2.7.14.** Define a relation  $\rho_R$  on  $G$  by setting

$$(a, b) \in \rho_R \text{ if } ab^{-1} \in H.$$

- (i) Show that  $\rho_R$  is an equivalence relation on  $G$ .
- (ii) Show that the  $\rho_R$ -equivalence class of  $a \in G$  in  $G$  is the subset of  $G$  defined by

$$[a]_{\rho_R} := \{b \in G : a^{-1}b \in H\} = \{ha : h \in H\} =: Ha.$$

The subset  $Ha \subseteq G$  is called the *right coset of  $H$  in  $G$  represented by  $a$* .

- (iii) Show that if  $G$  is abelian then  $aH = Ha$ , for all  $a \in G$ .
- (iv) Give an example of a group  $G$ , two subgroups  $H$  and  $K$  of  $G$ , and an element  $b \in G$  such that that  $bK \neq Kb$ , while  $aH = Ha$  holds, for all  $a \in G$ . (Hint: Take  $G = S_3$ , and

$$H := \{e, (1 \ 2 \ 3), (1 \ 3 \ 2)\} \subset S_3 \text{ and } K := \{e, (2 \ 3)\} \subset S_3.$$

Note that both  $H$  and  $K$  are subgroups of  $S_3$ . Verify that  $aH = Ha$ ,  $\forall a \in S_3$ , while for  $b = (1 \ 3 \ 2) \in S_3$  we have  $bK \neq Kb$ .)

- (v) Show that  $H$  and  $Ha$  have the same cardinality, for all  $a \in G$ .

The set of all distinct right cosets of  $H$  in  $G$  is denoted by

$$H \backslash G = \{Ha : a \in G\}.$$

**Lemma 2.7.15.** *Let  $H$  be a subgroup of a group  $G$ . Then there is a one-to-one correspondence between the set of all left cosets of  $H$  in  $G$  and the set of all right cosets of  $H$  in  $G$ . In other words, there is a bijective map  $\varphi : G/H \longrightarrow H \backslash G$ . Therefore, both the sets  $G/H$  and  $H \backslash G$  have the same cardinality.*

*Proof.* Define a map  $\varphi : \{aH : a \in G\} \longrightarrow \{Hb : b \in G\}$  by sending  $\varphi(aH) = Ha^{-1}$ , for all  $a \in G$ . Note that,  $aH = bH$  if and only if  $a^{-1}b \in H$  if and only if  $a^{-1}(b^{-1})^{-1} \in H$  if and only if  $Ha^{-1} = Hb^{-1}$ . Therefore,  $\varphi$  is well-defined and injective. To show  $\varphi$  bijective, note that given any  $Hb \in \{Hb : b \in G\}$  we have  $\varphi(b^{-1}H) = Hb$ . Thus,  $\varphi$  is surjective, and hence is a bijective map.  $\square$



**Definition 2.7.16.** Let  $H$  be a subgroup of a group  $G$ . We define the *index of  $H$  in  $G$* , denoted as  $[G : H]$ , to be the cardinality  $|G/H| = |H \backslash G|$ . In case, this is a finite number, the index  $[G : H]$  is the number of distinct left (and right) cosets of  $H$  in  $G$ .

**Exercise 2.7.17.** Let  $H$  and  $K$  be two subgroups of  $G$  of finite indices. Show that  $H \cap K$  is a subgroup of  $G$  of finite index.

**Example 2.7.18.** The index of  $n\mathbb{Z}$  in  $\mathbb{Z}$  is  $n$ . Indeed, given any two elements  $a, b \in \mathbb{Z}$ , we have  $a - b \in n\mathbb{Z}$  if and only if  $a \equiv b \pmod{n}$ . Therefore, the left coset of  $n\mathbb{Z}$  represented by  $a \in \mathbb{Z}$  is precisely the equivalence class

$$[a] := \{b \in \mathbb{Z} : a \equiv b \pmod{n}\} = a + n\mathbb{Z}.$$

Since there are exactly  $n$  such distinct equivalence classes by division algorithm, namely

$$a + n\mathbb{Z}, \text{ where } 0 \leq a \leq n-1;$$

(c.f. Example 2.1.32), we conclude that the index of  $n\mathbb{Z}$  in  $\mathbb{Z}$  is  $[\mathbb{Z} : n\mathbb{Z}] = n$ . We shall explain it later using group homomorphism and quotient group.

**Proposition 2.7.19 (Epimorphism of groups is surjective).** Let  $f : G \rightarrow H$  be a group homomorphism satisfying the following property:

- Given a group  $T$  and a group homomorphism  $\phi : H \rightarrow T$  with  $\phi \circ f$  trivial, we have  $\phi$  is trivial.

Then  $f$  is surjective.

*Proof.* Note that  $A := f(G)$  is a subgroup of  $H$ , and so we can consider the set

$$A \backslash H = \{Ah : h \in H\}$$

consisting of all distinct **right cosets of  $A$  in  $H$** . Let  $A'$  be a subset of  $H$  which is not a right coset of  $A$  in  $H$ , and let  $S = \{A'\} \cup H/A$ . Let  $T = \text{Aut}(S)$  be the symmetric group on  $S$ ; its elements are bijective maps from  $S$  onto itself and the group operation is given by composition of maps. Note that, given  $h \in H$ , consider the map

$$\varphi_h : A \backslash H \rightarrow A \backslash H$$

that sends  $Ah' \in A \backslash H$  to  $A(h'h) \in A \backslash H$ . Since  $(h'h)(h''h)^{-1} = h'h h^{-1} h''^{-1} = h'h''^{-1}$ , it follows that  $\varphi_h$  is well-defined and injective. Since  $\varphi_{h^{-1}} \circ \varphi_h = \text{Id}_{A \backslash H} = \varphi_h \circ \varphi_{h^{-1}}$ , the map  $\varphi_h$  is bijective.

Let  $\varphi : H \rightarrow T := \text{Aut}(S)$  be the map given by sending  $h \in H$  to the permutation  $\varphi(h) \in \text{Aut}(S)$  which is defined by

$$\varphi(h)(A') = A' \quad \text{and} \quad \varphi(h)|_{A \backslash H} = \varphi_h.$$

It is easy to verify that  $\varphi$  is a group homomorphism. Let  $\sigma \in T = \text{Aut}(S)$  be the permutation that interchanges  $A$  and  $A'$ , and keeps everything else fixed; i.e.,  $\sigma$  is the 2-cycle  $\sigma = (A \ A')$ . Then the map

$$(2.7.20) \quad \psi : H \rightarrow T, \quad h \mapsto \sigma^{-1} \varphi(h) \sigma,$$

is a group homomorphism (verify!).

If  $a \in A$ , then  $\varphi(a)(A) = Aa = A$  and  $\varphi(a)(A') = A'$ . Then  $\varphi(a) \in T$  is disjoint from the 2-cycle  $\sigma = (A \ A')$ , and hence they commute to give  $\psi(a) = \sigma^{-1} \varphi(a) \sigma = \varphi(a)$ . Therefore,  $\varphi|_A = \psi|_A$  and hence  $\varphi \circ f = \psi \circ f$ . Since  $f$  is an epimorphism, we have  $\varphi = \psi$ . Then  $\varphi(h) = \sigma^{-1} \varphi(h) \sigma$ , for all  $h \in H$ . Since  $\sigma = (A \ A')$  and  $\varphi(h)(A') = A'$ , we have  $\varphi(h)(A) =$

$(\sigma^{-1}\varphi(h)\sigma)(A) = \sigma^{-1}\varphi(h)(A') = \sigma^{-1}(A') = A$ . Since  $\varphi(h)(A) = Ah$  by definition, we have  $Ah = A$ , and hence  $h \in A$ . Since  $h \in H$  is arbitrary, we have  $A = H$ , as required.  $\square$

- Exercise 2.7.21.** (i) Does there exist a group isomorphism from  $(\mathbb{Q}, +)$  onto  $(\mathbb{Q}^*, \cdot)$ ?  
(ii) Does there exist a surjective group homomorphism from  $(\mathbb{Q}, +)$  onto  $(\mathbb{Q}^+, \cdot)$ ?  
(iii) Does there exist a non-trivial group homomorphism from  $\mathbb{Q}$  into  $\mathbb{Z}$ ?

## 2.8 Normal Subgroup & Quotient Group

In this section we introduce the notion of normal subgroup and give a construction of quotient of a group by its normal subgroup. Recall that the condition (QG1) in Definition 2.7.1 of quotient group suggests us to consider the set

$$G/H := \{gH : g \in G\}$$

consisting of all left cosets of  $H$  in  $G$  as a possible candidate for the set  $Q$ . Now question is what should be the appropriate group structure on it? Take any group homomorphism  $f : G \rightarrow T$  such that  $H \subseteq \text{Ker}(f)$ . Then we have  $f(a) = f(b)$  if  $a^{-1}b \in H$ . The commutativity of the diagram (2.7.2) tells us to send  $aH \in Q$  to  $f(a) \in T$  to define the map  $\tilde{f} : Q \rightarrow T$  which needs to be a group homomorphism. Then we should have

$$(2.8.1) \quad \tilde{f}((aH)(bH)) = f(ab) = \tilde{f}((ab)H), \forall a, b \in G.$$

This suggests us to define a binary operation on the set  $G/H = \{gH : g \in G\}$  by

$$(2.8.2) \quad (aH)(bH) := (ab)H, \forall a, b \in G.$$

**Proposition 2.8.3.** *The map  $G/H \times G/H \rightarrow G/H$  defined by sending  $(aH, bH)$  to  $(ab)H$  is well-defined if and only if*

$$(2.8.4) \quad g^{-1}hg \in H, \forall g \in G \text{ and } h \in H.$$

*Proof.* Suppose the above map is well-defined. Let  $h \in H$  and  $g \in G$  be arbitrary. Then  $hH = H$ , and hence  $(hH) \cdot (gH) = H \cdot (gH)$ . Since the above defined binary operation on  $G/H$  is well-defined, we have  $(hg)H = gH$  and hence  $g^{-1}hg \in H$ .

Conversely, suppose that  $g^{-1}hg \in H$ , for all  $g \in G$  and  $h \in H$ . Let  $a_1H = a_2H$  and  $b_1H = b_2H$ , for some  $a_1, a_2, b_1, b_2 \in G$ . Then  $h := a_1^{-1}a_2 \in H$  and  $b_1^{-1}b_2 \in H$ . Then

$$\begin{aligned} (a_1b_1)^{-1}(a_2b_2) &= b_1^{-1}a_1^{-1}a_2b_2 \\ &= b_1^{-1}hb_2, \text{ since } h := a_1^{-1}a_2. \\ &= (b_1^{-1}hb_1)(b_1^{-1}b_2) \in H, \end{aligned}$$

since  $H$  is a group and both  $b_1^{-1}hb_1$  and  $b_1^{-1}b_2$  are in  $H$ . Therefore,  $(a_1b_1)H = (a_2b_2)H$ , as required.  $\square$

Proposition 2.8.3 suggests us to reserve a terminology for those subgroups  $H$  of  $G$  that satisfies the property (2.8.4).

**Definition 2.8.5** (Normal Subgroup). A subgroup  $H$  of a group  $G$  is said to be *normal* in  $G$  if  $g^{-1}hg \in H$ ,  $\forall g \in G, h \in H$ . In this case we express it symbolically by  $H \trianglelefteq G$ .

**Exercise 2.8.6.** Let  $G$  be a group and  $H$  a subgroup of  $G$ . Given  $a \in G$ , let

$$Ha := \{ha : h \in H\} \subseteq G.$$

Show that the following are equivalent.

- (i)  $aH = Ha$ , for all  $a \in G$ .
- (ii)  $a^{-1}Ha = H$ , for all  $a \in G$ .
- (iii)  $a^{-1}Ha \subseteq H$ , for all  $a \in G$ .
- (iv)  $a^{-1}ha \in H$ , for all  $a \in G$  and  $h \in H$ .

**Proposition 2.8.7.** Any subgroup of index 2 is normal.

*Proof.* Let  $H$  be a subgroup of  $G$  such that  $[G : H] = 2$ . Then  $H$  has only two left (resp., right) cosets, namely  $H$  and  $aH$  (resp.,  $H$  and  $Ha$ ), where  $a \in G \setminus H$ . Since  $G = H \sqcup aH = H \sqcup Ha$ , for any  $a \in G \setminus H$ , we see that  $aH = Ha$ , for all  $a \in G$ , and hence  $aHa^{-1} = H$ , for all  $a \in G$ . This completes the proof.  $\square$

**Corollary 2.8.8.** For all  $n \geq 3$ ,  $A_n$  is a normal subgroup of  $S_n$ .

**Exercise 2.8.9.** (i) Show that any subgroups of an abelian group  $G$  is normal in  $G$ .

(ii) Let  $H = \langle (1 \ 2 \ 3) \rangle$  be the cyclic subgroup of  $S_3$  generated by the 3-cycle  $(1 \ 2 \ 3) \in S_3$ . Show that  $H$  is a normal subgroup of  $S_3$ .

(iii) Verify if the subgroup  $K := \langle (1 \ 2) \rangle$  of  $S_3$  is normal or not.

(iv) Determine all normal subgroups of  $S_3$ .

(v) Show that  $\text{SL}_n(\mathbb{R})$  is a normal subgroup of  $\text{GL}_n(\mathbb{R})$ , for all  $n \in \mathbb{N}$ .

**Exercise 2.8.10.** Show that  $S_4$  has no normal subgroup of order 3. (*Hint:* If  $\sigma \in S_4$  has order 3, then  $\sigma$  is a 3-cycle in  $S_4$ . Since there are  $\frac{4!}{3} = 8$  distinct 3-cycles in  $S_4$  (see Exercise 2.5.15), and all of them are conjugates (see Proposition 2.5.12), a normal subgroup  $H$  of  $S_4$  containing a 3-cycle contains at least 8 elements.)

**Exercise 2.8.11.** Let  $H$  be a subgroup of  $G$ . Let  $\rho = \{(a, b) \in G \times G : a^{-1}b \in H\} \subseteq G \times G$ . Note that  $\rho$  is an equivalence relation on  $G$ . Show that  $H$  is a normal subgroup of  $G$  if and only if  $\rho$  is a subgroup of the direct product group  $G \times G$  (see Exercise 2.1.34).

**Lemma 2.8.12.** The kernel of a group homomorphism  $f : G \rightarrow H$  is a normal subgroup of  $G$ .

*Proof.* For any  $a \in G$  and  $b \in \text{Ker}(f)$ , we have  $f(aba^{-1}) = f(a)f(b)f(a^{-1}) = f(a)e_H f(a)^{-1} = e_H$ , and hence  $aba^{-1} \in \text{Ker}(f)$ . Therefore,  $\text{Ker}(f)$  is a normal subgroup of  $G$ .  $\square$

**Exercise 2.8.13.** For  $n \geq 2$ , show that  $A_n$  is a normal subgroup of  $S_n$  by constructing a group homomorphism  $\varphi : S_n \rightarrow \mu_2 = \{1, -1\}$  such that  $\text{Ker}(\varphi) = A_n$ .

**Exercise 2.8.14.** For  $n \geq 1$ , show that  $\text{SL}_n(\mathbb{R})$  is a normal subgroup of  $\text{GL}_n(\mathbb{R})$  by constructing a group homomorphism  $\varphi : \text{GL}_n(\mathbb{R}) \rightarrow \mathbb{R}^*$  such that  $\text{Ker}(\varphi) = \text{SL}_n(\mathbb{R})$ .

**Lemma 2.8.15.** Let  $f : G \rightarrow H$  be a group homomorphism. If  $K$  is a normal subgroup of  $H$ , then  $f^{-1}(K)$  is a normal subgroup of  $G$ .

*Proof.* Suppose that  $K$  is a normal subgroup of  $H$ . Then for any  $a \in G$  and  $b \in f^{-1}(K)$ , we have  $f(aba^{-1}) = f(a)f(b)f(a^{-1}) \in K$ , and hence  $aba^{-1} \in f^{-1}(K)$ .  $\square$

**Exercise 2.8.16.** Show that  $N := \{A \in \text{GL}_n(\mathbb{C}) : |\det(A)| = 1\}$  is a normal subgroup of  $\text{GL}_n(\mathbb{C})$ .

**Remark 2.8.17.** Normal subgroup of a normal subgroup need not be normal. To elaborate it, there exists a group  $G$  together with a normal subgroup  $H$  of  $G$  such that  $H$  has a normal subgroup  $K$  which is not a normal subgroup of  $G$ . Can you give such an example?

**Theorem 2.8.18 (Existence of Quotient Group).** *Let  $H$  be a normal subgroup of a group  $G$ . Then the quotient group  $(Q, \pi)$  of  $G$  by  $H$  exists and is unique in the sense that if  $(Q, \pi)$  and  $(Q', \pi')$  are two quotients of  $G$  by  $H$ , then there exists a unique isomorphism of groups  $\varphi : Q \rightarrow Q'$  such that  $\varphi \circ \pi = \pi'$ . We denote  $Q$  by  $G/H$ .*

*Proof.* Since  $H$  is a normal subgroup of  $G$ ,

$$(aH)(bH) := (ab)H, \forall a, b \in G,$$

is a well-defined binary operation on the set  $G/H := \{aH : a \in G\}$ ; see Proposition 2.8.3. Given any  $a, b, c \in G$ , we have

$$(aH \cdot bH) \cdot cH = (ab)H \cdot cH = ((ab)c)H = (a(bc))H = aH \cdot (bc)H = aH \cdot (bH \cdot cH).$$

Therefore, the binary operation on  $G/H$  is associative. Given any  $aH \in G/H$ , we have

$$\begin{aligned} aH \cdot eH &= (ae)H = aH \\ \text{and } eH \cdot aH &= (ea)H = aH. \end{aligned}$$

Therefore,  $eH = H \in G/H$  is neutral element for the binary operation on  $G/H$ . Given any  $aH \in G/H$ , note that

$$\begin{aligned} aH \cdot a^{-1}H &= (aa^{-1})H = eH \\ \text{and } a^{-1}H \cdot aH &= (a^{-1}a)H = eH. \end{aligned}$$

Therefore,  $G/H$  is a group. Set  $Q := G/H$  and consider the map

$$(2.8.19) \quad \pi : G \longrightarrow Q \text{ defined by } \pi(a) = aH, \forall a \in G.$$

Clearly  $\pi$  is surjective and given  $a, b \in G$  we have  $\pi(ab) = (ab)H = (aH)(bH) = \pi(a)\pi(b)$ . Therefore,  $\pi$  is a group homomorphism. Since for any  $h \in H$ , we have  $\pi(h) = hH = eH = H$ , the neutral element of the group  $G/H$ , we see that  $H \subseteq \text{Ker}(\pi)$ . Let  $T$  be any group and  $t : G \rightarrow T$  be a group homomorphism satisfying  $t(h) = e_T$ , the neutral element of  $T$ , for all  $h \in H$ . Since  $aH = bH$  if and only if  $a^{-1}b \in H$ , applying  $\pi$  on  $a^{-1}b$  we see that  $\pi(a) = \pi(b)$ . Therefore, the map

$$(2.8.20) \quad \tilde{t} : G/H \rightarrow T, \quad aH \longmapsto t(a),$$

is well-defined. Since

$$\tilde{t}((aH)(bH)) = \tilde{t}((ab)H) = f(ab) = f(a)f(b) = \tilde{t}(aH)\tilde{t}(bH),$$

we conclude that  $\tilde{t}$  is a group homomorphism. Since  $(\tilde{t} \circ \pi)(a) = \tilde{t}(aH) = f(a)$ ,  $\forall a \in G$ , we have  $\tilde{t} \circ \pi = f$ . If  $\xi : G/H \rightarrow T$  is any group homomorphism satisfying  $\xi \circ \pi = f$ , then for any  $a \in G$  we have  $\tilde{t}(aH) = (\tilde{t} \circ \pi)(a) = f(a) = (\xi \circ \pi)(a) = \xi(aH)$ , and hence  $\tilde{t} = \xi$ . Therefore, the pair  $(G/H, \pi)$  satisfy the properties (QG1) and (QG2), and hence is a quotient of  $G$  by  $H$ . Uniqueness is already shown in Proposition 2.7.3.  $\square$

**Corollary 2.8.21.** *Let  $H$  be a normal subgroup of a group  $G$ , and let  $(G/H, \pi)$  be the associated quotient of  $G$  by  $H$ . Then  $\text{Ker}(\pi) = H$ .*

*Proof.* Since the group operation on the quotient group  $G/H := \{aH : a \in G\}$  is given by  $(aH)(bH) := (ab)H$ ,  $\forall aH, bH \in G/H$ , we have

$$\begin{aligned}\text{Ker}(\pi) &= \{a \in G : \pi(a) = H\} \\ &= \{a \in G : aH = H\} \\ &= \{a \in G : a \in H\} = H.\end{aligned}$$

This completes the proof.  $\square$

**Exercise 2.8.22.** Let  $G$  be a group such that  $G/Z(G)$  is cyclic. Show that  $G$  is abelian.

*Solution:* Let  $Z := Z(G)$ . Suppose that  $G/Z$  is cyclic. Then  $G/Z = \langle aZ \rangle$ , for some  $a \in G$ . Let  $x \in G$  be arbitrary. Then  $xZ = (aZ)^n = a^nZ$ , for some  $n \in \mathbb{Z}$ . Then  $a^{-n}x = (a^n)^{-1}x \in Z$ . Therefore,  $a^{-n}x = z$ , for some  $z \in Z$ , and so  $x = a^n z$ , for some  $z \in Z = Z(G)$ . Let  $y \in G$  be given. Then as before,  $y = a^m w$ , for some  $m \in \mathbb{Z}$  and  $w \in Z(G)$ . Since  $z, w \in Z(G)$ , we have  $xy = a^n z a^m w = a^m w a^n z = yx$ , as required.  $\square$

**Corollary 2.8.23.** There is no group  $G$  such that  $|G/Z(G)|$  is a prime number.

## 2.9 Isomorphism Theorems

Let  $G$  be a group. Given a normal subgroup  $K$  of  $G$ , let  $(G/K, \pi)$  be the associated quotient group of  $G$  by  $K$ , where

$$\pi : G \rightarrow G/K = \{aK : a \in G\}$$

is the natural quotient homomorphism given by

$$\pi(a) = aK, \quad \forall a \in G.$$

**Theorem 2.9.1.** Let  $f : G \rightarrow H$  be a group homomorphism. Let  $K$  be a normal subgroup of  $G$  such that  $K \subseteq \text{Ker}(f)$ . Then there is a unique group homomorphism  $\tilde{f} : G/K \rightarrow H$  such that  $\tilde{f} \circ \pi = f$ , where  $\pi : G \rightarrow G/K$  is the quotient homomorphism.

$$\begin{array}{ccc} G & \xrightarrow{f} & H \\ \pi \downarrow & \nearrow \tilde{f} & \\ G/K & & \end{array}$$

Furthermore,  $\tilde{f}$  is injective if and only if  $K = \text{Ker}(f)$ .

*Proof.* Since  $K$  is a normal subgroup of  $G$ , the quotient group  $G/K$  exists with the natural surjective group homomorphism  $\pi : G \rightarrow G/K$  defined by  $\pi(a) = aK$ ,  $\forall a \in G$ . Since  $K \subseteq \text{Ker}(f)$ , by universal property of quotient (see Definition 2.7.1) we have a unique group homomorphism  $\tilde{f} : G/K \rightarrow H$  such that  $\tilde{f} \circ \pi = f$ . The fact that  $\tilde{f}$  is a well-defined group homomorphism can also be directly checked by observing that

$$\tilde{f}(aK) = (\tilde{f} \circ \pi)(a) = f(a), \quad \forall a \in G.$$

Since  $\text{Ker}(\tilde{f}) = \{gK : f(g) = e_H\} = \{gK : g \in \text{Ker}(f)\}$ , we see that  $\text{Ker}(\tilde{f})$  is trivial (meaning that, it is a trivial subgroup) if and only if  $gK = K$ ,  $\forall g \in \text{Ker}(f)$ . This is equivalent to say that,  $g \in K$ ,  $\forall g \in \text{Ker}(f)$ , i.e.,  $\text{Ker}(f) \subseteq K$ . Since  $K \subseteq \text{Ker}(f)$  by assumption, it follows from Proposition 2.6.20 that  $\tilde{f}$  is injective if and only if  $K = \text{Ker}(f)$ .  $\square$

**Slogan:** To get a group homomorphism from a quotient group  $G/H$  to a group  $G'$ , thanks to Theorem 2.9.1 we just need to define a group homomorphism  $f : G \rightarrow G'$  such that  $H \subseteq \text{Ker}(f)$ .

**Example 2.9.2.** Let  $H_1$  and  $H_2$  be normal subgroups of  $G_1$  and  $G_2$ , respectively. Note that  $H_1 \times H_2$  is a normal subgroup of  $G_1 \times G_2$  (verify!). Let  $\pi_1 : G_1 \rightarrow G_1/H_1$  and  $\pi_2 : G_2 \rightarrow G_2/H_2$  be the natural quotient group homomorphisms. These give rise to a group homomorphism  $\phi : G_1 \times G_2 \rightarrow (G_1/H_1) \times (G_2/H_2)$  given by

$$\phi(a_1, a_2) = (\pi_1(a_1), \pi_2(a_2)) = (a_1H_1, a_2H_2), \forall (a_1, a_2) \in G_1 \times G_2.$$

Note that  $\phi$  is surjective because both  $\pi_1$  and  $\pi_2$  are so. Moreover,  $\text{Ker}(\phi) = H_1 \times H_2$  (verify!). Then by Theorem 2.9.1, given any normal subgroup  $K$  of  $G_1 \times G_2$  with  $K \leq H_1 \times H_2$ , there is a unique group homomorphism  $\tilde{\phi} : (G_1 \times G_2)/K \rightarrow (G_1/H_1) \times (G_2/H_2)$  such that  $\tilde{\phi} \circ \pi_K = \phi$ , where  $\pi_K : G_1 \times G_2 \rightarrow (G_1 \times G_2)/K$  is the natural quotient group homomorphism.

As an immediate corollary, we have the following.

**Corollary 2.9.3** (First Isomorphism Theorem). *Let  $f : G \rightarrow H$  be a surjective homomorphism of groups. Then  $f$  induces a natural isomorphism of groups  $\tilde{f} : G/\text{Ker}(f) \rightarrow H$ .*

*Proof.* Note that  $\text{Ker}(f)$  is a normal subgroup of  $G$ . It follows from Theorem 2.9.1 that the group homomorphism  $\tilde{f} : G/\text{Ker}(f) \rightarrow H$  induced by  $f$  is injective. Since  $f$  is surjective and  $\tilde{f} \circ \pi = f$ , where  $\pi : G \rightarrow G/\text{Ker}(f)$  is the natural surjective homomorphism, it follows that  $\tilde{f}$  is surjective. Therefore,  $\tilde{f}$  is a bijective group homomorphism, and hence is an isomorphism of groups.  $\square$

Let  $G$  be a group. Note that given a normal subgroup  $N$  of  $G$ , the quotient group  $G/N$  of  $G$  by  $N$  comes with a natural surjective group homomorphism  $\pi_N : G \rightarrow G/N$  such that  $\text{Ker}(\pi_N) = N$  (see Definition 2.7.1 and Corollary 2.8.21). On the other hand, given a group  $Q$  and a surjective group homomorphism  $\pi : G \rightarrow Q$ , its kernel  $\text{Ker}(\pi)$  is a normal subgroup of  $G$  such that  $G/\text{Ker}(\pi) \cong Q$  by the First isomorphism theorem (Corollary 2.9.3) for groups. This motivates us to define the following (c.f. Definition 2.7.1).

**Definition 2.9.4.** A *quotient group* of  $G$  is a pair  $(Q, \pi)$ , where  $Q$  is a group and  $\pi : G \rightarrow Q$  is a surjective group homomorphism.

As an immediate consequence, we have the following.

**Corollary 2.9.5.** *Given a group  $G$ , there is a one-to-one correspondence between the following two sets:*

- (i)  $\mathcal{N}_G :=$  the set of all normal subgroups of  $G$ , and
- (ii)  $\mathcal{Q}_G :=$  the set of all quotient groups of  $G$ .

*Proof.* Define a map  $\Phi : \mathcal{N}_G \rightarrow \mathcal{Q}_G$  by sending a normal subgroup  $N$  of  $G$  to the associated quotient group  $(G/N, \pi_N) \in \mathcal{Q}_G$ . Since  $\pi_N$  is a surjective group homomorphism with  $\text{Ker}(\pi_N) = N$ , the map  $\Phi$  admits an inverse, namely  $\Psi : \mathcal{Q}_G \rightarrow \mathcal{N}_G$  given by sending a quotient group  $(Q, \pi)$  of  $G$  to the kernel  $N := \text{Ker}(\pi) \in \mathcal{N}_G$ . Since the pairs  $(G/N, \pi_N)$  and  $(Q, \pi)$  are uniquely isomorphic, we conclude that  $\Phi$  and  $\Psi$  are inverse to each other. This completes the proof.  $\square$

**Proposition 2.9.6.** *The group  $\mathbb{Z}_n$  is isomorphic to  $\mathbb{Z}/n\mathbb{Z}$ .*

*Proof.* Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}_n$  be the map defined by

$$f(k) = [k], \forall k \in \mathbb{Z}.$$

Since

$$f(k_1 + k_2) = [k_1 + k_2] = [k_1] + [k_2] = f(k_1) + f(k_2), \forall k_1, k_2 \in \mathbb{Z},$$

we see that  $f$  is a group homomorphism. Clearly  $f$  is surjective (verify!). Note that  $\text{Ker}(f) = \{k \in \mathbb{Z} : [k] = [0]\} = n\mathbb{Z}$ . Then by first isomorphism theorem we have  $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}_n$ .  $\square$

**Proposition 2.9.7.** Any finite cyclic group of order  $n$  is isomorphic to  $\mathbb{Z}_n$ .

*Proof.* Let  $G$  be a finite cyclic group of order  $n$ . Then there exists  $a \in G$  such that  $\langle a \rangle = \{a^k : k \in \mathbb{Z}\} = G$ . Define a map  $f : \mathbb{Z} \rightarrow G$  by

$$f(k) = a^k, \forall k \in \mathbb{Z}.$$

Since

$$f(k_1 + k_2) = a^{k_1 + k_2} = a^{k_1} a^{k_2} = f(k_1) f(k_2), \forall k_1, k_2 \in \mathbb{Z},$$

$f$  is a group homomorphism. Clearly  $f$  is surjective because every element of  $G$  is of the form  $a^k$ , for some  $k \in \mathbb{Z}$ . Then by first isomorphism theorem  $G$  is isomorphic to  $\mathbb{Z}/\text{Ker}(f)$ . Note that,  $\text{Ker}(f) = \{k \in \mathbb{Z} : a^k = e\}$ . Since  $G$  is a cyclic group of order  $n$  generated by  $a$ , we have  $\text{ord}(a) = n$  (see Corollary 2.3.11). Then we have  $\text{Ker}(f) = \{k \in \mathbb{Z} : a^k = e\} = n\mathbb{Z}$ . Therefore,  $G \cong \mathbb{Z}/n\mathbb{Z}$ . Since  $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}_n$  by Theorem 2.9.6, we have  $G \cong \mathbb{Z}_n$ .  $\square$

**Exercise 2.9.8.** Show that any group of order 4 is isomorphic to either  $\mathbb{Z}_4$  or  $K_4$ .

**Exercise 2.9.9.** Show that any group of order 6 is isomorphic to either  $\mathbb{Z}_6$  or  $S_3$ .

**Exercise 2.9.10.** Use the signature homomorphism  $S_n \rightarrow \mu_2 = \{1, -1\}$  to show that  $A_n$  is the only index 2 subgroup of  $S_n$ .

**Exercise 2.9.11.** Show that  $\text{SL}_2(\mathbb{Z}_3)$  and  $S_4$  are two non-isomorphic non-commutative groups of order 24.

### 2.9.1 Inner Automorphisms

Let  $G$  be a group. Given  $a \in G$ , the map  $\varphi_a : G \rightarrow G$  defined by

$$\varphi_a(b) = aba^{-1}, \forall b \in G,$$

is a group homomorphism. Indeed,

$$\varphi_a(bc) = a(bc)a^{-1} = (aba^{-1})(aca^{-1}) = \varphi_a(b)\varphi_a(c), \forall b, c \in G.$$

Since  $\text{Ker}(\varphi_a) = \{b \in G : aba^{-1} = e\} = \{e\}$ ,  $\varphi_a$  is injective. Given  $c \in G$ , note that  $\varphi_a(a^{-1}ca) = a(a^{-1}ca)a^{-1} = c$ , and so  $\varphi_a$  is surjective. Therefore,  $\varphi_a$  is an isomorphism.

**Definition 2.9.12.** An automorphism  $\varphi \in \text{Aut}(G)$  is said to be an *inner automorphism* of  $G$  if there exists  $a \in G$  such that  $\varphi(b) = aba^{-1}$ , for all  $b \in G$ .

**Proposition 2.9.13.** Let  $G$  be a group. Let  $\text{Inn}(G)$  be the set of all inner automorphisms of  $G$ . Then  $\text{Inn}(G)$  is a subgroup of  $\text{Aut}(G)$ .

*Proof.* Note that the identity map  $\text{Id}_G : G \rightarrow G$  is in  $\text{Inn}(G)$ . Given  $f, g \in \text{Inn}(G)$ , there exists  $a, b \in G$  such that  $f$  and  $g(x) = bxb^{-1}$ , for all  $x \in G$ . Then  $f^{-1} = \varphi_{a^{-1}}$ , and that  $(\varphi_a^{-1} \circ \varphi_b)(x) = a^{-1}bxb^{-1}a = (a^{-1}b)x(a^{-1}b)^{-1} = \varphi_{a^{-1}b}(x)$ , for all  $x \in G$ . Therefore,  $\text{Inn}(G)$  is a subgroup of  $\text{Aut}(G)$ .  $\square$

**Proposition 2.9.14.** The map  $\varphi : G \rightarrow \text{Inn}(G)$  that sends  $a \in G$  to the map  $\varphi_a : G \rightarrow G$  defined by

$$\varphi(a)(b) = aba^{-1}, \forall b \in G,$$



is a surjective group homomorphism with kernel  $Z(G)$ . Consequently,  $G/Z(G) \cong \text{Inn}(G)$ .

*Proof.* Let  $a, b \in G$  be given. Then for any  $x \in G$  we have  $\varphi(ab)(x) = (ab)x(ab)^{-1} = a(bxb^{-1})a^{-1} = a(\varphi_b(x))a^{-1} = (\varphi_a \circ \varphi_b)(x)$ , and hence  $\varphi(ab) = \varphi(a) \circ \varphi(b)$ . Therefore,  $\varphi$  is a group homomorphism. Since every element of  $\text{Inn}(G)$  is of the form  $\varphi_a$ , for some  $a \in G$ , the map  $\varphi$  is surjective. Since  $\text{Ker}(\varphi) = \{a \in G : \varphi(a) = \text{Id}_G\} = \{a \in G : aba^{-1} = b, \forall b \in G\} = Z(G)$ , by the first isomorphism theorem for groups we have  $G/Z(G) \cong \text{Inn}(G)$ .  $\square$

**Exercise 2.9.15.** Let  $G$  be a group such that  $G/Z(G)$  is cyclic. Show that  $\text{Inn}(G)$  is a trivial subgroup of  $\text{Aut}(G)$ .

**Theorem 2.9.16** (Second Isomorphism Theorem). *Let  $G$  be a group. Let  $H$  and  $K$  be subgroups of  $G$  with  $K$  normal in  $G$ . Then*

- (i)  $HK$  is a subgroup of  $G$ ,
- (ii)  $K$  is a normal subgroup of  $HK$ , and
- (iii)  $H/(H \cap K) \cong HK/K$ .

*Proof.* (i) Let  $h \in H$  and  $k \in K$  be arbitrary. Since  $K$  is a normal subgroup of  $G$ , we have  $hk = (hkh^{-1})h \in KH$  and so  $HK \subseteq KH$ . Similarly,  $kh = h(h^{-1}kh) \in HK$  shows that  $KH \subseteq HK$ . Thus  $HK = KH$  and hence  $HK$  is a subgroup of  $G$  by Theorem 2.4.3.

(ii) Clearly  $K$  is a subgroup of  $HK$ . Since  $K$  is normal in  $G$ , given any  $a \in HK \subseteq G$  and  $k \in K$  we have  $aka^{-1} \in K$ , and hence  $K$  is a normal subgroup of  $HK$ .

(iii) Define a map  $\varphi : H \rightarrow HK/K$  by  $\varphi(a) = aK$ , for all  $a \in H$ . Since  $\varphi(ab) = (ab)K = (aK)(bK) = \varphi(a)\varphi(b)$ , for all  $a, b \in H$ ,  $\varphi$  is a group homomorphism. Since  $K \in HK/K$  is the neutral element, given any  $h \in H$  and  $k \in K$  we have  $(hk)K = (hK)(kK) = hK = \varphi(h)$ , and so  $\varphi$  is surjective. Since

$$\text{Ker}(\varphi) = \{h \in H : hK = K\} = \{h \in H : h \in K\} = H \cap K,$$

by first isomorphism theorem (see Corollary 2.9.3) we have  $H/(H \cap K) \cong HK/K$ .  $\square$

**Example 2.9.17.** Let  $m, n \in \mathbb{N}$  with  $\gcd(m, n) = 1$ . Consider the subgroups  $H = m\mathbb{Z}$  and  $K = n\mathbb{Z}$  of  $(\mathbb{Z}, +)$ . Since  $\mathbb{Z}$  is abelian,  $K$  is a normal subgroup of  $\mathbb{Z}$ . Since  $\gcd(m, n) = 1$ , there exists  $a, b \in \mathbb{Z}$  such that  $am + bn = 1$ , and so  $1 \in H + K$ . Since  $\gcd(m, n) = 1$ , we have  $\text{lcm}(m, n) = mn$ , and so  $H \cap K = mn\mathbb{Z}$ . Then by the second isomorphism theorem we have  $m\mathbb{Z}/mn\mathbb{Z} = H/(H \cap K) \cong (H + K)/K = \mathbb{Z}/n\mathbb{Z}$ . Generalize this to the case when  $m$  and  $n$  are not necessarily coprime.

**Exercise 2.9.18.** Use the second isomorphism theorem for groups to prove the following.

- (i)  $3\mathbb{Z}/15\mathbb{Z} \cong \mathbb{Z}/5\mathbb{Z}$ , and
- (ii)  $6\mathbb{Z}/30\mathbb{Z} \cong 2\mathbb{Z}/10\mathbb{Z}$ . (Hint: Take  $H = 6\mathbb{Z}$  and  $K = 10\mathbb{Z}$ ).

**Theorem 2.9.19** (Abelianization). *Let  $G$  be a group. Then upto isomorphism there exists a unique pair  $(G_{\text{ab}}, \Phi)$  consisting of an abelian group  $G_{\text{ab}}$  and a surjective group homomorphism  $\Phi : G \rightarrow G_{\text{ab}}$  satisfying the following universal property: given any abelian group  $H$  and a group homomorphism  $f : G \rightarrow H$ , there exists a unique group homomorphism  $\tilde{f} : G_{\text{ab}} \rightarrow H$  such that  $\tilde{f} \circ \Phi = f$ .*

$$\begin{array}{ccc} G & \xrightarrow{f} & H \\ \Phi \downarrow & \nearrow \tilde{f} & \\ G_{\text{ab}} & & \end{array}$$

The group  $G_{\text{ab}}$  is known as the maximal abelian quotient or the abelianization of  $G$ .



*Proof. Uniqueness:* First we prove uniqueness of the pair  $(G_{\text{ab}}, \Phi)$  upto unique isomorphism of groups. Suppose that  $(K, g)$  be another such pair consisting of an abelian group  $K$  and a surjective group homomorphism  $g : G \rightarrow K$  such that the pair  $(K, g)$  satisfies the above universal property. Taking  $(H, f) = (G_{\text{ab}}, \Phi)$  we find a unique group homomorphism  $\tilde{\Phi} : K \rightarrow G_{\text{ab}}$  such that  $\tilde{\Phi} \circ g = \Phi$ .

$$\begin{array}{ccccc} & & G & & \\ & g \swarrow & \downarrow \Phi & \searrow g & \\ K & \xrightarrow{\tilde{\Phi}} & G_{\text{ab}} & \xrightarrow{\tilde{g}} & K \end{array}$$

Applying universal property of  $(G_{\text{ab}}, \Phi)$  with  $(H, f) = (K, g)$ , we have a unique group homomorphism  $\tilde{g} : G_{\text{ab}} \rightarrow K$  such that  $\tilde{g} \circ \Phi = g$ . Since the composite map  $\tilde{g} \circ \tilde{\Phi} : K \rightarrow K$  is a group homomorphism, by the universal property of the pair  $(K, g)$  we have  $\tilde{g} \circ \tilde{\Phi} = \text{Id}_K$ , where  $\text{Id}_K : K \rightarrow K$  is the identity map of  $K$ . Similarly, we have  $\tilde{\Phi} \circ \tilde{g} = \text{Id}_{G_{\text{ab}}}$ . Therefore, both  $\tilde{g} : K \rightarrow G_{\text{ab}}$  and  $\tilde{\Phi} : G_{\text{ab}} \rightarrow K$  are isomorphism of groups. Since both  $\tilde{\Phi}$  and  $\tilde{g}$  are unique and  $\tilde{\Phi} \circ g = \Phi$  and  $\tilde{g} \circ \Phi = g$ , we conclude that the pair  $(K, g)$  is uniquely isomorphic to  $(G_{\text{ab}}, \Phi)$ .

*Existence:* To prove existence of the pair  $(G_{\text{ab}}, \Phi)$ , consider the elements of  $G$  of the form

$$[a, b] := aba^{-1}b^{-1},$$

where  $a, b \in G$ , called *commutators* in  $G$ . Clearly  $[a, b] = e$  if  $G$  is abelian. Let

$$[G, G] := \langle aba^{-1}b^{-1} : a, b \in G \rangle$$

be the subgroup of  $G$  generated by all commutators of elements of  $G$ . The subgroup  $[G, G]$  is known as the *commutator subgroup* or the *derived subgroup* of  $G$ . Since

$$ghg^{-1} = ghg^{-1}h^{-1}h = [g, h]h, \quad \forall g, h \in G,$$

taking  $h \in [G, G]$  we see that  $[G, G]$  is a normal subgroup of  $G$ . Let  $G_{\text{ab}} := G/[G, G]$  be the associated quotient group, and let  $\Phi : G \rightarrow G_{\text{ab}}$  be the natural quotient map which sends  $a \in G$  to the coset  $a[G, G] \in G/[G, G] = G_{\text{ab}}$ . Let us denote by  $\bar{a}$  the image of  $a \in G$  in  $G/[G, G]$  under the quotient map  $\Phi : G \rightarrow G/[G, G]$ . Since

$$(ab)(ba)^{-1} = aba^{-1}b^{-1} \in [G, G], \quad \forall a, b \in G,$$

we have  $\bar{a}\bar{b} = \bar{b}\bar{a}$  in  $G/[G, G]$ . Therefore,  $G/[G, G]$  is commutative. If  $f : G \rightarrow H$  is a group homomorphism, then

$$f([a, b]) = f(aba^{-1}b^{-1}) = [f(a), f(b)], \quad \forall a, b \in G.$$

Now suppose that  $H$  is abelian. Then for any  $a, b \in G$ , we have  $[f(a), f(b)] = e$ , and so  $[a, b] \in \text{Ker}(f)$ . Therefore,  $[G, G] \subseteq \text{Ker}(f)$ . Consequently, by universal property of quotient (see Definition 2.7.1) there is a unique homomorphism  $\tilde{f} : G/[G, G] \rightarrow H$  such that  $\tilde{f} \circ \Phi = f$ . This completes the proof of existence part.  $\square$

**Proposition 2.9.20.** *The commutator subgroup of  $S_n$  is  $A_n$ , for all  $n \geq 3$ .*

*Proof.* Since the signature map  $\text{sgn} : S_n \rightarrow \mu_2 = \{1, -1\}$  defined by

$$\text{sgn}(\sigma) = \begin{cases} 1, & \text{if } \sigma \text{ is even,} \\ -1, & \text{if } \sigma \text{ is odd,} \end{cases}$$

is a group homomorphism (see Lemma 2.6.4), we have  $\text{sgn}(\sigma)^{-1} = \text{sgn}(\sigma)$ , for all  $\sigma \in S_n$ . Therefore, given  $\sigma, \tau \in S_n$  we have

$$\text{sgn}([\sigma, \tau]) = \text{sgn}(\sigma \circ \tau \circ \sigma^{-1} \tau^{-1}) = \text{sgn}(\sigma) \text{sgn}(\tau) \text{sgn}(\sigma)^{-1} \text{sgn}(\tau)^{-1} = 1.$$

Therefore,  $[\sigma, \tau] \in A_n$ , for all  $\sigma, \tau \in S_n$ , and hence  $[S_n, S_n] \subseteq A_n$ . To show the reverse inclusion, note that  $A_n$  is generated by 3-cycles, for all  $n \geq 3$  (see Exercise 2.5.34), and any 3-cycle  $(i \ j \ k)$  in  $S_n$  can be written as

$$(i \ j \ k) = (i \ j) \circ (i \ k) \circ (i \ j)^{-1} \circ (i \ k)^{-1},$$

which is an element of  $[S_n, S_n]$ . Thus  $A_n \subseteq [S_n, S_n]$ . This completes the proof.  $\square$

**Exercise 2.9.21.** Show that the abelianization of  $S_n$  is isomorphic to  $\mathbb{Z}_2$ , for all  $n \geq 3$ .

**Exercise 2.9.22.** Given any two groups  $H$  and  $K$ , let  $\text{Hom}(H, K)$  be the set of all group homomorphisms from  $H$  into  $K$ . Fix an integer  $n \geq 3$ .

- (i) Given an abelian group  $G$ , show that there is a natural bijective map  $\text{Hom}(S_n, G) \longrightarrow \text{Hom}(\mathbb{Z}_2, G)$ .
- (ii) Find the number of elements in  $\text{Hom}(S_n, \mathbb{Z}_4 \times \mathbb{Z}_6)$ .

**Exercise 2.9.23.** Show that  $S_4$  has no normal subgroup of order 8. (*Hint:* If  $H$  is a normal subgroup of  $S_4$  of order 8, the quotient group  $S_4/H$  is abelian, and hence  $A_4 = [S_4, S_4] \subseteq H$ , a contradiction.)

**Theorem 2.9.24** (Third Isomorphism Theorem). *Let  $H$  and  $K$  be normal subgroups of  $G$  with  $K \subseteq H$ . Then we have an isomorphism of groups  $(G/K)/(H/K) \cong G/H$ .*

*Proof.* Since  $H$  and  $K$  are normal subgroups of  $G$  and  $K \subseteq H$ , that  $K$  is a normal subgroup of  $H$ , and the associated quotient groups

- (i)  $\phi : G \rightarrow G/H$ ,
- (ii)  $\psi : G \rightarrow G/K$ , and
- (iii)  $\eta : H \rightarrow H/K$

exist. Let  $\iota_H : H \hookrightarrow G$  be the inclusion of  $H$  into  $G$ . Then the composite map

$$H \xrightarrow{\iota_H} G \xrightarrow{\psi} G/K$$

is a group homomorphism with kernel  $K$ , and hence we get an injective group homomorphism

$$H/K \hookrightarrow G/K.$$

Given  $h \in H$  and  $a \in G$ , we have  $aha^{-1} \in H$ , and so  $(aK)(hK)(aK)^{-1} = (ah)K \cdot a^{-1}K = (aha^{-1})K \in H/K$ . Therefore,  $H/K$  is a normal subgroup of  $G/K$ , and hence the associated quotient group  $\pi : G/K \rightarrow (G/K)/(H/K)$  exists. Consider the diagram

$$\begin{array}{ccc} G & \xrightarrow{\psi} & G/K \\ \phi \downarrow & & \downarrow \pi \\ G/H & \xrightarrow{\widetilde{\pi \circ \psi}} & (G/K)/(H/K) \end{array}$$

Note that  $H/K \in (G/K)/(H/K)$  is the neutral element of the group  $(G/K)/(H/K)$ . Moreover, the composite map  $\pi \circ \psi$  is a surjective group homomorphism with kernel

$$\begin{aligned} \text{Ker}(\pi \circ \psi) &= \{a \in G : \pi(\psi(a)) = e\} \\ &= \{a \in G : \pi(aK) = e\} \\ &= \{a \in G : aK(H/K) = H/K\} \\ &= \{a \in G : aK \in H/K\} \\ &= \{a \in G : a \in H\}, \text{ since the map } H/K \hookrightarrow G/K \text{ is injective.} \\ &= H \end{aligned}$$

Then by first isomorphism theorem (Corollary 2.9.3) applied to the group homomorphism  $\pi \circ \psi$  we have the required isomorphism  $G/H \cong (G/K)/(H/K)$  of groups.  $\square$

**Corollary 2.9.25** (Correspondence Theorem). *Let  $f : G \rightarrow H$  be a surjective group homomorphism. Consider the following two sets:*

- (i)  $\mathcal{A} :=$  the set of a subgroups of  $G$  containing  $\text{Ker}(f)$ , and
- (ii)  $\mathcal{B} :=$  the set of all subgroups of  $H$ .

Then there is an inclusion preserving bijective map

$$\Phi : \mathcal{A} \rightarrow \mathcal{B}$$

such that a subgroup  $N \in \mathcal{A}$  of  $G$  is normal in  $G$  if and only if  $\Phi(N)$  is normal in  $H$ .

*Proof.* Define a map  $\Phi : \mathcal{A} \rightarrow \mathcal{B}$  by sending a subgroup  $N$  of  $G$  containing  $\text{Ker}(f)$  to its image  $f(N)$ . Note that  $f(N)$  is a subgroup of  $H$  by Proposition 2.6.7 (i), and hence is an element of  $\mathcal{B}$ . Conversely, given a subgroup  $K$  of  $H$ , its preimage  $f^{-1}(K)$  is a subgroup of  $G$  by Proposition 2.6.7 (ii). Since  $e_H \in K$  we have  $\text{Ker}(f) = f^{-1}(e) \subseteq f^{-1}(K)$ . Thus,  $f^{-1}(K) \in \mathcal{A}$ . This gives a map

$$\Psi : \mathcal{B} \rightarrow \mathcal{A}, \quad K \mapsto f^{-1}(K).$$

It remains to show that  $\Phi$  and  $\Psi$  are inverse to each other. Given  $N \in \mathcal{A}$ , we have  $(\Psi \circ \Phi)(N) = f^{-1}(f(N)) \supseteq N$ . If  $a \in f^{-1}(f(N))$ , then  $f(a) = f(b)$ , for some  $b \in N$ . Then  $f(ab^{-1}) = f(a)f(b)^{-1} = e_H$  implies  $ab^{-1} \in \text{Ker}(f) \subseteq N$ , and so  $a = (ab^{-1})b \in N$ . Therefore,  $(\Psi \circ \Phi)(N) = f^{-1}(f(N)) = N$ , for all  $N \in \mathcal{A}$ , and hence  $\Psi \circ \Phi = \text{Id}_{\mathcal{A}}$ . Conversely, given  $K \in \mathcal{B}$ , we have  $(\Phi \circ \Psi)(K) = f(f^{-1}(K)) = K$ , since  $f$  is surjective. Thus  $\Phi \circ \Psi = \text{Id}_{\mathcal{B}}$ . This completes the proof.  $\square$

**Exercise 2.9.26.** Let  $H$  be a normal subgroup of a group  $G$ . Show that every subgroup of  $G/H$  is of the form  $K/H$ , for some subgroup  $K$  of  $G$  containing  $H$ .

**Exercise 2.9.27.** Let  $\pi : G \rightarrow Q$  be a surjective group homomorphism. Let  $H$  be a normal subgroup of  $G$  and let  $\pi_H : H \rightarrow Q$  be the restriction of  $\pi$  on  $H$ . If  $K = H \cap \text{Ker}(\pi)$ , show that the induced map  $\widetilde{\pi}_H : H/K \rightarrow Q$  is injective, and it identifies  $H/K$  as a normal subgroup of  $Q$ .

## 2.10 Direct Product & Direct Sum of Groups

**Definition 2.10.1.** The *direct product* of a family of groups  $\{G_\alpha : \alpha \in \Lambda\}$  is a pair  $(G, \{\pi_\alpha\}_{\alpha \in \Lambda})$ , where  $G$  is a group and  $\{\pi_\alpha : G \rightarrow G_\alpha\}_{\alpha \in \Lambda}$  is a family of group homomorphisms such that given any group  $H$  and a family of group homomorphisms  $\{f_\alpha : H \rightarrow G_\alpha\}_{\alpha \in \Lambda}$  there exists a

unique group homomorphism  $f : H \rightarrow G$  such that  $\pi_\alpha \circ f = f_\alpha$ , for all  $\alpha \in \Lambda$ .

$$\begin{array}{ccc} H & & \\ \downarrow \exists! f & \searrow f_\alpha & \\ G & \xrightarrow{\pi_\alpha} & G_\alpha \end{array}$$

**Theorem 2.10.2 (Existence & Uniqueness of Product of Groups).** *The direct product of a family of groups exists and is unique upto a unique isomorphism in the sense that if  $(G, \{g_\alpha : G \rightarrow G_\alpha\}_{\alpha \in \Lambda})$  and  $(H, \{h_\alpha : H \rightarrow G_\alpha\}_{\alpha \in \Lambda})$  are direct products of the family of groups  $\{G_\alpha : \alpha \in \Lambda\}$ , then there exists a unique isomorphism of groups  $\phi : G \rightarrow H$  such that  $h_\alpha \circ \phi = g_\alpha$ , for all  $\alpha \in \Lambda$ . We denote by  $\prod_{\alpha \in \Lambda} G_\alpha$  the underlying group of the direct product of the family of groups  $\{G_\alpha : \alpha \in \Lambda\}$ .*

*Proof.* Since  $(G, \{g_\alpha\}_{\alpha \in \Lambda})$  is a direct product by assumption, for the test object  $(H, \{h_\beta : H \rightarrow G_\beta\}_{\beta \in \Lambda})$  we have a group homomorphism  $\varphi : G \rightarrow H$  such that  $\pi_\alpha \circ \varphi = h_\alpha$ ,  $\forall \alpha \in \Lambda$ . Interchanging the roles of  $(G, \{g_\alpha\}_{\alpha \in \Lambda})$  and  $(H, \{h_\alpha\}_{\alpha \in \Lambda})$  we have a group homomorphism  $\psi : H \rightarrow G$  such that  $\pi_\alpha \circ \psi = g_\alpha$ ,  $\forall \alpha \in \Lambda$ . Since both  $\psi \circ \varphi : G \rightarrow G$  and  $\text{Id}_G : G \rightarrow G$  are group homomorphisms satisfying

$$f_\alpha \circ (\psi \circ \varphi) = f_\alpha \quad \text{and} \quad f_\alpha \circ \text{Id}_G = f_\alpha, \quad \forall \alpha \in \Lambda,$$

it follows that  $\psi \circ \varphi = \text{Id}_G$ . Similarly,  $\varphi \circ \psi = \text{Id}_H$ , and hence  $\varphi : G \rightarrow H$  is the unique isomorphism such that  $h_\alpha \circ \varphi = g_\alpha$ ,  $\forall \alpha \in \Lambda$ .

For a construction, let

$$\prod_{\alpha \in \Lambda} G_\alpha := \{f : \Lambda \rightarrow \prod_{\alpha \in \Lambda} G_\alpha \mid f(\alpha) \in G_\alpha, \forall \alpha \in \Lambda\}.$$

Given  $f, g \in \prod_{\alpha \in \Lambda} G_\alpha$  we define

$$fg : \Lambda \rightarrow \prod_{\alpha \in \Lambda} G_\alpha$$

by

$$(fg)(\alpha) := f(\alpha)g(\alpha), \quad \forall \alpha \in \Lambda.$$

Clearly  $fg \in \prod_{\alpha \in \Lambda} G_\alpha$ , and  $(fg)h = f(gh)$ ,  $\forall f, g, h \in \prod_{\alpha \in \Lambda} G_\alpha$ . Let  $e_\alpha \in G_\alpha$  be the neutral element, for all  $\alpha \in \Lambda$ . Then the map  $e : \Lambda \rightarrow \prod_{\alpha \in \Lambda} G_\alpha$  given by  $e(\alpha) = e_\alpha$ ,  $\forall \alpha \in \Lambda$  satisfies  $ef = fe = f$ ,  $\forall f \in \prod_{\alpha \in \Lambda} G_\alpha$ . Given  $f \in \prod_{\alpha \in \Lambda} G_\alpha$  we define  $f^{-1} \in \prod_{\alpha \in \Lambda} G_\alpha$  by  $f^{-1}(\alpha) = (f_\alpha)^{-1} \in G_\alpha$ ,  $\forall \alpha \in \Lambda$ . Then  $ff^{-1} = e = f^{-1}f$ . Therefore,  $\prod_{\alpha \in \Lambda} G_\alpha$  is a group. For each  $\beta \in \Lambda$ , we define a map  $\pi_\beta : \prod_{\alpha \in \Lambda} G_\alpha \rightarrow G_\beta$  by  $\pi_\beta(f) = f(\beta)$ . Then  $\pi_\beta$  is a group homomorphism. Given a group  $H$  and a family  $\{h_\alpha : H \rightarrow G_\alpha\}_{\alpha \in \Lambda}$  of group homomorphisms, we define a map  $\psi : H \rightarrow \prod_{\alpha \in \Lambda} G_\alpha$  that sends  $a \in H$  to the function  $\psi_a : \Lambda \rightarrow \prod_{\alpha \in \Lambda} G_\alpha$  defined by  $\psi_a(\alpha) = h_\alpha(a)$ ,  $\forall \alpha \in \Lambda$ . Then it is straight forward to verify that  $\psi$  is a group homomorphism satisfying  $\pi_\alpha \circ \psi = h_\alpha$ ,  $\forall \alpha \in \Lambda$ .  $\square$

**Example 2.10.3 (External Direct Product of  $G_1, \dots, G_n$ ).** Let  $G_1, \dots, G_n$  be a finite family of groups, not necessarily distinct. Define a binary operation on the Cartesian product  $G := G_1 \times \dots \times G_n$  by

$$(2.10.4) \quad (a_1, \dots, a_n) \cdot (b_1, \dots, b_n) := (a_1 b_1, \dots, a_n b_n),$$

where  $a_i, b_i \in G_i$ , for all  $i = 1, \dots, n$ . Given  $a_i, b_i, c_i \in G_i$ , for each  $i \in \{1, \dots, n\}$ , we have

$$\begin{aligned} ((a_1, \dots, a_n) \cdot (b_1, \dots, b_n)) \cdot (c_1, \dots, c_n) &= (a_1 b_1, \dots, a_n b_n) \cdot (c_1, \dots, c_n) \\ &= ((a_1 b_1) c_1, \dots, (a_n b_n) c_n) \\ &= (a_1 (b_1 c_1), \dots, a_n (b_n c_n)) \\ &= (a_1, \dots, a_n) \cdot ((b_1, \dots, b_n) \cdot (c_1, \dots, c_n)) \end{aligned}$$

Therefore, the above defined binary operation on the set  $G$  is associative. Let  $e_i \in G_i$  be the neutral element of  $G_i$ , for all  $i \in \{1, \dots, n\}$ . Then given any  $a_i \in G_i$ , for each  $i$ , we have

$$(a_1, \dots, a_n) \cdot (e_1, \dots, e_n) = (a_1, \dots, a_n) = (e_1, \dots, e_n) \cdot (a_1, \dots, a_n).$$

Since

$$(a_1, \dots, a_n) \cdot (a_1^{-1}, \dots, a_n^{-1}) = (e_1, \dots, e_n) = (a_1^{-1}, \dots, a_n^{-1}) \cdot (a_1, \dots, a_n),$$

we conclude that  $(a_1, \dots, a_n)^{-1} = (a_1^{-1}, \dots, a_n^{-1}) \in G$ . Therefore,  $G = G_1 \times \dots \times G_n$  is a group with respect to the binary operation defined in (2.10.4).

For each  $i \in \{1, \dots, n\}$ , let

$$(2.10.5) \quad p_i : G_1 \times \dots \times G_n \rightarrow G_i$$

be the map defined by

$$(2.10.6) \quad p_i(a_1, \dots, a_n) = a_i, \quad \forall (a_1, \dots, a_n) \in G_1 \times \dots \times G_n.$$

Clearly  $p_i$  is a surjective group homomorphism (verify!). Let  $H$  be a group and let  $\{f_i : H \rightarrow G_i\}_{1 \leq i \leq n}$  be a family of group homomorphisms. Define a map  $f : H \rightarrow G_1 \times \dots \times G_n$  by

$$(2.10.7) \quad f(h) = (f_1(h), \dots, f_n(h)), \quad \forall h \in H.$$

Then given any  $a, b \in H$  we have

$$\begin{aligned} f(ab) &= (f_1(ab), \dots, f_n(ab)) \\ &= (f_1(a)f_1(b), \dots, f_n(a)f_n(b)) \\ &= (f_1(a), \dots, f_n(a))(f_1(b), \dots, f_n(b)) \\ &= f(a)f(b). \end{aligned}$$

Therefore,  $f$  is a group homomorphism. Clearly  $p_i \circ f = f_i$ , for all  $i \in \{1, \dots, n\}$ . Suppose that  $f' : H \rightarrow G_1 \times \dots \times G_n$  is any group homomorphism such that  $p_i \circ f' = f_i$ , for all  $i \in \{1, \dots, n\}$ . Let  $h \in H$  be arbitrary. Let  $f'(h) = (a_1, \dots, a_n) \in G_1 \times \dots \times G_n$ . Then  $f_i(h) = (p_i \circ f')(h) = p_i(a_1, \dots, a_n) = a_i$ , for all  $i \in \{1, \dots, n\}$ , and hence  $f'(h) = (a_1, \dots, a_n) = (f_1(h), \dots, f_n(h)) = f(h)$ . Therefore,  $f' = f$ , and hence by universal property of product of groups (see Definition 2.10.1) we conclude that  $G_1 \times \dots \times G_n$  is a direct product of  $G_1, \dots, G_n$ . The group  $G_1 \times \dots \times G_n$  is also known as the **external direct product of  $G_1, \dots, G_n$** .

**Corollary 2.10.8.** *The direct product of a finite family of finite groups  $G_1, \dots, G_n$  is a group of order  $|G_1| \cdots |G_n|$ . Moreover,  $G_1 \times \dots \times G_n$  is abelian if and only if  $G_i$  is abelian, for all  $i \in I_n$ .*

**Exercise 2.10.9.** Given any two groups  $G$  and  $H$ , show that  $Z(G \times H) = Z(G) \times Z(H)$ .

**Proposition 2.10.10.** *Let  $G := G_1 \times \dots \times G_n$  be the external direct product of the family of groups  $G_1, \dots, G_n$ . For each  $i \in I_n := \{1, \dots, n\}$ , let  $H_i = \{(a_1, \dots, a_n) \in G : a_j = e_j, \forall j \neq i\} \subseteq G$ . Then we have the following.*

- (i)  $H_i$  is a normal subgroup of  $G$ , for all  $i \in I_n$ .
- (ii) Every element  $a \in G$  can be uniquely expressed as  $a = h_1 \cdots h_n$ , with  $h_i \in H_i$ , for all  $i \in I_n$ .

(iii)  $H_i \cap (H_1 \cdots H_{i-1} H_{i+1} \cdots H_n) = \{e\}$ , for all  $i \in I_n$ .

(iv)  $G = H_1 \cdots H_n$ .

*Proof.* (i) Since  $(e_1, \dots, e_n) \in H_i$ , so  $H_i \neq \emptyset$ . Let  $a := (a_1, \dots, a_n), b := (b_1, \dots, b_n) \in H_i$ . Then  $a_j = e_j = b_j, \forall j \neq i$ , and hence  $a_j^{-1}b_j = e_j$ , for all  $j \neq i$ . Therefore,  $a^{-1}b = (a_1^{-1}b_1, \dots, a_n^{-1}b_n) \in H_i$ , and hence  $H_i$  is a subgroup of  $G$ . Let  $a = (a_1, \dots, a_n) \in G$  and  $b := (b_1, \dots, b_n) \in H_i$  be arbitrary. Then  $b_j = e_j$ , for all  $j \neq i$ , and so  $a_j b_j a_j^{-1} = a_j e_j a_j^{-1} = e_j$ , for all  $j \neq i$ . This shows that  $aba^{-1} = (a_1, \dots, a_n)(b_1, \dots, b_n)(a_1^{-1}, \dots, a_n^{-1}) \in H_i$ . Therefore,  $H_i$  is a normal subgroup of  $G$ , for all  $i \in I_n$ .

(ii) Let  $a \in G$  be given. Then  $a = (a_1, \dots, a_n)$ , where  $a_i \in G_i, \forall i \in I_n$ . Let  $h_i \in G$  be the element whose  $i$ -th entry is  $a_i$  and for  $j \neq i$ , its  $j$ -th entry is  $e_j \in G_j$ . In other words,  $h_i := (h_{i1}, \dots, h_{in}) \in G$ , where

$$h_{ij} := \begin{cases} e_j, & \text{if } j \neq i, \\ a_i, & \text{if } j = i. \end{cases}$$

Then  $h_i \in H_i$ , for all  $i \in I_n$ , and  $h_1 \cdots h_n = (a_1, \dots, a_n) = a$ . To show uniqueness of this expression, let  $a = k_1 \cdots k_n$ , where  $k_i \in H_i$ , for all  $i \in I_n$ . If  $k_{ij} \in G_j$  denote the  $j$ -th entry of  $k_i \in H_i$ , then  $k_{ij} = e_j$ , for  $j \neq i$ . Therefore,

$$(a_1, \dots, a_n) = a = h_1 \cdots h_n = k_1 \cdots k_n = (k_{11}, \dots, k_{nn}).$$

Then  $a_i = h_{ii}$ , for all  $i \in I_n$ . This shows that  $k_i = h_i$ , for all  $i \in I_n$ . This proves uniqueness.

(iii) Let  $a = (a_1, \dots, a_n) \in H_i \cap (H_1 \cdots H_{i-1} H_{i+1} \cdots H_n)$ . Since  $a \in H_i$ , we have  $a_j = e_j, \forall j \neq i$ . Since  $a \in H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$ , we have

$$(2.10.11) \quad a = h_1 \cdots h_{i-1} h_{i+1} \cdots h_n$$

for some  $h_j \in H_j, \forall j \neq i$ . Since  $h_j = (h_{1j}, \dots, h_{nj}) \in H_j$ , we have

$$h_{kj} = e_k \in G_k, \forall k \neq j.$$

If  $b_k$  denote the  $k$ -th component of the product  $h_1 \cdots h_{i-1} h_{i+1} \cdots h_n$  in  $G_1 \times \cdots \times G_n$ , then

$$(2.10.12) \quad b_k = \begin{cases} e_i, & \text{if } k = i, \\ h_{ik}, & \text{if } k \neq i. \end{cases}$$

Comparing the  $j$ -th component of both sides of the equation (2.10.11), we have

$$a_j = e_j \in G_j, \forall j \in I_n.$$

(iv) It follows from (ii) that  $G \subseteq H_1 \cdots H_n$ . Since  $H_i$  is a subgroup of  $G$ , for all  $i \in I_n$ , we have  $H_1 \cdots H_n \subseteq G$ . Hence the result follows.  $\square$

**Lemma 2.10.13.** Let  $G$  be a group. Let  $H, K$  be two normal subgroups of  $G$  such that  $H \cap K = \{e\}$ . Then given any  $h \in H$  and  $k \in K$  we have  $hk = kh$ . Consequently,  $[H, K] = \{e\}$ .

*Proof.* Since  $H$  is normal in  $G$ , we have  $(hk)(kh)^{-1} = h(kh^{-1}k^{-1}) \in H$ . Similarly, since  $K$  is normal in  $G$ , we have  $(hk)(kh)^{-1} = (hkh^{-1})k^{-1} \in K$ . Therefore,  $(hk)(kh)^{-1} \in H \cap K = \{e\}$ , and hence  $hk = kh$  in  $G$ .  $\square$

**Exercise 2.10.14.** Is the conclusion of the Lemma 2.10.13 still holds if we assume exactly one of  $H$  and  $K$  is normal in  $G$ ?

**Lemma 2.10.15.** Let  $G$  be a group. Let  $H$  and  $K$  be normal subgroups of  $G$ . Then  $HK$  is a normal subgroup of  $G$ .

*Proof.* Since  $H$  and  $K$  are normal in  $G$ , it follows that  $HK$  is a subgroup of  $G$ . Let  $a \in G$  and  $h \in H, k \in K$  be arbitrary. Then  $a(hk)a^{-1} = (aha^{-1})(aka^{-1}) \in HK$ . Therefore,  $HK$  is a normal subgroup of  $G$ .  $\square$

**Definition 2.10.16.** Let  $G$  be a group and let  $H_1, \dots, H_n$  be normal subgroups of  $G$ . Then  $G$  is said to be an *internal direct product of  $H_1, \dots, H_n$*  if every element  $a \in G$  can be **uniquely** expressed as  $a = h_1 \cdots h_n$  with  $h_i \in H_i$ , for all  $i \in \{1, \dots, n\}$ .

**Proposition 2.10.17.** Let  $G = G_1 \times \cdots \times G_n$  be the external direct product of a finite collection of (not necessarily distinct) groups  $G_1, \dots, G_n$ , and  $H_i := \{(a_1, \dots, a_n) \in G : a_j = e_j, \forall j \neq i\}$ , for each  $i \in I_n$ . Then  $G$  is an internal direct product of  $H_1, \dots, H_n$ , respectively.

*Proof.* It follows from Proposition 2.10.10 (ii) that given  $a \in G$  there exists  $a_i \in H_i$ , for each  $i \in I_n$ , such that  $a = a_1 \cdots a_n$ . To show that this expression for  $a$  is unique, let

$$a = a_1 \cdots a_n = b_1 \cdots b_n,$$

for some  $a_i, b_i \in H_i, \forall i \in I_n$ . Note that each  $H_i$  is a normal subgroup of  $G$  by Proposition 2.10.10 (i), and  $K_i := H_1 \cdots H_{i-1}H_{i+1} \cdots H_n$  is a normal subgroups of  $G$  by Lemma 2.10.15. Moreover,  $H_i \cap K_i = \{e\}$  by Proposition 2.10.10 (iii). Then using Lemma 2.10.13 we have

$$\begin{aligned} e &= a^{-1}a = (a_1 \cdots a_n)^{-1}b_1 \cdots b_n \\ &= a_n^{-1} \cdots a_1^{-1}b_1 \cdots b_n \\ &= (a_1^{-1}b_1) \cdots (a_n^{-1}b_n). \end{aligned}$$

Then for each  $i \in I_n$ , we have

$$b_i^{-1}a_i = (a_1^{-1}b_1) \cdots (a_{i-1}^{-1}b_{i-1})(a_{i+1}^{-1}b_{i+1}) \cdots (a_n^{-1}b_n) \in H_i \cap K_i = \{e\},$$

and hence  $a_i = b_i$ , for all  $i \in I_n$ . This completes the proof.  $\square$

**Theorem 2.10.18.** Let  $\{H_1, \dots, H_n\}$  be a finite collection of normal subgroups of  $G$ . Let  $K_i := H_1 \cdots H_{i-1}H_{i+1} \cdots H_n, \forall i \in I_n$ . Then  $G$  is an internal direct product of  $H_1, \dots, H_n$  if and only if

- (i)  $G = H_1 \cdots H_n$ , and
- (ii)  $H_i \cap K_i = \{e\}$ , for all  $i \in I_n$ .

Moreover, in this case we have an isomorphism of groups  $G \cong H_1 \times \cdots \times H_n$ .

*Proof.* Suppose that  $G$  is an internal direct product of  $H_1, \dots, H_n$ , respectively. Let  $a \in G$  be given. Then for each  $i \in I_n$ , there exists unique  $a_i \in H_i$  such that  $a = a_1 \cdots a_n$ . Therefore,  $G \subseteq H_1 \cdots H_n$ , and hence  $G = H_1 \cdots H_n$ . Let  $a \in H_i \cap K_i$ . Then  $a \in H_i$  gives  $a = e_1 \cdots e_{i-1}ae_{i+1} \cdots e_n$ , where  $e_j \in H_j$  is the neutral element of  $H_j$ , for all  $j$ . Again,  $a \in K_i = H_1 \cdots H_{i-1}H_{i+1} \cdots H_n$  gives  $a = a_1 \cdots a_{i-1}ea_{i+1} \cdots a_n$ , where  $a_j \in H_j, \forall j \neq i$ . Then from the uniqueness of representation of  $a$  as product of elements from  $H_j$ 's, we see that  $a = e$ . Therefore,  $H_i \cap K_i = \{e\}$ .

Conversely, suppose that (i) and (ii) holds. By (i) given  $a \in G$ , there exists  $a_i \in H_i$ , for each  $i \in I_n$ , such that  $a = a_1 \cdots a_n$ . Suppose that for each  $i \in I_n$ , there exists  $b_i \in H_i$  such that  $a = b_1 \cdots b_n$ . Then as shown in the proof of the above Proposition, we have

$$\begin{aligned} e &= a^{-1}a = (a_1 \cdots a_n)^{-1}b_1 \cdots b_n \\ &= a_n^{-1} \cdots a_1^{-1}b_1 \cdots b_n \\ &= (a_1^{-1}b_1) \cdots (a_n^{-1}b_n). \end{aligned}$$

Then for each  $i \in I_n$ , we have

$$b_i^{-1}a_i = (a_1^{-1}b_1) \cdots (a_{i-1}^{-1}b_{i-1})(a_{i+1}^{-1}b_{i+1}) \cdots (a_n^{-1}b_n) \in H_i \cap K_i = \{e\},$$

and hence  $a_i = b_i$ , for all  $i \in I_n$ . This completes the proof.  $\square$

**Exercise 2.10.19.** Let  $G$  be a finite group of order  $mn$ , where  $\gcd(m, n) = 1$ . If  $H$  and  $K$  are normal subgroups of  $G$  of orders  $m$  and  $n$ , respectively, show that  $G$  is isomorphic to the direct product group  $H \times K$ .

**Corollary 2.10.20.** If  $m, n \in \mathbb{Z}$  with  $\gcd(m, n) = 1$ , then  $\mathbb{Z}_{mn} \cong \mathbb{Z}_m \times \mathbb{Z}_n$ .

**Theorem 2.10.21 (Direct Sum of Groups).** Let  $\{G_\alpha : \alpha \in \Lambda\}$  be a family of groups. Then there is a pair  $(A, \{\iota_\alpha\}_{\alpha \in \Lambda})$ , consisting of a group  $A$  and a family of group monomorphisms

$$\{\iota_\alpha : G_\alpha \rightarrow A\}_{\alpha \in \Lambda}$$

satisfying the following universal property: Given any abelian group  $T$  and a family of group homomorphisms  $\{f_\alpha : G_\alpha \rightarrow T\}_{\alpha \in \Lambda}$ , there exists a unique group homomorphism  $f : A \rightarrow T$  such that  $f \circ \iota_\alpha = f_\alpha$ ,  $\forall \alpha \in \Lambda$ .

$$\begin{array}{ccc} G_\alpha & \xrightarrow{f_\alpha} & T \\ \downarrow \iota_\alpha & \nearrow f & \\ A & & \end{array}$$

The pair  $(A, \{\iota_\alpha\}_{\alpha \in \Lambda})$  is uniquely determined by the universal property, and is called the **direct sum** of the family of groups  $\{G_\alpha\}_{\alpha \in \Lambda}$ , and is denoted by  $\bigoplus_{\alpha \in \Lambda} G_\alpha$ .

*Proof.* Uniqueness of the pair  $(A, \{\iota_\alpha\}_{\alpha \in \Lambda})$  follows from the universal property. We now prove existence. We write the group operation of  $G_\alpha$  additively. Given  $\alpha \in \Lambda$ , let  $0_\alpha$  be the neutral element of  $G_\alpha$ , and  $\pi_\alpha : \prod_{\beta \in \Lambda} G_\beta \rightarrow G_\alpha$  be the natural projection homomorphism. Given  $x \in \prod_{\alpha \in \Lambda} G_\alpha$ , let  $x_\alpha := \pi_\alpha(x) \in G_\alpha$ . Consider the subset

$$A := \left\{ x \in \prod_{\alpha \in \Lambda} G_\alpha \mid \pi_\alpha(x) = 0_\alpha, \text{ for all but finitely many } \alpha \in \Lambda \right\}.$$

Clearly  $0 := (0_\alpha)_{\alpha \in \Lambda} \in A$ , and given any  $x, y \in A$ ,  $\pi_\alpha(x - y) = x_\alpha - y_\alpha = 0_\alpha$ , for all but finitely many  $\alpha \in \Lambda$ , and so  $x - y \in A$ . Therefore,  $A$  is a subgroup of  $\prod_{\alpha \in \Lambda} G_\alpha$ . For each  $\alpha \in \Lambda$ , let  $\iota_\alpha : G_\alpha \rightarrow A$  be the map defined by  $\iota_\alpha(g) = x$ , where

$$\pi_\beta(x) := \begin{cases} g, & \text{if } \beta = \alpha, \\ e_\beta, & \text{if } \beta \neq \alpha. \end{cases}$$

Clearly  $\iota_\alpha$  is an injective group homomorphism, for all  $\alpha \in \Lambda$ . Let  $T$  be an abelian group. Let  $f_\alpha : G_\alpha \rightarrow T$  be a group homomorphism, for each  $\alpha \in \Lambda$ . Define a map  $f : A \rightarrow T$  by

$$f(a) = \sum_{\alpha \in \Lambda} f_\alpha(\pi_\alpha(a)), \quad \forall a \in A.$$

Note that the above sum is finite. Since  $f_\alpha : G_\alpha \rightarrow T$  is a group homomorphism,  $f_\alpha(0_\alpha) = 0_T \in T$ , and hence  $f(\iota_\alpha(g)) = f_\alpha(g)$ , for all  $g \in G_\alpha$ . Therefore,  $f \circ \iota_\alpha = f_\alpha$ ,  $\forall \alpha \in \Lambda$ . Uniqueness of  $f$  is easy to see.  $\square$

**Remark 2.10.22.** Note that  $\bigoplus_{\alpha \in \Lambda} G_\alpha = \prod_{\alpha \in \Lambda} G_\alpha$ , whenever the index set  $\Lambda$  is finite.



**Exercise 2.10.23.** Let  $G$  and  $H$  be cyclic groups of prime order  $p$  generated by  $x \in G$  and  $y \in H$ , respectively. Show that  $G \times H$  is an abelian group of order  $p^2$  that is not cyclic. Show that

$$\langle x \rangle, \langle xy \rangle, \langle xy^2 \rangle, \dots, \langle xy^{p-1} \rangle \text{ and } \langle y \rangle$$

are all possible distinct subgroups of  $G \times H$  of order  $p$ .

**Exercise 2.10.24.** Find the number of distinct subgroups of order  $p$  of the cyclic group  $\mathbb{Z}_{p^n}$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ .

## 2.11 Group Action

Let  $G$  be a group and let  $X$  be a non-empty set.

**Definition 2.11.1.** A *left  $G$ -action* on  $X$  is a map

$$\sigma : G \times X \rightarrow X$$

satisfying the following conditions:

- (i)  $\sigma(e, x) = x, \forall x \in X$ , and
- (ii)  $\sigma(b, \sigma(a, x)) = \sigma(ba, x), \forall a, b \in G, x \in X$ .

For notational simplicity, we write  $ax$  for  $\sigma(a, x)$ .

**Remark 2.11.2.** We can define a *right  $G$ -action* on  $X$  to be a map

$$\tau : X \times G \rightarrow X$$

satisfying the following conditions:

- (i)  $\tau(x, e) = x, \forall x \in X$ , and
- (ii)  $\tau(\tau(x, a), b) = \tau(x, ab), \forall a, b \in G, x \in X$ .

For notational simplicity, we write  $xa$  for  $\tau(x, a)$ .

**Example 2.11.3.** (i) Given a group  $G$  and a non-empty set  $X$ , the map

$$\sigma : G \times X \rightarrow X$$

defined by

$$\sigma(a, x) = x, \forall a \in G \text{ and } x \in X,$$

is a left  $G$ -action on  $X$ , known as the *trivial left  $G$ -action on  $X$* . Similarly, we have a trivial right  $G$ -action  $\tau : X \times G \rightarrow X$  on  $X$  that sends  $(x, a) \in X \times G$  to  $x \in X$ .

- (ii) For each integer  $n \geq 2$ , the group  $S_n$  acts on the set  $I_n := \{k \in \mathbb{N} : 1 \leq k \leq n\}$  by sending  $(\sigma, i) \in S_n \times I_n$  to  $\sigma(i) \in I_n$ . Clearly for  $\sigma = e \in S_n$  we have  $\sigma(i) = i, \forall i \in I_n$ , and  $(\sigma\tau)(i) = \sigma(\tau(i)), \forall i \in I_n, \sigma, \tau \in S_n$ .
- (iii) Given a non-empty set  $X$ , let  $S(X)$  be the group of all symmetries on  $X$ ; its elements are bijective maps from  $X$  onto itself, and the group operation is given by composition of maps. Then the group  $S(X)$  acts on  $X$  from the left.

(iv) Let  $H$  be a normal subgroup of a group  $G$ . For example,  $H = Z(G)$ . Then the map  $\varphi : G \times H \rightarrow H$  defined by

$$\varphi(a, h) = aha^{-1}, \forall a \in G, h \in H,$$

is a  $G$ -action on  $H$ . Indeed,  $\varphi(e, h) = ehe^{-1} = h, \forall h \in H$ , and

$$\varphi(a, \varphi(b, h)) = \varphi(a, bhb^{-1}) = a(bhb^{-1})a^{-1} = (ab)h(ab)^{-1} = \varphi(ab, h), \forall a, b \in G, h \in H.$$

**Lemma 2.11.4** (*Permutation representation of a  $G$ -action*). Given a group  $G$  and a non-empty set  $X$ , there is a one-to-one correspondence between the set of all left  $G$ -actions on  $X$  and the set of all group homomorphisms from  $G$  into the symmetric group  $S(X)$  on  $X$ .

*Proof.* Let  $\mathcal{A}$  be the set of all left  $G$ -actions on  $X$ , and let  $\mathcal{B} := \text{Hom}(G, S(X))$  be the set of all group homomorphisms from  $G$  into  $S(X)$ . Define a map  $\Phi : \mathcal{A} \rightarrow \mathcal{B}$  by sending a left  $G$ -action  $\sigma : G \times X \rightarrow X$  to the map

$$(2.11.5) \quad f_\sigma : G \rightarrow S(X)$$

that sends  $a \in G$  to the map

$$(2.11.6) \quad f_\sigma(a) : X \rightarrow X, x \mapsto \sigma(a, x).$$

We first show that  $f_\sigma(a)$  is bijective and hence is an element of  $S(X)$ . Let  $x, y \in X$  be such that  $\sigma(a, x) = \sigma(a, y)$ . Then we have

$$\begin{aligned} x &= \sigma(e, x) = \sigma(a^{-1}, \sigma(a, x)) \\ &= \sigma(a^{-1}, \sigma(a, y)) \\ &= \sigma(e, y) = y. \end{aligned}$$

Therefore,  $f_\sigma(a)$  is injective. Given  $y \in X$ , note that  $x := \sigma(a^{-1}, y) \in X$ , and that

$$f_\sigma(a)(x) = \sigma(a, x) = \sigma(a, \sigma(a^{-1}, y)) = \sigma(e, y) = y.$$

This shows that  $f_\sigma(a)$  is surjective. Therefore,  $f_\sigma(a) \in S(X)$ , for all  $a \in G$ . To show  $f_\sigma : G \rightarrow S(X)$  is a group homomorphism, note that given  $a, b \in G$  we have

$$\begin{aligned} f_\sigma(ab)(x) &= \sigma(ab, x) = \sigma(a, \sigma(b, x)) \\ &= f_\sigma(a)(f_\sigma(b)(x)) \\ &= (f_\sigma(a) \circ f_\sigma(b))(x), \forall x \in X, \end{aligned}$$

and hence  $f_\sigma(ab) = f_\sigma(a) \circ f_\sigma(b), \forall a, b \in G$ . Therefore,  $f_\sigma$  is a group homomorphism, known as the *permutation representation* of  $G$  associated to the left  $G$ -action  $\sigma$  on  $X$ . Thus,  $f_\sigma \in \mathcal{B}$ .

Given a group homomorphism  $f : G \rightarrow S(X)$ , consider the map  $\sigma_f : G \times X \rightarrow X$  defined by

$$\sigma_f(a, x) = f(a)(x), \forall a \in G, x \in X.$$

We show that  $\sigma_f$  is a left  $G$ -action on  $X$ . Since  $f : G \rightarrow S(X)$  is a group homomorphism,  $f(e) = \text{Id}_X$  in  $S(X)$ . Therefore,  $\sigma_f(e, x) = f(e)(x) = x, \forall x \in X$ . Since  $f : G \rightarrow S(X)$  is a group homomorphism, given  $a, b \in G$  we have  $f(ab) = f(a) \circ f(b)$ , and hence given any  $x \in X$  we have

$$\begin{aligned} f(ab)(x) &= (f(a) \circ f(b))(x) \\ \Rightarrow \sigma_f(ab, x) &= f(a)(\sigma_f(b, x)) \\ \Rightarrow \sigma_f(ab, x) &= \sigma_f(a, \sigma_f(b, x)). \end{aligned}$$

Therefore,  $\sigma_f$  is a left  $G$ -action on  $X$ . Thus we get a map  $\Psi : \mathcal{B} \rightarrow \mathcal{A}$  defined by

$$\Psi(f) = \sigma_f, \forall f \in \mathcal{B}.$$

It remains to check that  $\Psi \circ \Phi = \text{Id}_{\mathcal{A}}$  and  $\Phi \circ \Psi = \text{Id}_{\mathcal{B}}$ . Given a left  $G$ -action  $\tau : G \times X \rightarrow X$  on  $X$ , we have  $(\Psi \circ \Phi)(\tau) = \Psi(f_\tau) = \sigma_{f_\tau}$ . Since

$$\sigma_{f_\tau}(a, x) = f_\tau(a)(x) = \tau(a, x), \forall (a, x) \in G \times X,$$

we have  $(\Psi \circ \Phi)(\tau) = \tau, \forall \tau \in \mathcal{A}$ . Therefore,  $\Psi \circ \Phi = \text{Id}_{\mathcal{A}}$ . Conversely, given a group homomorphism  $g : G \rightarrow S(X)$ , we have  $(\Phi \circ \Psi)(g) = \Phi(\sigma_g) = f_{\sigma_g}$ . Since  $f_{\sigma_g}(a) = \sigma_g(a, -) = g(a)$ ,  $\forall a \in G$ , we conclude that  $(\Phi \circ \Psi)(g) = g, \forall g \in \mathcal{B}$ . Therefore,  $\Phi \circ \Psi = \text{Id}_{\mathcal{B}}$ . This completes the proof.  $\square$

**Definition 2.11.7 (Faithful action).** A left  $G$ -action  $\sigma : G \times X \rightarrow X$  on a non-empty set  $X$  is said to be *faithful* if  $\text{Ker}(f_\sigma) = \{e\}$ , where  $f_\sigma : G \rightarrow S(X)$  is the permutation representation of  $G$  associated to  $\sigma$  (see (2.11.5) and (2.11.6) in Lemma 2.11.4).

**Example 2.11.8.** The multiplicative group  $\mathbb{R}^* := \mathbb{R} \setminus \{0\}$  acts on  $V := \mathbb{R}^n$  by scalar multiplication

$$\sigma : \mathbb{R}^* \times V \rightarrow V$$

defined by

$$\sigma(t, (a_1, \dots, a_n)) := (ta_1, \dots, ta_n), \forall t \in \mathbb{R}^*, (a_1, \dots, a_n) \in \mathbb{R}^n.$$

Note that  $\sigma$  is a left  $\mathbb{R}^*$ -action on  $V = \mathbb{R}^n$ . The permutation representation

$$f_\sigma : \mathbb{R}^* \rightarrow S(V)$$

associated to  $\sigma$  is given by sending  $t \in \mathbb{R}^*$  to the map

$$f_\sigma(t) : V \rightarrow V, (a_1, \dots, a_n) \mapsto (ta_1, \dots, ta_n).$$

Since

$$\begin{aligned} \text{Ker}(f_\sigma) &= \{t \in \mathbb{R}^* : f_\sigma(t) = \text{Id}_V\} \\ &= \{t \in \mathbb{R}^* : tv = v, \forall v \in V\} \\ &= \{1\} \end{aligned}$$

is trivial, we conclude that  $\sigma$  is a faithful left  $\mathbb{R}^*$ -action on  $V = \mathbb{R}^n$ .

**Example 2.11.9.** Recall that Cayley's theorem (Theorem 2.6.23) says that any group  $G$  is isomorphic to a subgroup of the permutation group  $S(G)$  on  $G$ . This can be explained using group action as follow. Consider the left translation map

$$\sigma : G \times G \rightarrow G$$

defined by

$$\sigma(a, x) = ax, \forall a, x \in G.$$

Note that  $\sigma$  is a left  $G$ -action on itself, called the *left regular action of  $G$  on itself*, and the associated permutation representation  $f_\sigma : G \rightarrow S(G)$  that sends  $a \in G$  to the bijective map

$$f_\sigma(a) : G \rightarrow G, x \mapsto ax,$$

Then  $f_\sigma$  is a group homomorphism with

$$\begin{aligned}\text{Ker}(f_\sigma) &= \{a \in G : f_\sigma(a) = \text{Id}_G\} \\ &= \{a \in G : ax = x, \forall x \in G\} \\ &= \{e_G\}\end{aligned}$$

is trivial, and hence  $\sigma$  is a faithful action.

Given a left  $G$ -action  $\sigma : G \times X \rightarrow X$  on  $X$ , we define a relation  $\sim_\sigma$  on  $X$  by setting

$$(2.11.10) \quad x \sim_\sigma y \text{ if } y = \sigma(a, x), \text{ for some } a \in G.$$

Note that  $\sim_\sigma$  is an equivalence relation on  $X$  (verify!). The  $\sim_\sigma$ -equivalence class of  $x \in X$  is the subset

$$(2.11.11) \quad \text{Orb}_G(x) := \{\sigma(a, x) : a \in G\} \subseteq X,$$

called the *orbit* of  $x$  under the left  $G$ -action  $\sigma$  on  $X$ . Note that

- (i)  $x \in \text{Orb}_G(x)$ ,  $\forall x \in X$ , and
- (ii) given  $x, y \in X$ , either  $\text{Orb}_G(x) = \text{Orb}_G(y)$  or  $\text{Orb}_G(x) \cap \text{Orb}_G(y) = \emptyset$ .

Therefore,  $X$  is a disjoint union of distinct  $G$ -orbits of elements of  $X$ . A  $G$ -action  $\sigma : G \times X \rightarrow X$  is said to be *transitive* if  $\text{Orb}_G(x) = \text{Orb}_G(y)$ , for all  $x, y \in X$ . Therefore,  $\sigma$  is transitive if and only if given any two elements  $x, y \in X$ , there exists  $a \in G$  such that  $\sigma(a, x) = y$ .

**Proposition 2.11.12.** Let  $\sigma : G \times X \rightarrow X$  be a left  $G$ -action on  $X$ . For each  $x \in X$  the subset

$$G_x := \{a \in G : \sigma(a, x) = x\}$$

is a subgroup of  $G$ , called the *stabilizer* or the *isotropy subgroup* of  $x$ , and sometimes it is also denoted by  $\text{Stab}_G(x)$ .

*Proof.* Since  $\sigma(e, x) = x$ ,  $e \in G_x$ . Let  $a, b \in G_x$  be arbitrary. Then  $x = \sigma(a, x)$  gives

$$\sigma(a^{-1}, x) = \sigma(a^{-1}, \sigma(a, x)) = \sigma(a^{-1}a, x) = \sigma(e, x) = x.$$

Since  $\sigma(b, x) = x$ , we have  $\sigma(a^{-1}b, x) = \sigma(a^{-1}, \sigma(b, x)) = \sigma(a^{-1}, x) = x$ . Therefore,  $a^{-1}b \in G_x$ . Thus  $G_x$  is a subgroup of  $G$ .  $\square$

**Exercise 2.11.13.** Let  $\sigma : G \times X \rightarrow X$  be a left  $G$ -action on  $X$ . If  $f_\sigma : G \rightarrow S(X)$  is the group homomorphism induced by  $\sigma$ , then show that  $\text{Ker}(f_\sigma) = \bigcap_{x \in X} G_x$ , where  $G_x$  is the isotropy subgroup of  $x \in X$ .

**Corollary 2.11.14.** Let  $X$  be a non-empty set equipped with a left  $G$ -action  $\sigma : G \times X \rightarrow X$ . Let  $H$  be a normal subgroup of  $G$ . Then the  $G$ -action  $\sigma$  induces a left  $G/H$ -action  $\tilde{\sigma} : (G/H) \times X \rightarrow X$  making the following diagram commutative

$$\begin{array}{ccc} G \times X & \xrightarrow{\sigma} & X \\ \pi_H \times \text{Id}_X \downarrow & & \parallel \\ (G/H) \times X & \xrightarrow{\tilde{\sigma}} & X \end{array}$$

if and only if  $H \subseteq \bigcap_{x \in X} G_x$ , where  $G_x := \{g \in G : \sigma(g, x) = x\}$ ,  $\forall x \in X$ .

*Proof.* Let  $f_\sigma : G \rightarrow S(X)$  be the permutation representation of  $G$  in  $S(X)$  associated to the  $G$ -action  $\sigma$  on  $X$ . Note that  $\text{Ker}(f_\sigma) = \bigcap_{x \in X} G_x$ .

Let  $H$  be a normal subgroup of  $G$ . Let  $\pi_H : G \rightarrow G/H$  be the associated quotient group homomorphism. Suppose that  $H \subseteq \bigcap_{x \in X} G_x = \text{Ker}(f_\sigma)$ . Then by universal property of quotient, there exists a unique group homomorphism  $\widetilde{f}_\sigma : G/H \rightarrow S(X)$  such that  $\widetilde{f}_\sigma \circ \pi_H = f_\sigma$ . Then  $\widetilde{f}_\sigma$  induces a left  $G/H$ -action  $\tilde{\sigma} : (G/H) \times X \rightarrow X$  which sends  $(aH, x) \in (G/H) \times X$  to  $\tilde{\sigma}(aH, x) := \widetilde{f}_\sigma(aH)(x) = f_\sigma(a)(x) = \sigma(a, x) \in X$ .

Conversely, suppose that  $\tilde{\sigma} : (G/H) \times X \rightarrow X$  be a left  $G/H$ -action on  $X$  making the above diagram commutative. Let

$$f_{\tilde{\sigma}} : G/H \rightarrow S(X)$$

be the permutation representation of  $G/H$  into  $S(X)$  associated to  $\tilde{\sigma}$ . Then  $\sigma$  can be recovered from the group homomorphism

$$G \xrightarrow{\pi_H} G/H \xrightarrow{f_{\tilde{\sigma}}} S(X)$$

using the construction given in Lemma 2.11.4. From this, we have  $H \subseteq \text{Ker}(f_\sigma)$ .  $\square$

**Exercise 2.11.15.** Let  $\sigma : G \times X \rightarrow X$  be a left  $G$ -action on  $X$ . Given  $x \in X$  and  $a \in G$ , show that  $G_y = aG_xa^{-1}$ , where  $y = \sigma(a, x) \in X$ . Deduce that if  $\sigma$  is a transitive  $G$ -action on  $X$ , show that  $\text{Ker}(f_\sigma) = \bigcap_{a \in G} aG_xa^{-1}$ .

**Exercise 2.11.16.** Let  $X$  be a non-empty set. Let  $G$  be a subgroup of the symmetric group  $S(X)$  on  $X$ . Given  $\sigma \in G$  and  $x \in X$  we have  $\sigma G_x \sigma^{-1} = G_{\sigma(x)}$ . Deduce that if  $G$  acts transitively on  $X$ , then  $\bigcap_{\sigma \in G} \sigma G_x \sigma^{-1} = \{e\}$ .

**Corollary 2.11.17** (Generalized Cayley's Theorem). *Let  $H$  be a subgroup of  $G$ , and let  $X = \{aH : a \in G\}$  be the set of all distinct left cosets of  $H$  in  $G$ . Let  $S(X)$  be the symmetric group on the set  $X$ . Then there exists a group homomorphism  $\varphi : G \rightarrow S(X)$  such that  $\text{Ker}(\varphi) \subseteq H$ .*

*Proof.* Consider the map  $\sigma : G \times X \rightarrow X$  defined by

$$\sigma(a, bH) = (ab)H, \forall a \in G, bH \in X.$$

If  $bH = cH$ , for some  $b, c \in G$ , then given any  $a \in G$ , we have  $(ab)^{-1}(ac) = b^{-1}a^{-1}ac = b^{-1}c \in H$ . Therefore,  $\sigma$  is well-defined. Note that  $\sigma(e, bH) = bH$ ,  $\forall bH \in X$ , and  $\sigma(a_1, \sigma(a_2, bH)) = \sigma(a_1, a_2bH) = (a_1a_2b)H = \sigma(a_1a_2, bH)$ , for all  $a_1, a_2 \in G$  and  $bH \in X$ . Therefore,  $\sigma$  is a left  $G$ -action on  $X$ . Then  $\sigma$  give rise to the group homomorphism

$$f_\sigma : G \rightarrow S(X)$$

that sends  $a \in G$  to the map

$$\sigma(a, -) : X \rightarrow X, \quad x \mapsto \sigma(a, x).$$

Since  $\text{Ker}(f_\sigma) \subseteq G_x$ , for all  $x \in X$  by Exercise 2.11.13, taking  $x = H \in X$  we see that

$$G_H = \{a \in G : \sigma(a, H) = H\} = \{a \in G : a \in H\} = H,$$

and hence  $\text{Ker}(f_\sigma) \subseteq H$ .  $\square$

**Exercise 2.11.18.** Let  $H$  be a subgroup of  $G$ , and let  $X$  be the set of all left cosets of  $H$  in  $G$ . Let  $\sigma : G \times X \rightarrow X$  be the left  $G$ -action on  $X$  defined by  $\sigma(a, bH) = (ab)H$ ,  $\forall a, b \in G$ . Show that  $\sigma$  is a transitive action.

**Exercise 2.11.19.** Let  $G$  be a group and  $H$  a subgroup of  $G$  with  $[G : H] = n < \infty$ . Show that there is a normal subgroup  $K$  of  $G$  with  $K \subseteq H$  and  $[G : K] \leq n!$ .

**Corollary 2.11.20** (Cayley's Theorem). *Any group  $G$  is isomorphic to a subgroup of the symmetric group  $S(G)$  on  $G$ .*

*Proof.* Take  $H = \{e\}$  in Corollary 2.11.17. □

**Exercise 2.11.21.** Let  $G$  be a group of order  $2n$ , where  $n \geq 1$  is an odd integer. Show that  $G$  has a normal subgroup of order  $n$ .

*Solution:* By Cayley's theorem (Theorem 2.6.23)  $G$  is isomorphic to a subgroup, say  $H$ , of the symmetric group  $S(G)$  via the monomorphism  $\varphi : G \rightarrow S(G) \cong S_{2n}$  defined by sending  $a \in G$  to the bijective map  $\varphi_a : G \rightarrow G$  that sends  $b \in G$  to  $ab$ , for all  $b \in G$ . Since 2 divides  $|G| = 2n$ ,  $G$  has an element, say  $a \in G$ , of order 2 by Exercise 2.2.37. Since for any  $b \in G$  we have  $\varphi_a(b) = ab$  and  $\varphi_a(ab) = a^2b = eb = b$ , we see that  $\varphi_a \in S(G)$  is a product of transpositions of the form  $(b \ ab)$ . Since  $|G| = 2n$ , the number of transpositions appearing in the factorization of  $\varphi_a$  is  $n$ , an odd number. So  $\varphi_a$  is an odd permutation. This shows that the subgroup  $H := \varphi(G)$  contains an odd permutation. Define a map

$$f : H \rightarrow \{-1, 1\}$$

by sending  $\sigma \in H$  to

$$f(\sigma) := \begin{cases} 1, & \text{if } \sigma \text{ is an even permutation,} \\ -1, & \text{if } \sigma \text{ is an odd permutation.} \end{cases}$$

Note that  $f$  is a surjective group homomorphism, and hence by first isomorphism theorem (Theorem 2.9.3) we have

$$H/\text{Ker}(f) \cong \{-1, 1\}.$$

Then we have

$$2 = |\{-1, 1\}| = |H/\text{Ker}(f)| = \frac{|H|}{|\text{Ker}(f)|} = \frac{2n}{|\text{Ker}(f)|}.$$

Therefore,  $\text{Ker}(f)$  is a normal subgroup of  $H$  of order  $|\text{Ker}(f)| = n$ . Since  $G \cong H$  via  $\varphi$ , taking inverse image of  $\text{Ker}(f) \subseteq H$  along the isomorphism  $\varphi$  we get a required normal subgroup of  $G$  of order  $n$ . □

**Corollary 2.11.22.** *Let  $G$  be a finite group of order  $n$ . Let  $p > 0$  be a smallest prime number that divides  $n$ . If  $H$  is subgroup of  $G$  with  $[G : H] = p$ , then  $H$  is normal in  $G$ .*

*Proof.* Let  $H$  be a subgroup of index  $p$  in  $G$ . Let  $X := \{aH : a \in G\}$  be the set of all distinct left cosets of  $H$  in  $G$ . Then  $|X| = p$ . Let  $f : G \rightarrow S(X)$  be the map that sends  $a \in G$  to

$$f(a) : X \rightarrow X, \quad bH \mapsto (ab)H.$$

Then  $f$  is a group homomorphism. Then  $K := \text{Ker}(f) \subseteq H$  by Corollary 2.11.17, and  $[G : K] = [G : H] \cdot [H : K] = pk$ , where  $k := [H : K]$ . Since  $|X| = [G : H] = p$ , the quotient group  $G/K$  is isomorphic to a subgroup of the symmetric group  $S_p$  by first isomorphism theorem (see Theorem 2.9.3). Then by Lagrange's theorem  $pk = |G/K|$  divides  $|S_p| = p!$ . Then  $k$  divides  $(p-1)!$ . Since  $k$  is a divisor of  $n$  and  $p$  is the smallest prime divisor of  $n$ , unless  $k = 1$ , any prime divisor of  $k$  must be greater than or equal to  $p$ . But since  $k$  divides  $(p-1)!$ , any prime divisor of  $k$  is less than  $p$ . Thus we get a contradiction unless  $k = 1$ . Therefore,  $[H : K] = k = 1$ , and so  $H = K = \text{Ker}(f)$ . Thus  $H$  is a normal subgroup of  $G$ . □

**Warning:** The above Corollary 2.11.22 does not ensure existence of a subgroup  $H$  of  $G$  of index smallest prime factor of  $|G|$ .

**Exercise 2.11.23.** Let  $G$  be a finite group of order  $p^n$ , for some prime number  $p$  and integer  $n > 0$ . Show that every subgroup of  $G$  of index  $p$  is normal in  $G$ . Deduce that every group of order  $p^2$  has a normal subgroup of order  $p$ .

**Exercise 2.11.24.** Let  $G$  be a non-abelian group of order 6. Show that  $G$  has a non-normal subgroup of order 2. Use this to classify groups of order 6. (*Hint*: Produce a monomorphism into  $S_3$ ).

**Proposition 2.11.25.** Let  $\sigma : G \times X \rightarrow X$  be a left  $G$ -action on  $X$ . Fix  $x \in X$ , and let  $G/G_x = \{aG_x : a \in G\}$  be the set of all distinct left cosets of  $G_x$  in  $G$ . Then the map  $\varphi : G/G_x \rightarrow \text{Orb}_G(x)$  defined by  $\varphi(aG_x) = \sigma(a, x)$ ,  $\forall a \in G$ , is a well-defined bijective map. Consequently,  $[G : G_x] = |\text{Orb}_G(x)|$ .

*Proof.* Let  $a, b \in G$  be such that  $aG_x = bG_x$ . Then  $a^{-1}b \in G_x$ , and so  $\sigma(a^{-1}b, x) = x$ . Applying  $\sigma(a, -)$  both sides, we have  $\sigma(b, x) = \sigma(a, \sigma(a^{-1}b, x)) = \sigma(a, x)$ . Therefore, the map  $\varphi$  is well-defined. To show that  $\varphi$  is injective, suppose that  $\sigma(a, x) = \sigma(b, x)$ , for some  $a, b \in G$ . Then  $\sigma(a^{-1}b, x) = \sigma(a^{-1}, \sigma(b, x)) = \sigma(a^{-1}, \sigma(a, x)) = \sigma(e, x) = x$ . Therefore,  $a^{-1}b \in G_x$ , and hence  $aG_x = bG_x$ . Thus  $\varphi$  is injective. To show  $\varphi$  is surjective, note that  $\sigma(a, x) = \varphi(aG_x)$ , for all  $a \in G$ . Therefore,  $\varphi$  is bijective.  $\square$

**Corollary 2.11.26 (Class Equation).** Let  $\sigma : G \times X \rightarrow X$  be a left  $G$ -action on a non-empty finite set  $X$ , and let  $\mathcal{O}$  be a subset of  $X$  containing exactly one element from each  $G$ -orbits in  $X$ . Then we have

$$|X| = \sum_{x \in \mathcal{O}} [G : G_x].$$

*Proof.* Since  $X = \bigsqcup_{x \in \mathcal{O}} \text{Orb}_G(x)$ , the result follows from Proposition 2.11.25.  $\square$

**Exercise 2.11.27.** Let  $G$  be a group. Let  $H$  be a subgroup of  $G$  such that  $|H| = 11$  and  $[G : H] = 4$ . Show that  $H$  is a normal subgroup of  $G$ .

**Exercise 2.11.28.** Fix  $n \in \mathbb{N}$ . Show that the map  $\sigma : \text{GL}_n(\mathbb{R}) \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  defined by

$$\sigma(A, v) = Av, \forall A \in \text{GL}_n(\mathbb{R}), v = (v_1, \dots, v_n)^t \in \mathbb{R}^n,$$

is a left  $\text{GL}_n(\mathbb{R})$ -action on  $\mathbb{R}^n$ . Is  $\sigma$  transitive? Find the set of all  $\text{GL}_n(\mathbb{R})$ -orbits in  $\mathbb{R}^n$ .

**Exercise 2.11.29.** Let  $\sigma : G \times G \rightarrow G$  be the left  $G$ -action on itself given by

$$\sigma(a, b) = aba^{-1}, \forall a, b \in G.$$

If  $f_\sigma : G \rightarrow S(G)$  is the permutation representation of  $G$  associated to  $\sigma$ , show that  $\text{Ker}(f_\sigma) = Z(G)$ .

**Theorem 2.11.30 (Burnside's Theorem).** Let  $G$  be a finite group acting from the left on a non-empty finite set  $X$ . Then the number of distinct  $G$ -orbits in  $X$  is equal to

$$\frac{1}{|G|} \sum_{a \in G} F(a),$$

where  $F(a) = \#\{x \in X : ax = x\}$ , the number of elements of  $X$  fixed by  $a$ .

*Proof.* Let  $T := \{(a, x) \in G \times X : ax = x\}$ . Note that  $|T| = \sum_{a \in G} F(a)$ . Also  $|T| = \sum_{x \in X} |G_x|$ , where  $G_x$  is the stabilizer of  $x \in X$ . Let  $\{x_1, \dots, x_n\}$  be the subset of  $X$  consisting of exactly one element from each of the  $G$ -orbits in  $X$ . Note that two elements  $x$  and  $y$  of  $X$  are in the

same  $G$ -orbit if and only if  $\text{Orb}_G(x) = \text{Orb}_G(y)$ . Since  $|G|/|G_x| = [G : G_x] = |\text{Orb}_G(x)|$ , we conclude that  $|G_x| = |G_y|$  whenever  $x$  and  $y$  are in the same  $G$ -orbit. Then we have

$$\begin{aligned} \sum_{a \in G} F(a) &= |T| = \sum_{x \in X} |G_x| \\ &= \sum_{i=1}^n |\text{Orb}_G(x_i)| |G_{x_i}| \\ &= \sum_{i=1}^n |G| = n|G|, \end{aligned}$$

and hence  $n = \frac{1}{|G|} \sum_{a \in G} F(a)$ . This completes the proof.  $\square$

## 2.12 Conjugacy Action & Class Equations

Let  $G$  be a group. Consider the map

$$(2.12.1) \quad \sigma : G \times G \rightarrow G, (a, b) \mapsto aba^{-1}.$$

Note that  $\sigma$  is a left action of  $G$  on itself, known as the conjugation action. Given  $a \in G$ , its  $\sigma$ -stabilizer

$$G_a = \{g \in G : gag^{-1} = a\} = \{g \in G : ga = ag\}.$$

is a subgroup of  $G$ , called the *centralizer* or the *normalizer* of  $a$  in  $G$ . The equivalence relation  $\sim_\sigma$  on  $G$  induced by the conjugation action of  $G$  on itself is known as the *conjugate* relation on  $G$ . An element  $b \in G$  is said to be a *conjugate* of  $a \in G$  if there exists  $g \in G$  such that  $b = gag^{-1}$ . Given  $a \in G$ , its  $G$ -orbit

$$(2.12.2) \quad \text{Orb}_G(a) = \{gag^{-1} : g \in G\}$$

consists of all conjugates of  $a$  in  $G$ , and is called the *conjugacy class* of  $a$  in  $G$ .

**Definition 2.12.3.** A partition of an integer  $n \geq 1$  is a finite sequence of positive integers  $(n_1, \dots, n_r)$  such that  $n_1 \geq \dots \geq n_r$  and  $\sum_{j=1}^r n_j = n$ .

**Exercise 2.12.4.** Fix an integer  $n \geq 2$ . Show that the number of conjugacy classes in  $S_n$  is the number of partitions of  $n$ .

*Solution:* Let  $\mathcal{C} = \{C_1, \dots, C_k\}$  be the set of all distinct conjugacy classes in  $S_n$ . Let  $\mathcal{P}_n$  be the set of all partitions of  $n$ . Define a map  $t : \mathcal{C} \rightarrow \mathcal{P}_n$  by sending  $C_i \in \mathcal{C}$  to the cycle type of an element of  $C_i$ , for all  $i$ . Since two elements of  $S_n$  are conjugate in  $S_n$  if and only if they have the same cycle type by Theorem 2.5.22, the map  $t$  is well-defined and injective. Given a partition  $(n_1, \dots, n_r)$  of  $n$ , we have a permutation  $\sigma = (1 \dots n_1) \circ \dots \circ (n_1 + \dots + n_{r-1} + 1 \dots n_1 + \dots + n_r) \in S_n$  whose cycle type is precisely  $(n_1, \dots, n_r)$ . Therefore,  $t$  is surjective, and hence is bijective, as required.  $\square$

More generally,  $G$  acts on its power set  $X := \mathcal{P}(G)$  by conjugation:

$$(2.12.5) \quad \sigma : G \times \mathcal{P}(G) \rightarrow \mathcal{P}(G), (a, S) \mapsto aSa^{-1},$$

where

$$aSa^{-1} := \begin{cases} \{aga^{-1} \in G : g \in S\}, & \text{if } S \neq \emptyset, \text{ and} \\ \emptyset, & \text{if } S = \emptyset. \end{cases}$$



Two non-empty subset  $S$  and  $T$  of  $G$  are said to be conjugates if there exists  $a \in G$  such that  $T = aSa^{-1}$ . Given a subset  $S \subseteq G$ , its stabilizer

$$(2.12.6) \quad N_G(S) := \{a \in G : aSa^{-1} = S\}$$

for the conjugation action in (2.12.5), is a subgroup of  $G$ , known as the *normalizer* of  $S$  in  $G$ . Then we have the following.

**Corollary 2.12.7.** *Let  $S$  be a non-empty subset of  $G$ . Then the number of distinct conjugates of  $S$  in  $G$  is the index  $[G : N_G(S)]$ . In particular, the number of distinct conjugates of an element  $a \in G$  is  $[G : C_G(a)]$ , where  $C_G(a)$  is the centralizer of  $a$  in  $G$ .*

*Proof.* Follows from Proposition 2.11.25. □

**Exercise 2.12.8.** Let  $\sigma = (k_1 \ \cdots \ k_r) \in S_n$  be a  $r$ -cycle in  $S_n$ . Let  $I_n \setminus \sigma := I_n \setminus \{k_1, \dots, k_r\} \subset I_n$ , and let

$$S(I_n \setminus \sigma) := \left\{ \tau \in S_n : \tau|_{\{k_1, \dots, k_r\}} = \text{Id}_{\{k_1, \dots, k_r\}} \right\}.$$

- (i) Show that  $S(I_n \setminus \sigma)$  is a subgroup of  $S_n$ .
- (ii) Show that  $|C_{S_n}(\sigma)| = r(n-r)!$ .
- (iii) Deduce that  $C_{S_n}(\sigma) = \{\sigma^i \tau \in S_n : \tau \in S(I_n \setminus \sigma)\}$ . (*Hint:* Note that  $\sigma$  commutes with  $e, \sigma, \dots, \sigma^{r-1}$ , and with all  $\tau \in S_n$  whose cycles are disjoint from that of  $\sigma$  (precisely elements of  $S(I_n \setminus \sigma)$ ). Then use part (ii).)
- (iv) Compute  $C_{S_7}(\sigma)$ , where  $\sigma = (1 \ 2 \ 3) \in S_7$ .

**Exercise 2.12.9.** Let  $G$  be a group and  $S$  a non-empty subset of  $G$ . If  $H$  is the subgroup of  $G$  generated by  $S$ , show that  $N_G(S) \leq N_G(H)$ .

Note that given  $a \in G$  we have  $C_G(a) = G$  if and only if  $a \in Z(G)$ . Therefore, we have the following.

**Theorem 2.12.10** (Class Equation). *Let  $G$  be a finite group, and let  $\{a_1, \dots, a_n\}$  be the subset of  $G$  consisting of exactly one element from each conjugacy class that are not contained in  $Z(G)$ . Then we have*

$$|G| = |Z(G)| + \sum_{i=1}^n [G : C_G(a_i)].$$

*Proof.* Follows from Corollary 2.11.26 by taking  $X = G$  and  $\sigma$  to be the conjugation action of  $G$  on itself. □

**Corollary 2.12.11.** *Let  $G$  be a group of order  $p^n$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Then  $G$  has non-trivial center.*

*Proof.* The class equation (see Theorem 2.12.10) for the conjugacy action of  $G$  on itself gives

$$p^n = |G| = |Z(G)| + \sum_{i=1}^r [G : C_G(a_i)],$$

where  $\{a_1, \dots, a_n\}$  is a subset consisting of exactly one element from each conjugacy class that are not in the center  $Z(G)$ . Since  $C_G(a_i)$  is a subgroup of  $G$ , by Lagrange's theorem  $|C_G(a_i)|$  divides  $|G| = p^n$ , and hence its index  $[G : C_G(a_i)] = |G|/|C_G(a_i)|$  is of the form  $p^{n_i}$ , for some  $n_i \in \mathbb{N} \cup \{0\}$ . Since  $a_i \notin Z(G)$ , we have  $C_G(a_i) \neq G$ , and so  $n_i \geq 1$ , for all  $i$ . Since  $Z(G)$  is a subgroup of  $G$ , we have  $|Z(G)| \geq 1$ . Then by above class equation we see that  $|Z(G)| = p^n - \sum_{i=1}^r p^{n_i}$  is divisible by  $p$ . Therefore,  $Z(G) \neq \{e\}$ . □

**Corollary 2.12.12.** *Let  $G$  be a group of order  $p^2$ , where  $p > 0$  is a prime number. Then  $G$  is isomorphic to either  $\mathbb{Z}_{p^2}$  or  $\mathbb{Z}_p \times \mathbb{Z}_p$ .*

*Proof.* Since  $Z(G) \neq \{e\}$  by Corollary 2.12.11, we see that  $G/Z(G)$  has order  $p$  or 1, and hence is cyclic. Then  $G$  is abelian by Exercise 2.8.22. If  $G$  has an element of order  $p^2$ , then  $G$  is cyclic. Suppose that  $G$  has no element of order  $p^2$ . Then every non-neutral element of  $G$  has order  $p$ . Fix an  $a \in G \setminus \{e\}$ , and take  $b \in G \setminus \langle a \rangle$ . Then we have  $|\langle a, b \rangle| > |\langle a \rangle| = p$ , and hence  $\langle a, b \rangle = G$ . Since both  $a$  and  $b$  has order  $p$ , it follows that  $\langle a \rangle \times \langle b \rangle \cong \mathbb{Z}_p \times \mathbb{Z}_p$ . Note that both  $H := \langle a \rangle$  and  $K := \langle b \rangle$  are normal subgroups of  $G$  of order  $p$ . Since  $H \cap K$  is a subgroup of both  $H$  and  $K$ ,  $|H \cap K|$  is either  $p$  or 1 by Lagrange's theorem (Theorem 2.7.5). If  $|H \cap K| = p$ , then  $K = H \cap K = H$ , which contradicts the choice of  $b \in G \setminus H$ . Therefore,  $H \cap K = \{e\}$ . Since  $HK$  is a subgroup of  $G$  by Theorem 2.4.3 with

$$|HK| = \frac{|H| \cdot |K|}{|H \cap K|} = p^2 = |G|$$

by Lemma 2.4.7, we have  $G = HK$ . Then  $G \cong H \times K$  by Theorem 2.10.18.  $\square$

**Proposition 2.12.13.** *Let  $G$  be a finite abelian group of order  $n \geq 2$ . If  $p > 0$  is a prime number dividing  $n$ , then  $G$  has an element of order  $p$ .*

*Proof.* We prove this by induction on  $n = |G|$ . The case  $n = 2$  is trivial. Assume that  $n > 2$ , and the result holds for any abelian group of order  $r$  with  $2 \leq r < n$ . Let  $a \in G \setminus \{e\}$  be given. If  $\langle a \rangle = G$ , then we are done by Proposition 2.3.14. Assume that  $H := \langle a \rangle$  is a proper non-trivial subgroup of  $G$ . Let  $m := \text{ord}(a)$ . Then  $1 < m < n$ . If  $p \mid m$ , then by induction hypothesis  $H$  has an element, say  $b$ , of order  $p$ , and we are done. Assume that  $p \nmid m$ . Since  $G$  is abelian,  $H$  is a normal subgroup of  $G$ . Then  $p$  divides the order of the quotient group  $G/H$ . Since  $|G/H| = n/m < n$ , by induction hypothesis  $G/H$  has an element, say  $bH \in G/H$ , of order  $p$ . Then  $b^p H = (bH)^p = H$  in  $G/H$ , and so  $b^p \in H$ . Since  $H = \langle a \rangle$  is a cyclic group of order  $m$ , we have  $(b^m)^p = (b^p)^m = e$ . Then  $\text{ord}(b^m) \mid p$ . Since  $p$  is a prime number, either  $b^m = e$  or  $\text{ord}(b^m) = p$ . If  $b^m = e$ , then  $(bH)^m = b^m H = eH = H$ , and so  $p = \text{ord}(bH) \mid m$ . This contradicts our assumption that  $p \nmid m$ . Therefore,  $b^m \neq e$ , and hence  $\text{ord}(b^m) = p$ .  $\square$

**Theorem 2.12.14 (Cauchy).** *Let  $G$  be a finite group of order  $n$ . Then for each prime number  $p > 0$  dividing  $n$ ,  $G$  has an element of order  $p$ .*

*Proof.* Fix a prime number  $p > 0$  that divides  $n$ . The case  $n = 2$  is trivial. Suppose that  $n > 2$ , and the statement holds for any finite group of order  $r$  with  $2 \leq r < n$ . The class equation for  $G$  associated to the conjugacy action of  $G$  on itself is given by

$$(2.12.15) \quad |G| = |Z(G)| + \sum_{i=1}^r [G : C_G(a_i)],$$

where  $\{a_1, \dots, a_r\}$  is the subset of  $G$  consisting of exactly one element from each  $G$ -orbits of that does not intersect  $Z(G)$ . Since  $a \in Z(G)$  if and only if  $C_G(a) = G$ , we see that  $|C_G(a_i)| < n$ , for all  $i \in \{1, \dots, r\}$ . If  $p \mid |C_G(a_i)|$ , for some  $i \in \{1, \dots, r\}$ , then by induction hypothesis  $C_G(a_i) \subseteq G$  has an element of order  $p$ , and we are done. Suppose that  $p \nmid |C_G(a_i)|$ ,  $\forall i \in \{1, \dots, r\}$ . Since  $p \mid n = |G|$  and  $|G| = |C_G(a_i)|[G : C_G(a_i)]$ , we see that  $p \mid [G : C_G(a_i)]$ ,  $\forall i \in \{1, \dots, r\}$ . Since  $Z(G)$  is a subgroup of  $G$ ,  $|Z(G)| \geq 1$ . Then from class equation above, we see that  $p$  divides  $|Z(G)|$ . Since  $Z(G)$  is abelian, it contains an element of order  $p$  by Proposition 2.12.13. This completes the proof.  $\square$

As an immediate corollary, we have the following result, known as the *converse of Lagrange's theorem for finite abelian groups*.

**Corollary 2.12.16.** *Let  $G$  be a finite abelian group of order  $n$ . Let  $m > 0$  be an integer that divides  $n$ . Then  $G$  has a subgroup of order  $m$ .*

*Proof.* The cases  $n = 2$  and  $m = 1$  are trivial. So we assume that  $m > 1$  and  $n > 2$ , and we prove it by induction on  $n$ . Suppose that the statement holds for any finite abelian group of order  $r$  with  $2 \leq r < n$ . Let  $G$  be an abelian group of order  $n$ . Since  $m > 1$ , there is a prime number, say  $p \in \mathbb{N}$ , such that  $p \mid m$ . Then  $m = pk$ , for some  $k \in \mathbb{N}$ . Then by Cauchy's theorem (Theorem 2.12.14)  $G$  has a subgroup, say  $H$ , of order  $p$ . Since  $G$  is abelian, that  $H$  is normal in  $G$ . Then the quotient group  $G/H$  exists and we have  $1 \leq |G/H| = n/p < n$ . Since  $m \mid n$ , we have  $n = m\ell$ , for some  $\ell \in \mathbb{N}$ . Then

$$|G/H| = \frac{n}{p} = \frac{m\ell}{p} = \frac{pk\ell}{p} = k\ell.$$

Since  $G/H$  is abelian group with  $|G/H| < n$  and  $k \mid |G/H|$ , by induction hypothesis  $G/H$  has a subgroup, say  $S$ , of order  $k$ . Now  $S = K/H$ , for some subgroup  $K$  of  $G$  containing  $H$  by Exercise 2.9.26. Since  $|K| = |S| \cdot |H| = kp = m$ , that  $K$  is a required subgroup of  $G$  of order  $m$ . This completes the proof.  $\square$

### 2.12.1 $p$ -groups

**Definition 2.12.17 ( $p$ -group).** Let  $p \in \mathbb{N}$  be a prime number. A group  $G$  is said to be a  $p$ -group if every element of  $G$  has order equal to a power of  $p$ . A subgroup  $H$  of  $G$  is called a  $p$ -subgroup of  $G$  if  $H$  is a  $p$ -group.

**Example 2.12.18.**  $D_4$  and  $K_4$  are 2-groups.

**Example 2.12.19.** Given a prime number  $p > 0$ , let

$$\mathbb{Z}_{(p)} := \left\{ \frac{m}{p^n} \in \mathbb{Q} : m, n \in \mathbb{Z} \right\}.$$

Clearly  $\mathbb{Z}_{(p)}$  is a non-empty subset of  $\mathbb{Q}$ . Since given  $m/p^n, k/p^\ell \in \mathbb{Z}_{(p)}$ , we have

$$\frac{m}{p^n} - \frac{k}{p^\ell} = \frac{mp^\ell - np^n}{p^{n+\ell}} \in \mathbb{Z}_{(p)},$$

we conclude that  $\mathbb{Z}_{(p)}$  is a subgroup of  $\mathbb{Q}$ . Note that  $\text{ord}(m/p^n)$  is a power of  $p$ , and hence  $\mathbb{Z}_{(p)}$  is a  $p$ -group.

**Proposition 2.12.20.** *A finite group  $G$  is a  $p$ -group if and only if  $|G| = p^n$ , for some  $n \in \mathbb{N}$ .*

*Proof.* If  $|G| = p^n$ , for some  $n \in \mathbb{N}$ , then given  $a \in G$ ,  $\text{ord}(a) \mid p^n$  by Lagrange's theorem (Theorem 2.7.5), and hence  $\text{ord}(a) = p^r$ , for some  $r \in \{1, \dots, n\}$ , since  $p$  is a prime number.

Conversely suppose that  $G$  is a finite  $p$ -group. If  $|G| \neq p^n$ , for all  $n \in \mathbb{N} \cup \{0\}$ , then there exists a prime number  $q \neq p$  such that  $q \mid |G|$ . Then by Cauchy's theorem  $G$  has an element of order  $q$ , which is not of the form  $p^n$ , for any  $n \in \mathbb{N}$ . This contradicts our assumption that  $G$  is a  $p$ -group. This completes the proof.  $\square$

**Lemma 2.12.21.** *Subgroup of a  $p$ -group is a  $p$ -group.*

*Proof.* Follows from the definition.  $\square$

**Lemma 2.12.22.** *Let  $G$  be a group (not necessarily finite), and  $p > 0$  a prime number. Then any  $p$ -subgroup of  $G$  is contained in a maximal  $p$ -subgroup of  $G$ .*

*Proof.* Let  $P$  be a  $p$ -subgroup of  $G$ . Let  $\mathcal{P}$  be the set of all  $p$ -subgroups of  $G$  containing  $P$ . Given  $P, Q \in \mathcal{P}$  we define  $P \leq Q$  if  $P \subseteq Q$ . Clearly this is a partial order relation on  $\mathcal{P}$ . Given a chain  $(P_n)_{n \geq 0}$  of elements from  $\mathcal{P}$  with  $P = P_0 \leq P_1 \leq \dots$ , the subset  $P := \bigcup_{n \geq 0} P_n$  is a  $p$ -subgroup of  $G$  (verify!), and hence is an element of  $\mathcal{P}$ . Then by Zorn's lemma  $\mathcal{P}$  has a maximal element, say  $P_{\max} \in \mathcal{P}$ . This completes the proof.  $\square$

**Proposition 2.12.23.** *Any finite non-trivial  $p$ -group have non-trivial center.*

*Proof.* Let  $G$  be a  $p$ -group of order  $p^n$ , for some prime number  $p > 0$  and positive integer  $n > 0$ . Then the class equation for the conjugacy action of  $G$  on itself gives

$$|G| = |Z(G)| + \sum_{a \in \mathcal{O} \setminus Z(G)} [G : C_G(a)],$$

where  $\mathcal{O}$  is a subset of  $G$  consisting of exactly one element from each  $G$ -orbits. Since  $C_G(a) = G$  if and only if  $a \in Z(G)$ , we see that  $[G : C_G(a)] > 1$  for all  $a \in \mathcal{O} \setminus Z(G)$ . Since  $|G| = p^n$ , it follows from Lagrange's theorem that  $p$  divides  $[G : C_G(a)]$ ,  $\forall a \in \mathcal{O} \setminus Z(G)$ . Then from the class equation above we see that  $p$  divides  $|Z(G)|$ . Since  $|Z(G)| \geq 1$ , it follows that  $Z(G) \neq \{e\}$ .  $\square$

**Corollary 2.12.24.** *Let  $p > 0$  be a prime number. Then every group of order  $p^2$  is abelian.*

*Proof.* Let  $G$  be a group of order  $p^2$ . Then by Proposition 2.12.23 above,  $Z(G) \neq \{e\}$ . Then  $|Z(G)| \in \{p, p^2\}$  by Lagrange's theorem. If  $|Z(G)| = p$ , then the quotient group  $G/Z(G)$  has order  $p$ , and hence is cyclic by Corollary 2.7.9. Then  $G$  is abelian by Exercise 2.8.22, which is a contradiction. Therefore,  $|Z(G)| = p^2 = |G|$ , and hence  $G = Z(G)$ . Therefore,  $G$  is abelian.  $\square$

**Lemma 2.12.25.** *Let  $G$  be a group of order  $p^n$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Let  $X$  be a non-empty finite set admitting a left  $G$ -action. Let*

$$X_0 := \{x \in X : ax = x, \forall a \in G\}$$

*be the subset of  $X$  consisting of elements with singleton  $G$ -orbits. Then  $|X| \equiv |X_0| \pmod{p}$ . In particular, if  $p \nmid |X|$ , there exists  $x \in X$  with singleton  $G$ -orbit.*

*Proof.* The class equation for the left  $G$ -action on  $X$  gives

$$|X| = |X_0| + \sum_{x \in \mathcal{O} \setminus X_0} [G : G_x],$$

where  $\mathcal{O}$  is the subset of  $X$  consisting of exactly one element from each  $G$ -orbits of  $X$ . Since  $[G : G_x] = |\text{Orb}_G(x)| > 1$ , for all  $x \in \mathcal{O} \setminus X_0$ , and  $|G| = p^n$ , we conclude that  $p$  divides  $[G : G_x]$ , for all  $x \in \mathcal{O} \setminus X_0$ . Then the result follows by reducing the class equation above modulo  $p$ . If  $p \nmid |X|$ , then  $|X_0| \neq 0 \pmod{p}$ , and hence the second part follows.  $\square$

**Corollary 2.12.26.** *Let  $G$  be a finite group having a subgroup  $H$  of order  $p^n$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Then  $[G : H] \equiv_p [N_G(H) : H]$ . In particular, if  $p \mid [G : H]$ , then  $N_G(H) \neq H$ .*

*Solution:* Take  $X = \{aH : a \in G\}$  to be the set of all left cosets of  $H$  in  $G$ . Then  $H$  acts on  $X$  by

$$\sigma : H \times X \rightarrow X, (h, aH) \mapsto (ha)H.$$

Note that  $\sigma$  is a well-defined map and is a left  $H$ -action on  $X$ . Moreover the subset of  $X$  consisting of singleton  $H$ -orbits is given by

$$\begin{aligned} X_0 &= \{aH \in X : \sigma(h, aH) = aH, \forall h \in H\} \\ &= \{aH \in X : a^{-1}ha \in H, \forall h \in H\} \\ &= \{aH \in X : a \in N_G(H)\}, \end{aligned}$$

we have  $|X_0| = [N_G(H) : H]$ . Since  $|X| = [G : H]$ , the result follows from Lemma 2.12.25.  $\square$

## 2.13 Simple Groups

**Definition 2.13.1.** A group is said to be *simple* if it has no non-trivial proper normal subgroup.

**Example 2.13.2.** Any group of prime order is simple (c.f. Lagrange's theorem).

**Lemma 2.13.3.** A finite abelian group  $G$  is simple if and only if  $|G|$  is a prime number.

*Proof.* If  $|G| = p$ , for some prime number, then its only subgroups are  $\{e\}$  and  $G$ , and hence  $G$  is simple in this case. To see the converse, note that if  $|G|$  is composite, then  $|G| = pk$ , for some prime number  $p$  and an integer  $k > 1$ . Then by Cauchy's theorem (Theorem 2.12.14)  $G$  has an element, say  $a \in G$ , of order  $p$ . Since  $G$  is abelian, the cyclic subgroup  $H := \langle a \rangle$  of  $G$  is normal in  $G$ . Since  $1 < |H| = p < |G|$ , it follows that  $H$  is a non-trivial proper normal subgroup of  $G$ . Thus  $G$  is not simple.  $\square$

**Exercise 2.13.4.** Let  $G$  be a finite group of order  $pq$ , where  $p$  and  $q$  are primes (not necessarily distinct). Show that  $G$  is not simple.

*Solution:* If  $p = q$ , then  $|G| = p^2$ , and so  $G$  is abelian by Corollary 2.12.24. Then  $G$  is not simple by Lemma 2.13.3. If  $p \neq q$ , without loss of generality we assume that  $p > q$ . Then by Cauchy's theorem  $G$  has a subgroup, say  $H$ , of order  $p$ . To show  $G$  is not simple, it suffices to show that  $H$  is normal. If possible suppose that there exists  $a \in G$  such that  $aHa^{-1} \neq H$ . Since both  $H$  and  $K_a := aHa^{-1}$  are subgroups of  $G$  of order  $p$ , their intersection  $H \cap K_a$  is a subgroup (see Lemma 2.2.18) of order 1 or  $p$  by Lagrange's theorem (Theorem 2.7.5). Since  $H \neq K_a$  by assumption,  $|H \cap K_a| = 1$ . Then the subset  $HK_a \subseteq G$  has cardinality

$$|HK_a| = \frac{|H| \cdot |K_a|}{|H \cap K_a|} = p^2 > pq = |G|,$$

which is a contradiction. Therefore,  $aHa^{-1} = H$ ,  $\forall a \in G$ , and hence  $H$  is normal in  $G$ .  $\square$

**Exercise 2.13.5.** Let  $G$  be an abelian group having finite subgroups  $H$  and  $K$  of orders  $m$  and  $n$ , respectively. Show that  $G$  has a subgroup of order  $d := \text{lcm}(m, n)$ .

*Solution.* Since  $G$  is abelian, both  $H$  and  $K$  are normal in  $G$ , and hence  $HK$  is a subgroup of  $G$  of order at most  $|H| \cdot |K| = mn$ . Since  $H$  and  $K$  are subgroups of  $HK$ , by Lagrange's theorem both  $m$  and  $n$  divides  $|HK|$ , and hence  $d := \text{lcm}(m, n)$  divides  $|HK|$ . Since  $G$  is abelian, so is its subgroup  $HK$ . Then by Corollary 2.12.16  $HK$  has a subgroup, say  $V$  of order  $d$ . Since  $V$  is also a subgroup of  $G$ , we are done.  $\square$

**Exercise 2.13.6.** Let  $G$  be a non-abelian group of order  $p^3$ , where  $p$  is a prime number. Show that  $|Z(G)| = p$ .

*Solution:* Since  $G$  has order  $p^3$ , it has non-trivial center. Since  $G$  is non-abelian, so  $Z(G) \neq G$ . Then by Lagrange's theorem  $Z(G)$  has order  $p$  or  $p^2$ . If  $|Z(G)| = p^2$ , then  $G/Z(G)$  has order

$p$ , and hence is a cyclic group. Then  $G$  is abelian by Exercise 2.8.22, which is a contradiction. Therefore,  $|Z(G)| = p$ .  $\square$

**Exercise 2.13.7.** Let  $G$  be a finite abelian group. Let  $n \in \mathbb{N}$  be such that  $n \mid |G|$ . Show that the number of solutions of the equation  $x^n = e$  in  $G$  is a multiple of  $n$ .

*Solution:* The set of all solutions of  $x^n = e$  in  $G$  is given by

$$H := \{a \in G : a^n = e\}.$$

Since  $e^n = e$ , we see that  $H \neq \emptyset$ . Let  $a, b \in H$  be given. Since  $G$  is abelian, we have  $(a^{-1}b)^n = (a^n)^{-1}b^n = e^{-1}e = e$ , and so  $a^{-1}b \in H$ . Therefore,  $H$  is a subgroup of  $G$ . Since  $G$  is a finite abelian group and  $n \mid |G|$ , by Corollary 2.12.16  $G$  has a subgroup, say  $K$  of order  $n$ . Then by Corollary 2.7.7 we have  $a^n = e$ ,  $\forall a \in K$ , and hence  $K \subseteq H$ . Since  $|K| = n$ , by Lagrange's theorem we have  $n \mid |H|$ .  $\square$

**Exercise 2.13.8.** Let  $G$  be a group of order  $p^n$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Let  $H$  be a subgroup of  $G$  of order  $p^{n-1}$ . Show that  $H$  is normal in  $G$ .

*Solution:* Follows from Corollary 2.11.22.  $\square$

**Exercise 2.13.9.** Show that  $N := \{e, (1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)\} \subset A_4$  is the unique subgroup of order 4 in  $A_4$ , and hence is normal in  $A_4$ . Conclude that  $A_4$  is not simple.

### 2.13.1 Simplicity of $A_n$ , for $n \geq 5$

Next we show that  $A_n$  is simple, for all  $n \geq 5$ . We begin with some useful observations.

**Lemma 2.13.10.** Fix an integer  $n \geq 5$ , and let  $H$  be a normal subgroup of  $A_n$ . If  $H$  contains a 3-cycle, then  $H = A_n$ .

*Proof.* Suppose that  $H$  contains a 3-cycle, say  $\sigma = (a\ b\ c) \in H$ . Since  $A_n$  is generated by 3-cycles, it suffices to show that any 3-cycle is contained in  $H$ . Let  $\tau = (u\ v\ w)$  be any 3-cycle. Let  $\pi \in S_n$  be such that

$$\pi(a) = u, \pi(b) = v \text{ and } \pi(c) = w.$$

Then by Proposition 2.5.12 we have

$$\pi\sigma\pi^{-1} = (\pi(a)\ \pi(b)\ \pi(c)) = (u\ v\ w) = \tau.$$

Since  $H$  is a normal subgroup of  $A_n$ , it follows that  $\tau \in H$  whenever  $\pi \in A_n$ .

If  $\pi$  is odd, then we replace  $\pi$  with  $\pi\delta$ , where  $\delta = (d\ f) \in S_n$  for some  $d, f \in I_n \setminus \{a, b, c\}$  with  $d \neq f$ , and we can always do this because of our assumption  $n \geq 5$ . Since the 2-cycle  $\delta = \delta^{-1}$  is disjoint from  $\sigma$ , they commute, and so  $(\pi\delta)\sigma(\pi\delta)^{-1} = \pi\sigma\pi^{-1} = \tau$ , as required. This completes the proof.  $\square$

**Corollary 2.13.11.** Fix an integer  $n \geq 5$ , and let  $H$  be a normal subgroup of  $A_n$ . If  $H$  contains a product of two disjoint transpositions, then  $H = A_n$ .

*Proof.* Let  $(a\ b)$  and  $(c\ d)$  be two disjoint transpositions in  $S_n$  such that  $(a\ b) \circ (c\ d) \in H$ . To show that  $H = A_n$ , in view of Lemma 2.13.10, it suffices to show that  $H$  contains a 3-cycle. Since  $n \geq 5$ , we can choose an element  $f \in I_n \setminus \{a, b, c, d\}$ . Then the 3-cycle  $\pi := (c\ d\ f) \in A_n$ . Since  $H$  is normal in  $A_n$ , we have  $\pi \circ (a\ b) \circ (c\ d) \circ \pi^{-1} \in H$ . But

$$\begin{aligned} \pi \circ (a\ b) \circ (c\ d) \circ \pi^{-1} &= (c\ d\ f) \circ (a\ b) \circ (c\ d) \circ (c\ f\ d) \\ &= (a\ b) \circ (d\ f). \end{aligned}$$

Since  $H$  is a group containing  $(a\ b) \circ (c\ d)$  and  $(a\ b) \circ (d\ f)$ , we have

$$\pi = (c\ d\ f) = (a\ b) \circ (c\ d) \circ (a\ b) \circ (d\ f) \in H,$$

as required. This completes the proof.  $\square$

**Theorem 2.13.12.** *The alternating group  $A_n$  is simple, for all  $n \geq 5$ .*

*Proof.* Let  $H$  be a non-trivial normal subgroup of  $A_n$ . To show  $A_n$  is simple, thanks to Lemma 2.13.10, it suffices to show that  $H$  contains a 3-cycle.

Let  $\sigma \in H \setminus \{e\}$  be a permutation that moves the smallest number of elements, say  $r$ , of  $I_n := \{1, \dots, n\}$ . If  $r = 2$ , then  $\sigma$  must be a transposition, which is not possible since then  $\sigma$  would be odd while  $H \subseteq A_n$ . Therefore,  $r \geq 3$ . If we can show that  $r = 3$ , then  $\sigma$  must be a 3-cycle and we are done.

Suppose on the contrary that  $r > 3$ . Write  $\sigma$  as a product of finite number of disjoint cycles, say  $\sigma = \sigma_1 \circ \dots \circ \sigma_k$ , where  $\sigma_j$  is a cycle in  $S_n$ , for all  $j \in \{1, \dots, k\}$ .

**Step 1:** Suppose that  $\sigma_j$  is a transposition, for all  $j \in \{1, \dots, k\}$ . Then  $k \geq 2$ , for otherwise  $\sigma = \sigma_1$  would be odd, a contradiction. Let  $\sigma_1 = (a\ b)$  and  $\sigma_2 = (c\ d)$  in  $S_n$ . Since  $\sigma_1$  and  $\sigma_2$  are disjoint cycles and  $n \geq 5$ , there exists an element  $f \in I_n \setminus \{a, b, c, d\}$ . Let  $\tau := (c\ d\ f) \in S_n$ . Since  $\tau$  is even,  $\tau \in A_n$ . Since  $\sigma \in H$  and  $H$  is normal in  $A_n$ , we have  $\tau\sigma\tau^{-1} \in H$ . Since  $H$  is a group,

$$\sigma' := [\sigma^{-1}, \tau] = \sigma^{-1}\tau\sigma\tau^{-1} \in H.$$

Since  $\sigma$  permutes  $a$  and  $b$ , we see that  $\sigma'(a) = a$  and  $\sigma'(b) = b$ . If  $u \in I_n \setminus \{a, b, c, d, f\}$  is such that  $\sigma(u) = u$ , then  $\sigma'(u) = (\sigma^{-1}\tau\sigma\tau^{-1})(u) = u$ . Since  $\sigma'(f) = c$ , we have  $\sigma' \neq e$ . Therefore,  $\sigma' \in H \setminus \{e\}$  moves fewer elements of  $I_n$  than  $\sigma$ , which is a contradiction. Therefore, at least one  $\sigma_i$  must be a cycle of length  $\geq 3$ . Since disjoint cycles commutes, we may assume that  $\sigma_1 = (a\ b\ c\ \dots)$  is a cycle of length  $\geq 3$ .

**Step 2:** If  $r = 4$ , then either  $\sigma$  is a product of two disjoint transpositions or is a 4-cycle. The first possibility is ruled out by step 1 and the second possibility is ruled out since a 4-cycle is odd and  $\sigma \in H \subseteq A_n$ . Therefore,  $r \geq 5$ .

**Step 3:** Since  $n \geq 5$ , we can choose  $d, f \in I_n \setminus \{a, b, c\}$  with  $d \neq f$ . Let  $\tau = (c\ d\ f) \in A_n$ . As before,  $H$  being a normal subgroup of  $A_n$  containing  $\sigma$ , we have  $\sigma' := \sigma^{-1}\tau\sigma\tau^{-1} \in H$ . Since  $\sigma'(b) \neq b$ , we have  $\sigma' \neq e$ . Given any  $u \in I_n \setminus \{a, b, c, d, f\}$ , if  $\sigma(u) = u$ , then  $\sigma'(u) = (\sigma^{-1}\tau\sigma\tau^{-1})(u) = u$ . Moreover  $\sigma(a) \neq a$  while  $\sigma'(a) = a$ . Therefore,  $\sigma' \in H \setminus \{e\}$  moves fewer elements of  $I_n$  than  $\sigma$ , which is a contradiction. Therefore, we must have  $r = 3$ , and hence  $\sigma$  must be a 3-cycle. Hence the result follows.  $\square$

### 2.13.2 \*Simplicity of $\text{PSL}_n(F)$ , for $n \geq 3$

**Definition 2.13.13.** A *field* is a triple  $(F, +, \cdot)$  consisting of a non-empty set  $F$  together with two binary operations  $+: F \times F \rightarrow F$  and  $\cdot: F \times F \rightarrow F$ , called the *addition* and *multiplication* of scalars, respectively, satisfying the following properties:

- (F1) The pair  $(F, +)$  is an abelian group.
- (F2) The pair  $(F^*, \cdot)$  is an abelian group, where  $F^* := F \setminus \{0\}$ .
- (F3) The multiplication operation  $\cdot$  distribute from the left and right over the addition operation, i.e.,

$$a(b + c) = ab + ac, \text{ and } (a + b)c = ac + bc, \forall a, b, c \in F.$$



**Example 2.13.14.** (i) The triples  $(\mathbb{Q}, +, \cdot)$ ,  $(\mathbb{R}, +, \cdot)$  and  $(\mathbb{C}, +, \cdot)$  are fields. The triple  $(\mathbb{Z}, +, \cdot)$  is not a field because  $2 \neq 0$  in  $\mathbb{Z}$ , and it has no multiplicative inverse in  $\mathbb{Z}$ .

(ii)  $(\mathbb{Z}_p, +, \cdot)$  is a field if and only if  $p > 0$  is a prime number.

Fix an integer  $n \geq 2$  and a field  $F$ . For  $i, j \in \{1, \dots, n\}$  with  $i \neq j$  and a non-zero scalar  $c \in F^*$ , let  $A_{ij}(c)$  be the  $n \times n$  matrix whose  $(i, j)$ -th entry is  $c$ , and all other entries are  $0 \in F$ . Then the matrices of the form

$$E_{ij}(c) := I_n + A_{ij}(c)$$

are called the  $n \times n$  elementary matrices over  $F$ .

**Exercise 2.13.15.** Show that  $E_{ij}(c)$  is invertible with  $E_{ij}(c)^{-1} = E_{ij}(-c)$ .

**Exercise 2.13.16.** Given  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ , show that the map

$$F^* \rightarrow \text{GL}_n(F), \quad c \mapsto E_{ij}(c),$$

is a group homomorphism.

**Lemma 2.13.17.** (i) The group  $\text{SL}_n(F)$  is generated by the elementary matrices.

(ii) If  $A \in \text{GL}_n(F)$ , then  $A$  can be written as  $A = SD$ , where  $S \in \text{SL}_n(F)$  and

$$D = \begin{pmatrix} I_{n-1} & 0 \\ 0 & d \end{pmatrix}$$

is the  $n \times n$  diagonal matrix over  $F$  whose  $(n, n)$ -th entry is  $d := \det(A)$  and all other diagonal entries are 1.

*Proof.* Use elementary row operations. □

**Corollary 2.13.18.** Fix an integer  $n \geq 3$ . Then  $\text{SL}_n(F)$  is equal to its own commutator subgroup; i.e.,  $[\text{SL}_n(F), \text{SL}_n(F)] = \text{SL}_n(F)$ .

*Proof.* Since  $\text{SL}_n(F)$  is generated by the elementary matrices, it suffices to show that  $E_{ij}(c)$  is a commutator, for all  $i \neq j$  and  $c \in F^*$ . Since  $n \geq 3$ , we can choose an element  $k \in \{1, \dots, n\} \setminus \{i, j\}$ . Then by a direct computation (do it!) we have

$$E_{ij}(c) = E_{ik}(c)E_{kj}(1)E_{ik}(-c)E_{kj}(-1).$$

This completes the proof. □

**Exercise 2.13.19.** Fix an integer  $n \geq 3$ . Given any abelian group  $G$ , show that any group homomorphism from  $\text{SL}_n(F)$  into  $G$  is trivial.

**Exercise 2.13.20.** (i) Show that  $Z(\text{GL}_n(F)) = \{\lambda I_n : \lambda \in F^*\} \cong F^*$ .

(ii) Show that  $Z(\text{SL}_n(F)) = \{\lambda I_n : \lambda \in F^*, \lambda^n = 1\} \cong \mu_n(F)$ , where  $\mu_n(F) = \{\lambda \in F : \lambda^n = 1\}$  is the group of all  $n$ -th roots of unity in  $F$ .

**Definition 2.13.21.** The quotient groups

$$\begin{aligned} \text{PGL}_n(F) &:= \text{GL}_n(F)/Z(\text{GL}_n(F)) \\ \text{and } \text{PSL}_n(F) &:= \text{SL}_n(F)/Z(\text{SL}_n(F)) \end{aligned}$$

are called the *projective general linear group over  $F$*  and the *projective special linear group over  $F$* , respectively.

**Exercise 2.13.22.** Fix an integer  $n \geq 2$  and a field  $F$ .



- (i) Show that  $(F^*)^n := \{a^n \in F^* : a \in F^*\}$  is a subgroup of  $F^*$  and the map  $p_n : F^* \rightarrow (F^*)^n$ , that sends every element  $a \in F^*$  to its  $n$ -th power  $a^n \in (F^*)^n$ , is a group homomorphism.
- (ii) Show that the map  $\det : \mathrm{GL}_n(F) \rightarrow F^*$ , defined by sending  $A \in \mathrm{GL}_n(F)$  to its determinant  $\det(A) \in F^*$ , is a surjective group homomorphism.
- (iii) Show that  $Z(\mathrm{GL}_n(F))$  is sitting inside the kernel of the composite group homomorphism

$$\varphi : \mathrm{GL}_n(F) \xrightarrow{\det} F^* \xrightarrow{\pi_n} F^*/(F^*)^n,$$

and hence give rise to a surjective group homomorphism

$$\tilde{\varphi} : \mathrm{PGL}_n(F) \rightarrow F^*/(F^*)^n.$$

- (iv) Let  $\pi : \mathrm{GL}_n(F) \rightarrow \mathrm{PGL}_n(F)$  be the natural quotient group homomorphism. Show that  $\mathrm{Ker}(\pi \circ \iota) = Z(\mathrm{SL}_n(F))$ , where  $\iota : \mathrm{SL}_n(F) \hookrightarrow \mathrm{GL}_n(F)$  is the inclusion map (group homomorphism).
- (v) Conclude that the induced group homomorphism

$$\widetilde{\pi \circ \iota} : \mathrm{PSL}_n(F) \longrightarrow \mathrm{PGL}_n(F)$$

is injective, and it identifies  $\mathrm{PSL}_n(F)$  as a normal subgroup of  $\mathrm{PGL}_n(F)$ . (Hint: Use Exercise 2.9.27).

- (vi) Show that the associated quotient group  $\mathrm{PGL}_n(F)/\mathrm{PSL}_n(F)$  is naturally isomorphic to  $F^*/(F^*)^n$ . The following diagram of group homomorphisms commutes.

$$\begin{array}{ccccc} Z(\mathrm{SL}_n(F)) & \hookrightarrow & Z(\mathrm{GL}_n(F)) \cong F^* & \xrightarrow{z \mapsto z^n} & (F^*)^n \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{SL}_n(F) & \hookrightarrow & \mathrm{GL}_n(F) & \xrightarrow{\det} & F^* \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{PSL}_n(F) & \hookrightarrow & \mathrm{PGL}_n(F) & \twoheadrightarrow & F^*/(F^*)^n \end{array}$$

- (vii) Conclude that  $\mathrm{PSL}_n(F) \cong \mathrm{PGL}_n(F)$  if  $F^*$  contains  $n$ -th roots of all of its elements.
- (viii) Let  $F$  be a finite field having  $q := p^n$  elements, where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Find the orders of  $\mathrm{PGL}_n(F)$  and  $\mathrm{PSL}_n(F)$ .

Note that the group  $\mathrm{GL}_n(F)$  acts on the  $n$ -dimensional  $F$ -vector space  $V := F^n$  by

$$(A, v) \mapsto Av, \quad \forall (A, v) \in \mathrm{GL}_n(F) \times V.$$

Given a non-zero  $F$ -linear functional  $\lambda : V \rightarrow F$ , the subset

$$H_\lambda := \mathrm{Ker}(\lambda) = \{v \in V : \lambda(v) = 0\}$$

is a  $F$ -linear subspace of  $V$  of dimension  $n - 1$ , called the *hyperplane section associated to  $\lambda$* . Note that every  $F$ -linear subspace  $H$  of  $V$  of dimension  $n - 1$  is a hyperplane section associated to some  $F$ -linear functional on  $V$ . Indeed, given  $H$ , the associated quotient space  $V/H$  has dimension 1, and hence is  $F$ -linearly isomorphic to  $F$ . Then the quotient map

$$q : V \rightarrow V/H \cong F$$

is the required  $F$ -linear functional on  $V$  whose kernel is precisely  $H$ , as required.

**Definition 2.13.23** (Transvection). An element  $T \in \text{GL}_n(F)$  is said to be a *transvection* if there exists a hyperplane section  $H \subset V$  of  $V$  such that

- (i)  $T$  fixes each vectors of  $H$ , i.e.,  $Tv = v$ ,  $\forall v \in H$ , and
- (ii) for each  $v \in V$ , there exists a vector  $h(v) \in H$  such that  $Tv = v + h(v)$ .

To emphasis dependency on  $H$ , we may call  $T$  a *transvection with respect to the hyperplane  $H$* .

**Exercise 2.13.24.** Let  $H_\lambda = \text{Ker}(\lambda)$  be a hyperplane section in  $V$  associated to a linear functional  $\lambda : V \rightarrow F$ . Given a vector  $u \in H_\lambda$ , define a map  $T_u : V \rightarrow V$  by

$$T_u(v) := v + \lambda(v)u, \forall v \in V.$$

- (i) Show that  $T_u$  is an invertible  $F$ -linear map, and hence with respect to the standard ordered basis for  $V = F^n$  it defines an element of  $\text{GL}_n(F)$ ; for notational simplicity, we denote it by the same symbol  $T_u$ .
- (ii) If  $T \in \text{GL}_n(F) \setminus \{I_n\}$  is a transvection, show that there exists a  $F$ -linear functional  $\lambda : V \rightarrow F$  and a vector  $u \in H_\lambda = \text{Ker}(\lambda)$  such that  $T(v) = v + \lambda(v)u$ ,  $\forall v \in V$ . (Hint: Since the map  $x \mapsto h(x) := T(x) - x$  is an  $F$ -linear with kernel  $\text{Ker}(h) = H$  (because  $T \neq I_n$ ), the image of  $h$  is an 1-dimensional  $F$ -linear subspace of  $V$ , and hence is generated by a non-zero vector, say  $u \in H$ , so that for each  $x \in V$ , we have  $h(x) = T(x) - x = \lambda(x)u$ , for some  $\lambda(x) \in F$ , from which one can easily verify that  $\lambda \in V^*$ , as required.)
- (iii) Given  $u_1, u_2 \in H_\lambda$ , show that  $T_{u_1} \circ T_{u_2} = T_{u_1+u_2} = T_{u_2} \circ T_{u_1}$ .
- (iv) If  $T \in \text{GL}_n(F)$  is a transvection with respect to a hyperplane  $H$  in  $V$ , show that  $ATA^{-1}$  is a transvection with respect to the hyperplane  $A(H)$  in  $V$ , for all  $A \in \text{GL}_n(F)$ .
- (v) If  $S$  and  $T$  are transvections in  $\text{GL}_n(F) \setminus \{I_n\}$  with respect to the same hyperplane  $H$  in  $V$ , so is their product  $ST$ . Show by an example that product of two transvections in  $\text{GL}_n(F)$  need not be a transvection, in general.
- (vi) Show that the elementary matrices  $E_{ij}(c) \in \text{GL}_n(F)$  are transvections, for all  $c \in F$ . (Hint: Let  $e_1, \dots, e_n$  be the standard ordered basis for  $V = F^n$  over  $F$ . Then for given  $i, j \in \{1, \dots, n\}$  with  $i \neq j$ , we have  $E_{ij}(c)(e_k) = e_k$ , for all  $k \neq j$ , and  $E_{ij}(c)(e_j) = ce_j$ .)
- (vii) If  $T \in \text{GL}_n(F)$  is a transvection, show that  $\det(T) = 1$ . (Hint: Let  $H$  be the hyperplane section in  $V = F^n$  that is point-wise fixed by  $T$ . Choose an ordered basis for  $H$  and then extend it to an ordered basis for  $V$  by choosing one vector from  $V \setminus H$ .)

**Lemma 2.13.25.** Fix an integer  $n \geq 3$ .

- (i) The set of all transvections in  $\text{GL}_n(F)$  is a subset of  $\text{SL}_n(F)$ .
- (ii) The set of all transvections in  $\text{SL}_n(F) \setminus \{I_n\}$  forms a single conjugacy class in  $\text{SL}_n(F)$ .

*Proof.* The first statement follows from Exercise 2.13.24 (vii). To prove the second part, since conjugate of a transvection by an element of  $\text{GL}_n(F)$  is again a transvection (see Exercise 2.13.24 (iv)), it suffices to show that given any two transvections  $T, T' \in \text{SL}_n(F) \setminus \{I_n\}$ , there exists  $A \in \text{SL}_n(F)$  such that  $T' = ATA^{-1}$ .

Let  $H = \text{Ker}(\lambda)$  and  $H' = \text{Ker}(\lambda')$  be the hyperplane sections in  $V = F^n$  that are point-wise fixed by  $T$  and  $T'$ , respectively (c.f. Exercise 2.13.24 (ii)). Then there exists  $u \in H$  and  $u' \in H'$  such that

$$T(v) = v + \lambda(v)u \quad \text{and} \quad T'(v) = v + \lambda'(v)(u'), \forall v \in V.$$

Since  $T \neq I_n \neq T'$ , we have  $u \neq 0$  and  $u' \neq 0$ . Choose some ordered bases  $\{v_1, \dots, v_{n-1}\}$  and  $\{v'_1, \dots, v'_{n-1}\}$  for  $H$  and  $H'$ , respectively, with  $v_1 = u$  and  $v'_1 = u'$ . Since  $\lambda$  and  $\lambda'$  are surjective,

we can get non-zero vectors  $v_n, v'_n \in F^n$  such that  $\lambda(v_n) = 1$  and  $\lambda(v'_n) = 1$ . Then the ordered bases  $\{v_1, \dots, v_n\}$  and  $\{v'_1, \dots, v'_n\}$  for  $V = F^n$  gives an invertible matrix  $A \in \text{GL}_n(F)$  such that

$$Av_i = v'_i, \quad \forall i \in \{1, \dots, n\}.$$

Note that,

$$T(v_i) = \begin{cases} v_i, & \text{if } 1 \leq i \leq n-1, \\ v_n + u, & \text{if } i = n; \end{cases}$$

and similarly,

$$T'(v'_i) = \begin{cases} v'_i, & \text{if } 1 \leq i \leq n-1, \\ v'_n + u', & \text{if } i = n. \end{cases}$$

Then by looking at the images of  $v'_i, \forall i \in \{1, \dots, n\}$ , under the composite map

$$V \xrightarrow{A^{-1}} V \xrightarrow{T} V \xrightarrow{A} V,$$

we see that  $ATA^{-1} = T'$  (verify!). Thus,  $T$  and  $T'$  are conjugate in  $\text{GL}_n(F)$ . There is no reason to expect to have  $\det(A) = 1$ , in general. Since  $n \geq 3$ , we can replace  $A$  with the matrix  $B$ , where

$$B(v_i) = \begin{cases} v'_i, & \text{if } i \neq n-1, \\ \det(A)^{-1}v'_{n-1}, & \text{if } i = n-1; \end{cases}$$

so that  $\det(B) = 1$  and that  $BTB^{-1} = T'$  (verify!). This completes the proof.  $\square$

**Definition 2.13.26.** A subgroup  $G$  of  $\text{GL}_n(F)$  is said to be  $\text{SL}_n(F)$ -invariant if

$$aGa^{-1} \subseteq G, \quad \forall a \in \text{SL}_n(F).$$

**Lemma 2.13.27.** Fix an integer  $n \geq 3$ . Let  $G$  be an  $\text{SL}_n(F)$ -invariant subgroup of  $\text{GL}_n(F)$ . If  $G$  contains a transvection  $T \neq I_n$ , then  $\text{SL}_n(F) \subseteq G$ .

*Proof.* Let  $T \neq I_n$  be a transvection in  $\text{GL}_n(F)$  such that  $T \in G$ . Since  $\text{SL}_n(F)$  is generated by elementary matrices  $\{E_{ij}(c) : c \in F^*, i \neq j\}$  by Lemma 2.13.17, to show  $\text{SL}_n(F) \subseteq G$ , it suffices to show that elementary matrices are in  $G$ . Since all elementary matrices are transvections by Exercise 2.13.24, they are conjugate to  $T$  in  $\text{SL}_n(F)$  by Lemma 2.13.25. Now since  $G$  is  $\text{SL}_n(F)$ -invariant, the result follows.  $\square$

**Exercise 2.13.28.** Let  $V$  be a finite dimensional  $F$ -vector space and let  $T : V \rightarrow V$  be an invertible  $F$ -linear map. If  $W \subseteq V$  is an  $F$ -linear subspace of  $V$  of dimension  $r$ , so is its image  $T(W)$ .

**Lemma 2.13.29.** Fix an integer  $n \geq 3$ . Let  $G$  be a  $\text{SL}_n(F)$ -invariant subgroup of  $\text{GL}_n(F)$ . If  $G$  is not contained in the center of  $\text{GL}_n(F)$ , then  $\text{SL}_n(F) \subseteq G$ .

*Proof.* By Lemma 2.13.27 it suffices to show that  $G \setminus \{I_n\}$  contains a transvection. Since  $G \not\subseteq Z(\text{GL}_n(F)) = \{cI_n : c \in F^*\}$ , there exists  $A \in G$  that moves a straight-line; i.e., there exists  $u \in F^n \setminus \{0\}$  such that  $Au \notin \{cu : c \in F\}$ . Fix a hyperplane  $H = \text{Ker}(\lambda)$  in  $F^n$  containing  $u$  and  $v := Au$ , and consider the transvection

$$T = T_u = I_n + u\lambda \in \text{SL}_n(F).$$

Since  $\lambda : F^n \rightarrow F$  is a non-zero  $F$ -linear functional, there exists  $x \in F^n$  such that  $\lambda(x) = 1$ . Then we have

$$\begin{aligned} (AT_u)(x) - (T_uA)(x) &= A(x + \lambda(x)u) - (Ax + \lambda(Ax)u) \\ &= \lambda(x)Au - \lambda(Ax)u \\ &= Au - \lambda(Ax)u. \end{aligned}$$

Since  $Au \notin \{cu \in V : c \in F\}$  by assumption, it follows that  $AT_u \neq T_uA$ , and hence  $B := AT_uA^{-1}T_u^{-1} \neq I_n$ . Since  $A \in G$  and  $G$  is  $\mathrm{SL}_n(F)$ -invariant subgroup of  $\mathrm{GL}_n(F)$ , it follows that  $B = A(TA^{-1}T^{-1}) \in G$ .

By a direct computation we have

$$Bx - x = (\lambda(A^{-1}x) - \lambda(x)\lambda(A^{-1}u))v - \lambda(x)u \in \mathrm{Span}_F\{u, v\} \subseteq H, \forall x \in V.$$

Then it follows that  $Bx = (Bx - x) + x \in H$ ,  $\forall x \in H$ , and hence  $B(H) \subseteq H$ . Since  $B = ATA^{-1}T^{-1}$  is invertible and  $H$  is finite dimensional, we have  $B(H) = H$ . Now we have the following two cases.

**Case 1:** Suppose that  $B$  commutes with all transvections with respect to  $H$ . Let  $w \in H$  be given. Since  $BT_w = T_wB$ , for given any  $x \in V$  we have

$$\begin{aligned} B(x + \lambda(x)w) &= B(x) + \lambda(B(x))w \\ \Rightarrow \lambda(x)B(w) &= \lambda(B(x))w = \lambda(x)w, \end{aligned}$$

where the last equality follows because  $B(x) - x \in H$  and  $\lambda : V \rightarrow V/H \cong F$  is the quotient map. Since  $\lambda \neq 0$ , it follows that  $B(w) = w$ . Therefore,  $B$  point-wise fixes every vector of  $H$ . Since  $B(x) - x \in H$ , for all  $x \in V$ , it follows from Definition 2.13.23 that  $B$  is a transvection in  $\mathrm{GL}_n(F)$ , as required.

**Case 2:** Suppose that  $BT \neq TB$ , for some transvection  $T \in \mathrm{GL}_n(F)$  with respect to  $H$ . Now  $T = T_w$ , for some  $w \in H$ . Then  $C := BT_wB^{-1}T_w^{-1} \neq I_n$ . Since  $B \in G$ ,  $T_w \in \mathrm{SL}_n(F)$ , and  $G$  is  $\mathrm{SL}_n(F)$ -invariant subgroup, it follows that  $C = B(T_wB^{-1}T_w^{-1}) \in G$ . Note that  $T_w^{-1} = T_{-w}$  and  $BT_wB^{-1}$  are transvections with respect to the hyperplanes  $H$  and  $B(H)$ , respectively. Since  $B(H) = H$ , it follows from Exercise 2.13.24 (v) that the product  $C = (BT_wB^{-1})T_w^{-1}$  is a transvection with respect to  $H$ . This completes the proof.  $\square$

**Theorem 2.13.30.** *The group  $\mathrm{PSL}_n(F)$  is simple, for all  $n \geq 3$ .*

*Proof.* Let  $H$  be a non-trivial normal subgroup of  $\mathrm{PSL}_n(F) = \mathrm{SL}_n(F)/Z(\mathrm{SL}_n(F))$ . Then  $H = N/Z(\mathrm{SL}_n(F))$ , for some normal subgroup  $N$  of  $\mathrm{SL}_n(F)$  that properly contains  $Z(\mathrm{SL}_n(F))$ . Since  $Z(\mathrm{SL}_n(F)) = Z(\mathrm{GL}_n(F)) \cap \mathrm{SL}_n(F)$  by Exercise 2.13.20, it follows that  $N$  is not contained in the center of  $\mathrm{GL}_n(F)$ . Then  $\mathrm{SL}_n(F) \subseteq N$  by Lemma 2.13.29, and hence  $N = \mathrm{SL}_n(F)$ . Then  $H = N/Z(\mathrm{SL}_n(F)) = \mathrm{PSL}_n(F)$ . This completes the proof.  $\square$

**Remark 2.13.31.** For  $n = 2$ , it turns out that  $\mathrm{PSL}_2(\mathbb{Z}_5) \cong A_5$ , and hence is simple, while  $\mathrm{PSL}_2(\mathbb{Z}_3) \cong A_4$ , and hence is not simple.

## 2.14 Sylow's Theorems

**Theorem 2.14.1** (Sylow's Theorem I). *Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r$  and  $m$  are positive integers, and  $\gcd(m, p) = 1$ . Then for each  $k \in \{1, \dots, r\}$ ,  $G$  has a subgroup of order  $p^k$ .*

*Proof.* Since  $p \mid |G|$ , by Cauchy's theorem  $G$  has a subgroup of order  $p$ , which proves the statement for the case  $k = 1$ . Suppose that  $1 < k \leq r$ , and  $G$  has a subgroup, say  $H$ , of order  $p^{k-1}$ . Since  $H$  is a finite  $p$ -subgroup of the finite group  $G$ , considering the class equation of the set  $\mathcal{L}_H := \{aH : a \in G\}$  associated to the left translation action of  $H$  on  $\mathcal{L}_H$  we see that

$$[G : H] \equiv_p [N_G(H) : H]$$

(see Corollary 2.12.26). Since  $p \mid [G : H]$  and  $[N_G(H) : H] \geq 1$ , we conclude that  $p \mid [N_G(H) : H]$ , and hence  $H$  is a proper normal subgroup of  $N_G(H)$ . Then  $p$  divides the order of the

quotient group  $N_G(H)/H$ . Then by Cauchy's theorem  $N_G(H)/H$  has a subgroup, say  $S$ , of order  $p$ . By Exercise 2.9.26 we have  $S = K/H$ , for some subgroup  $K$  of  $N_G(H)$  containing  $H$ . Therefore,  $K$  is a subgroup of  $G$  of order  $|K| = |K/H| \cdot |H| = p \cdot p^{k-1} = p^k$ , as required. This completes the proof by induction on  $k$ .  $\square$

**Exercise 2.14.2.** Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Let  $H$  be a subgroup of  $G$  of order  $p^k$ , for some  $k \in \{1, \dots, r-1\}$ . Show that there exists a subgroup  $K$  of  $G$  containing  $H$  such that  $|K| = p^{k+1}$  and  $H$  is normal in  $K$ .

*Solution:* Follows from the proof of Sylow's first theorem (Theorem 2.14.1) by noting that if  $K$  is a subgroup of  $N_G(H)$  containing  $H$ , then  $H$  is a normal subgroup of  $K$ .  $\square$

As an immediate corollary, we have the following generalization of Cauchy's theorem.

**Corollary 2.14.3.** Let  $G$  be a finite group. If  $p^n \mid |G|$ , for some prime number  $p > 0$  and an integer  $n \geq 0$ , then  $G$  has a subgroup of order  $p^n$ .

**Definition 2.14.4 (Sylow  $p$ -subgroup).** Let  $G$  be a finite group and  $p > 0$  a prime number. A subgroup  $P$  of  $G$  is said to be a Sylow  $p$ -subgroup of  $G$  if  $P$  is a maximal  $p$ -subgroup of  $G$ ; i.e.,  $P$  is a  $p$ -subgroup of  $G$  that is not properly contained in any other  $p$ -subgroup of  $G$ .

**Example 2.14.5.** The symmetric group  $S_3$  has three Sylow 2-subgroups, namely

$$H_1 := \{e, (1\ 2)\}, H_2 := \{e, (1\ 3)\}, \text{ and } H_3 := \{e, (2\ 3)\},$$

and one Sylow 3-subgroup, namely  $K := \{e, (1\ 2\ 3), (1\ 3\ 2)\}$ . Therefore, a Sylow  $p$ -subgroup, for certain prime  $p$ , need not be unique. In this case, the unique Sylow 3-subgroup  $K$  of  $S_3$  is normal; this is not a coincidence. We shall see later that a unique Sylow  $p$ -subgroup of  $G$  must be normal.

**Proposition 2.14.6.** Let  $G$  be a finite group. Then for each prime number  $p > 0$ ,  $G$  has a Sylow  $p$ -subgroup.

*Proof.* If  $p \nmid |G|$ , then  $\{e\}$  is the Sylow  $p$ -subgroup of  $G$ . If  $p \mid |G|$ , then there exists  $r \in \mathbb{N}$  such that  $|G| = p^r m$ , for some integer  $m \geq 1$  with  $\gcd(p, m) = 1$  by the fundamental theorem of arithmetic. Then  $G$  has a subgroup, say  $P$ , of order  $p^r$  by Sylow's first theorem (Theorem 2.14.1). If  $Q$  is any  $p$ -subgroup of  $G$  containing  $P$ , then  $|Q| = p^k$ , for some  $k \in \mathbb{N}$  and that  $p^r = |P|$  divides  $p^k = |Q|$  by Lagrange's theorem. Then  $r \leq k$ . Since  $p^k = |Q|$  divides  $|G| = p^r m$  by Lagrange's theorem and since  $\gcd(p, m) = 1$ , we must have  $k \leq r$ , and hence  $k = r$ . Then  $|P| = p^r = |Q|$  gives  $P = Q$ . Therefore,  $P$  is a required Sylow  $p$ -subgroup of  $G$ .  $\square$

**Exercise 2.14.7.** Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Let  $P$  be a subgroup of  $G$ . Show that  $P$  is a Sylow  $p$ -subgroup of  $G$  if and only if  $|P| = p^r$ .

**Exercise 2.14.8.** Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Let  $P$  be a subgroup of  $G$ . Prove the following statements.

- (i) If  $P$  is a  $p$ -subgroup of  $G$ , so is  $aPa^{-1}$ , for all  $a \in G$ .
- (ii) If  $P$  is a Sylow  $p$ -subgroup of  $G$ , so is  $aPa^{-1}$ , for all  $a \in G$ .
- (iii) If  $P$  is the only Sylow  $p$ -subgroup of  $G$ , then  $P$  is normal in  $G$ .

*Solution:* (i) Follows from the fact that  $|aPa^{-1}| = |P|$ , for all  $a \in G$ .

(ii) Follows from (i) and Exercise 2.14.7.

(iii) Since  $P$  is the only Sylow  $p$ -subgroup of  $G$ , using part (ii) we have  $P = aPa^{-1}$ , for all  $a \in G$ . Therefore,  $P$  is normal in  $G$ .  $\square$

**Lemma 2.14.9.** *Let  $H$  be a normal subgroup of a group  $G$ . If both  $H$  and  $G/H$  are  $p$ -groups, then  $G$  is a  $p$ -group.*

*Proof.* Let  $a \in G$  be arbitrary. Since  $G/H$  is a  $p$ -group,  $\text{ord}(aH) = p^r$ , for some integer  $r \geq 0$ . If  $r = 0$ , then  $a \in H$ , and then the result follows since  $H$  is a  $p$ -group. Assume that  $r > 0$ . Then  $a^{p^r}H = (aH)^{p^r} = H$  gives  $a^{p^r} \in H$ . Since  $H$  is a  $p$ -group,  $\text{ord}(a^{p^r}) = p^n$ , for some integer  $n \geq 0$ . Then  $a^{p^{r+n}} = e$ , and hence  $\text{ord}(a) \mid p^{r+n}$ . Hence the result follows. Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ .  $\square$

**Exercise 2.14.10.** Let  $G$  be a finite group. Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Let  $a \in G$  be such that  $\text{ord}(a) = p^r$ , for some integer  $r \geq 0$ . If  $aPa^{-1} = P$ , show that  $a \in P$ .

*Solution:* Since  $aPa^{-1} = P$ , we have  $a \in N_G(P)$ . Note that  $P \subseteq N_G(P)$ . Let  $b \in N_G(P) \setminus P$  be arbitrary. If  $\text{ord}(b) = p^k$ , for some integer  $k \geq 0$ , then considering the quotient homomorphism  $\pi : N_G(P) \rightarrow N_G(P)/P$  we see that  $\text{ord}(bP)$  divides  $\text{ord}(b) = p^k$ . Then the cyclic subgroup  $\langle bP \rangle$  of  $N_G(P)/P$  has order  $p^m$ , for some integer  $m \geq 0$ . Now  $\langle bP \rangle = K/P$ , for some subgroup  $K$  of  $N_G(P)$  containing  $P$  by Exercise 2.9.26. Since  $b \notin P$  by assumption,  $P \subsetneq K$ . Since both  $P$  and  $K/P$  are  $p$ -groups, so is  $K$  by Lemma 2.14.9. But this contradicts the maximality of  $P$  as it is a Sylow  $p$ -subgroup of  $G$ . Therefore,  $\text{ord}(b)$  cannot be a power of  $p$ , and hence  $a \in P$ .  $\square$

**Theorem 2.14.11** (Sylow's Theorem II). *Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Then any two Sylow  $p$ -subgroups of  $G$  are conjugate, and hence are isomorphic.*

*Proof.* Let  $H$  and  $K$  be two Sylow  $p$ -subgroups of  $G$ . Let  $\mathcal{L}_H := \{aH : a \in G\}$  be the set of all left cosets of  $H$  in  $G$ . Since  $G$  is finite, so is the set  $\mathcal{L}_H$ . Define a map

$$\sigma : K \times \mathcal{L}_H \rightarrow \mathcal{L}_H$$

by

$$\sigma(b, aH) = (ba)H, \forall (b, aH) \in K \times \mathcal{L}_H.$$

If  $aH = a'H$ , for some  $a, a' \in G$ , then  $(ba)^{-1}(ba') = a^{-1}b^{-1}ba' = a^{-1}a' \in H$ , and hence  $(ba)H = (ba')H$ , for all  $b \in K$ . Therefore, the map  $\sigma$  is well-defined. It is easy to check that  $\sigma$  is a left action of  $K$  on  $\mathcal{L}_H$ . The subset of all elements of  $\mathcal{L}_H$  with singleton  $K$ -orbits is given by

$$\begin{aligned} \mathcal{L}_{H,0} &:= \{aH \in \mathcal{L}_H : baH = aH, \forall b \in K\} \\ &= \{aH \in \mathcal{L}_H : aba^{-1} \in H, \forall b \in K\} \\ &= \{aH \in \mathcal{L}_H : aKa^{-1} \subseteq H\} \\ &= \{aH \in \mathcal{L}_H : aKa^{-1} = H\}, \end{aligned}$$

since both  $H$  and  $K$  are finite Sylow  $p$ -subgroups of  $G$ , they have the same cardinality. Since  $K$  is a finite  $p$ -group, considering the class equation for  $\mathcal{L}_H$  associated to the  $K$ -action  $\sigma$  on it, we have

$$[G : H] \equiv_p |\mathcal{L}_{H,0}|.$$

Since  $H$  is a Sylow  $p$ -subgroup of  $G$ , we see that  $p \nmid [G : H]$ . Therefore,  $|\mathcal{L}_{H,0}| \geq 1$ , and hence there exists  $a \in G$  such that  $aKa^{-1} = H$ . Since for any  $a \in G$ , the conjugation by  $a$  map

$$c_a : G \rightarrow G, c_a(g) = a^{-1}ga, \forall g \in G,$$

is an automorphism of  $G$ , we conclude that any two Sylow  $p$ -subgroup of  $G$  are isomorphic. This completes the proof.  $\square$

**Theorem 2.14.12** (Sylow's Theorem III). *Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Let  $n_p$  be the number of Sylow  $p$ -subgroups of  $G$ . Then  $n_p = 1 + kp$ , for some  $k \in \mathbb{N} \cup \{0\}$ , and  $n_p \mid m$ .*

*Proof.* Let  $X = \text{Syl}_p(G)$  be the set of all Sylow  $p$ -subgroups of  $G$ . Note that  $X \neq \emptyset$  by Sylow's first theorem. Fix a Sylow  $p$ -subgroup  $P \in X$ . Note that  $P$  acts on  $X$  by conjugation:

$$P \times X \rightarrow X, (a, Q) \mapsto aQa^{-1}.$$

Let  $X_0 := \{Q \in X : aQa^{-1} = Q, \forall a \in P\}$ . Since  $aPa^{-1} = P, \forall a \in P$ , we have  $P \in X_0$ . So  $X_0 \neq \emptyset$ . Let  $Q \in X_0$  be arbitrary. Then  $aQa^{-1} = Q$ , for all  $a \in P$ , and so  $P \subseteq N_G(Q)$ . Since both  $P$  and  $Q$  are Sylow  $p$ -subgroups of  $G$  contained in  $N_G(Q)$ , we conclude that  $P$  and  $Q$  are also Sylow  $p$ -subgroups of  $N_G(Q)$ . Then by Sylow's second theorem (Theorem 2.14.11)  $P$  and  $Q$  are conjugate in  $N_G(Q)$ . So there exists  $a \in N_G(Q)$  such that  $aQa^{-1} = P$ . But  $aQa^{-1} = Q$ , since  $a \in N_G(Q)$ . Therefore,  $P = Q$ , and hence  $X_0 = \{P\}$  is singleton. Then by Lemma 2.12.25 we have  $n_p = |X| \equiv_p 1$ , and hence  $n_p = 1 + kp$ , for some  $k \in \mathbb{N} \cup \{0\}$ .

For the second part, we consider the conjugation action of  $G$  on  $X$ . Since any two Sylow  $p$ -subgroups of  $G$  are conjugate by Sylow's second theorem (Theorem 2.14.11), we see that  $X$  has only one  $G$ -orbit, i.e.,

$$X = \text{Orb}_G(P) = \{aPa^{-1} : a \in G\},$$

for any Sylow  $p$ -subgroup  $P$  of  $G$ . Since the stabilizer of  $P \in X$  is

$$G_P = \{a \in G : aPa^{-1} = P\} = N_G(P),$$

the normalizer of  $P$  in  $G$ , we see that

$$\begin{aligned} n_p = |X| &= [G : N_G(P)] \\ &= \frac{[G : P]}{[N_G(P) : P]}, \text{ since } P \subseteq N_G(P) \subseteq G. \\ &= \frac{m}{[N_G(P) : P]}, \text{ since } |P| = p^r \text{ by Exercise 2.14.7,} \end{aligned}$$

and hence  $n_p \mid m$ . This completes the proof.  $\square$

**Definition 2.14.13.** Let  $G$  be a group. A subgroup  $H$  of  $G$  is said to be a *characteristic subgroup* of  $G$  if  $f(H) \subseteq H$ , for all  $f \in \text{Aut}(G)$ .

For example, for any group  $G$ , its trivial subgroup  $\{e\}$  and  $G$  itself are characteristic subgroups of  $G$ .

**Proposition 2.14.14.** Let  $G$  be a group and  $H$  a subgroup of  $G$ .

- (i) If  $H$  is a characteristic subgroup of  $G$ , then  $H$  is normal in  $G$ .
- (ii) If  $H$  is the unique subgroup of a given finite order, then  $H$  is a characteristic subgroup of  $G$ .
- (iii) If  $K$  is a characteristic subgroup of  $H$  and  $H$  is normal in  $G$ , then  $K$  is normal in  $G$ .

*Proof.* (i) Take inner automorphisms  $b \mapsto a^{-1}ba$ , for all  $a \in G$ .

(ii) Follows from the fact that automorphisms preserve the order of subgroups.

(iii) Given  $a \in G$  let

$$\varphi_a : G \rightarrow G$$

be the map defined by

$$\varphi_a(b) = a^{-1}ba, \forall b \in G.$$

Then  $\varphi_a \in \text{Aut}(G)$ . Since  $H$  is a normal subgroup of  $G$ , we have  $\varphi_a(H) = a^{-1}Ha = H$ , and so the restriction map  $\varphi_a|_H : H \rightarrow H$  is an automorphism of  $H$ . Since  $K$  is a characteristic subgroup of  $H$ , we have  $a^{-1}Ka = \varphi_a|_H(K) \subseteq K$ . Therefore,  $K$  is normal in  $G$ .  $\square$



**Corollary 2.14.15.** *Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Then the following are equivalent.*

- (i)  $P$  is the unique Sylow  $p$ -subgroup of  $G$ , i.e.,  $n_p = 1$ .
- (ii)  $P$  is normal in  $G$ .
- (iii)  $P$  is characteristic in  $G$ .
- (iv) All subgroups generated by elements of  $p$ -power order are  $p$ -groups, i.e., if  $S$  is any subset of  $G$  such that for any  $x \in S$  we have  $\text{ord}(x) = p^n$ , for some integer  $n \geq 0$ , then  $\langle S \rangle$  is a  $p$ -group.

*Proof.* (i)  $\Rightarrow$  (ii): Since for any  $a \in G$ ,  $a^{-1}Pa$  is also a Sylow  $p$ -subgroup of  $G$ , we have  $a^{-1}Pa = P$  by (i), and hence  $P$  is normal in  $G$ .

(ii)  $\Rightarrow$  (iii): Let  $f \in \text{Aut}(G)$ . Since  $P$  is a Sylow  $p$ -subgroup of  $G$ , so is its image  $f(P)$  in  $G$  by Exercise 2.14.7. Since any two Sylow  $p$ -subgroups of  $G$  are conjugate by Sylow's second theorem (Theorem 2.14.11), we have  $f(P) = a^{-1}Pa$ , for some  $a \in G$ . Since  $P$  is normal by assumption (ii), we have  $f(P) = a^{-1}Pa = P$ . This proves (iii).

(iii)  $\Rightarrow$  (iv): Let  $S$  be a subset of  $G$  such that for any  $x \in S$  we have  $\text{ord}(x) = p^n$ , for some integer  $n \geq 0$ , and let  $H = \langle S \rangle$  be the subgroup of  $G$  generated by  $S$ . We show that  $H$  is a  $p$ -group. Let  $x \in S$  be arbitrary. Since  $\text{ord}(x) = p^n$ , for some integer  $n \geq 0$ , the cyclic group  $\langle x \rangle$  is a  $p$ -group by Proposition 2.12.20, and hence it is contained in a maximal  $p$ -subgroup, say  $Q$ , of  $G$  by Lemma 2.12.22. Then  $Q$  is a Sylow  $p$ -subgroup of  $G$ , and hence by Sylow's second theorem  $Q = a^{-1}Pa$ , for some  $a \in G$ . Since  $P$  is a characteristic subgroup of  $G$ , we have  $Q = a^{-1}Pa = \varphi_a(P) = P$ , where  $\varphi_a$  is the inner automorphism of  $G$  defined by  $\varphi_a(b) = a^{-1}ba$ ,  $\forall b \in G$ . Therefore,  $x \in P$ ,  $\forall x \in S$ , and hence  $H := \langle S \rangle \subseteq P$ . Therefore,  $H$  is a  $p$ -group by Lemma 2.12.21.

(iv)  $\Rightarrow$  (i): Let  $P$  and  $Q$  be Sylow  $p$ -subgroups of  $G$ . Since both  $P$  and  $Q$  are  $p$ -subgroups of  $G$ , all of their elements have order power of  $p$ . Then  $\langle P \cup Q \rangle$  is a  $p$ -group by assumption (iv). Since Sylow  $p$ -subgroups of  $G$  are maximal  $p$ -subgroups of  $G$ , we have  $P = \langle P \cup Q \rangle = Q$ . This proves (i).  $\square$

**Corollary 2.14.16.** *If  $G$  is a finite abelian group, then for every prime number  $p > 0$ ,  $G$  has a unique Sylow  $p$ -subgroup, known as the  $p$ -primary component of  $G$ .*

*Proof.* Let  $G$  be a finite abelian group of order  $n$ . If  $p \nmid n$ , then the trivial subgroup  $\{e\}$  is the unique Sylow  $p$ -subgroup of  $G$ . If  $p \mid n$ , then the result follows from Corollary 2.14.15.  $\square$

**Exercise 2.14.17.** Show that  $S_4$  has no normal subgroups of order 8 and 3.

*Solution:* If possible suppose that  $S_4$  has a normal subgroup, say  $H$ , of order 8. Then the quotient group  $S_4/H$  is isomorphic to  $\mathbb{Z}_3$ . Since the abelianization of  $S_n$  is isomorphic to  $\mathbb{Z}_2$  by Exercise 2.9.21, we get a surjective group homomorphism  $f : \mathbb{Z}_2 \rightarrow \mathbb{Z}_3$  by Theorem 2.9.19, which is a contradiction. Therefore,  $S_4$  cannot have a normal subgroup of order 8.

If possible suppose that  $S_4$  has a normal subgroup, say  $H$ , of order 3. Since  $H$  has only two elements of order 3 and  $S_4$  has more than 3 elements of order 3, there exists  $x \in S_4 \setminus H$  with  $\text{ord}(x) = 3$ . Then the cyclic subgroup  $K := \langle x \rangle$  of  $S_4$  intersects  $H$  trivially, and hence  $HK$  is a subgroup of  $S_4$  of order  $|HK| = |H| \cdot |K| = 9$ , which is not possible by Lagrange's theorem since  $9 \nmid 24$ . Therefore,  $S_4$  cannot have a normal subgroup of order 3.  $\square$

**Example 2.14.18.** Let  $G = S_4$ . Then  $|G| = 4! = 24 = 2^3 \cdot 3$ . If  $n_p$  denotes the number of Sylow  $p$ -subgroups of  $G$ , then  $n_2 = 1 + 2k$ , for some integer  $k \geq 0$ , and  $n_2 \mid 3$ . Then  $n_2 = 1$  or 3. Since  $S_4$  cannot have a normal subgroup of order 8 by Exercise 2.14.17 we see that  $n_2 = 3$ . Since  $n_3 = 1 + 3k$ , for some integer  $k \geq 0$  and  $n_3 \mid 8$ , we have  $n_3 \in \{1, 4\}$ . Since  $S_4$  has no normal subgroup of order 8 by Exercise 2.14.17, we have  $n_3 = 4$ .



**Exercise 2.14.19.** Let  $p > 0$  be a prime number. Let  $P$  be a non-trivial  $p$ -subgroup of  $S_p$ . Show that  $|N_{S_p}(P)| = p(p-1)$ .

*Solution:* Since  $|S_p| = p!$ , the largest  $p$ -th power appearing in the prime factorization of  $|S_p|$  is  $p$ . Since  $P$  is a non-trivial  $p$ -subgroup of  $S_p$ , we see that  $|P| = p$ , and hence  $P$  is a Sylow  $p$ -subgroup of  $S_p$ . Since any two Sylow  $p$ -subgroups are conjugates by Sylow's second theorem (Theorem 2.14.11), and since the stabilizer of  $P$  is  $N_{S_p}(P)$ , we see that  $[S_p : N_{S_p}(P)] = n_p$ , the number of Sylow  $p$ -subgroups of  $S_p$ . Since any two distinct Sylow  $p$ -subgroups of  $S_p$  has order  $p$ , they intersect trivially. Let  $\text{Syl}_p(S_p)$  be the set of all Sylow  $p$ -subgroups of  $S_p$ . Note that, any non-identity element of  $P \in \text{Syl}_p(S_p)$  has order  $p$ , and hence is a  $p$ -cycle in  $S_p$ . Since there are  $p-1$  number of non-identity elements in each  $P \in \text{Syl}_p(S_p)$  and there are total  $(p-1)!$  number of distinct  $p$ -cycles in  $S_p$  by Exercise 2.5.15, we conclude that  $n_p(p-1) = (p-1)!$ . Therefore,  $[S_p : N_{S_p}(P)] = n_p = (p-2)!$ , and hence  $|N_{S_p}(P)| = |S_p|/n_p = p(p-1)$ .  $\square$

## 2.15 Miscellaneous Exercises

Let  $G$  be a group.

Q1. Given a subset  $A \subseteq G$ , we define  $N_G(A) := \{a \in G : a^{-1}Aa = A\}$ . Show that

- (i)  $N_G(A)$  is a subgroup of  $G$ .
- (ii) If  $H$  is a subgroup of  $G$ , show that  $H \leq N_G(H)$ .
- (iii) If  $H$  is a subgroup of  $G$ , show that  $N_G(H)$  is the largest subgroup of  $G$  in which  $H$  is normal.
- (iv) Show by an example that  $A$  need not be a subset of  $N_G(A)$ .

Q2. Given a subset  $A$  of  $G$ , let  $C_G(A) := \{a \in G : aba^{-1} = b, \forall b \in A\}$ .

- (i) Show that  $C_G(A)$  is a subgroup of  $G$ .
- (ii) If  $H$  is a subgroup of  $G$ , show that  $H \leq C_G(H)$  if and only if  $H$  is abelian.

Q3. If  $\mathcal{N}$  is a family of normal subgroups of  $G$ , show that  $\bigcap_{N \in \mathcal{N}} N$  is normal in  $G$ .

Q4. If  $N$  is a normal subgroup of  $G$ , show that  $H \cap N$  is normal in  $H$ , for any subgroup  $H$  of  $G$ .

Q5. Let  $N$  be a finite subgroup of  $G$ . Suppose that  $N = \langle S \rangle$  and  $G = \langle T \rangle$ , for some subsets  $S$  and  $T$  of  $G$ . Show that  $N$  is normal in  $G$  if and only if  $tSt^{-1} \subseteq N$ , for all  $t \in T$ .

Q6. Find all normal subgroups of the dihedral group  $D_8 = \langle r, s : \text{ord}(r) = 4, \text{ord}(s) = 2, sr = r^{-1}s \rangle$ , and identify the associated quotient groups.

Q7. Fix an integer  $n \geq 3$ , and let  $D_{2n} = \langle r, s : \text{ord}(r) = n, \text{ord}(s) = 2, sr = r^{-1}s \rangle$  be the dihedral group of degree  $n$  and order  $2n$ .

- (a) Show that

$$Z(D_{2n}) = \begin{cases} \{e\}, & \text{if } n \text{ is odd, and} \\ \{e, r^k\}, & \text{if } n = 2k \text{ is even.} \end{cases}$$

- (b) If  $k \in \mathbb{N}$  divides  $n$ , show that  $\langle r^k \rangle$  is a normal subgroup of  $D_{2n}$ , and the associated quotient group  $D_{2n}/\langle r^k \rangle$  is isomorphic to  $D_{2k}$ .

Q8. Let  $G$  and  $H$  be groups.

- (i) Show that  $\{(a, e_H) : a \in G\}$  is a normal subgroup of  $G \times H$  and the associated quotient group is isomorphic to  $H$ .

- (ii) If  $G$  is abelian, show that the diagonal  $\Delta_G := \{(a, a) : a \in G\}$  of  $G$  is a normal subgroup of  $G \times G$ , and the associated quotient group isomorphic to  $G$ .
- (iii) Show that the diagonal subgroup  $\Delta_{S_3} \subseteq S_3 \times S_3$  is not normal in  $S_3 \times S_3$ .
- Q9. Let  $H$  and  $K$  be subgroups of  $G$  with  $H \leq K$ . Show that  $[G : H] = [G : K][K : H]$ .
- Q10. Let  $G$  be a finite group. Let  $H$  and  $N$  be subgroups of  $G$  with  $N$  normal in  $G$ . If  $\gcd(|H|, [G : N]) = 1$ , show that  $H$  is a subgroup of  $N$ .
- Q11. Let  $N$  be a normal subgroup of a finite group  $G$ . If  $\gcd(|N|, [G : N]) = 1$ , show that  $N$  is the unique subgroup of order  $|N|$  in  $G$ .
- Q12. Let  $H$  be a normal subgroup of  $G$ . Given any subgroup  $K$  of  $G$ , show that  $H \cap K$  is normal in  $HK$ .
- Q13. Show that  $\mathbb{Q}$  has no proper subgroup of finite index. Deduce that  $\mathbb{Q}/\mathbb{Z}$  has no proper subgroup of finite index.
- Q14. Let  $H$  and  $K$  be subgroups of  $G$  with  $[G : H] = m < \infty$  and  $[G : K] = n < \infty$ . Show that  $\text{lcm}(m, n) \leq [G : H \cap K] \leq mn$ . Deduce that  $[G : H \cap K] = [G : H][G : K]$  whenever  $\gcd(m, n) = 1$ .
- Q15. Show that  $S_4$  cannot have normal subgroups of orders 8 and 3.
- Q16. Find the last two digits of  $3^{3^{100}}$ .
- Q17. Let  $H$  and  $K$  be subgroups of  $G$ . If  $H \subseteq N_G(K)$ , then show that
- (i)  $HK$  is a subgroup of  $G$ ,
  - (ii)  $K$  is normal in  $HK$ ,
  - (iii)  $H \cap K$  is normal in  $H$ , and
  - (iv)  $H/(H \cap K) \cong HK/K$ .
- Q18. If  $H$  is a normal subgroup of  $G$  with  $[G : H] = p$ , a prime number, show that for any subgroup  $K$  of  $G$ , either
- (i)  $K$  is a subgroup of  $H$ , or
  - (ii)  $G = HK$  and  $[K : H \cap K] = p$ .
- Q19. Let  $H$  and  $K$  be normal subgroups of  $G$  such that  $G = HK$ . Show that  $G/(H \cap K) \cong (G/H) \times (G/K)$ .
- Q20. Let  $G$  be a finite group of order  $p^r m$ , where  $p > 0$  is a prime number,  $r, m \in \mathbb{N}$  and  $\gcd(p, m) = 1$ . Let  $P$  be a subgroup of order  $p^r$ . Let  $N$  be a normal subgroup of  $G$  of order  $p^s n$ , where  $\gcd(p, n) = 1$ . Show that  $|P \cap N| = p^s$  and  $|PN/N| = p^{r-s}$ . Conclude that intersection of a Sylow  $p$ -subgroup of  $G$  with a normal subgroup  $N$  of  $G$  is a Sylow  $p$ -subgroup of  $N$ .
- Q21. A subgroup  $H$  of a finite group  $G$  is said to be a *Hall subgroup of  $G$*  if its index in  $G$  is relatively prime to its order; i.e., if  $\gcd([G : H], |H|) = 1$ .  
If  $H$  is a Hall subgroup of  $G$  and  $N$  is a normal subgroup of  $G$ , show that  $H \cap N$  is a Hall subgroup of  $N$  and  $HN/N$  is a Hall subgroup of  $G/N$ .
- Q22. A non-trivial abelian group  $G$  is said to be *divisible* if for each  $a \in G$  and non-zero integer  $n \in \mathbb{Z} \setminus \{0\}$ , there exists an element  $b \in G$  such that  $b^n = a$ ; i.e., each element of  $G$  has a  $n$ -th root in  $G$ , for all  $n \in \mathbb{Z} \setminus \{0\}$ . Prove the following.
- (i) Show that  $(\mathbb{Q}, +)$  is a divisible group.
  - (ii) Show that any non-trivial divisible group is infinite.

- (iii) Show by an example that subgroup of a divisible group need not be divisible.
  - (iv) If  $G$  and  $H$  are non-trivial abelian groups, show that  $G \times H$  is divisible if and only if both  $G$  and  $H$  are divisible.
  - (v) Show that quotient of a divisible group by a proper subgroup is divisible.
- Q23. Find all generators and subgroups of  $\mathbb{Z}_{48}$ .
- Q24. Let  $G$  be a group. Given an element  $a \in G$ , show that there is a unique group homomorphism  $f : \mathbb{Z} \rightarrow G$  such that  $f(1) = a$ .
- Q25. Let  $G$  be a group. Let  $a \in G$  be such that  $a^n = e$ , for some integer  $n \geq 0$ , show that there is a unique group homomorphism  $\varphi : \mathbb{Z}_n \rightarrow G$  such that  $\varphi([1]) = a$ .
- Q26. Fix an integer  $n \geq 2$ . Given an integer  $k$ , let  $f_k : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$  be the map defined by  $f_k(x) = x^k$ ,  $\forall x \in \mathbb{Z}_n$ .
- (i) Show that  $f_k$  is a well-defined map.
  - (ii) Show that  $f_k \in \text{Aut}(\mathbb{Z}_n)$  if and only if  $\gcd(n, k) = 1$ .
  - (iii) Show that  $f_k = f_\ell$  if and only if  $\ell \equiv k \pmod{n}$ .
  - (iv) Show that every group automorphism of  $\mathbb{Z}_n$  is of the form  $f_k$ , for some  $k \in \mathbb{Z}$ .
  - (v) Show that  $f_k \circ f_\ell = f_{k\ell}$ ,  $\forall k, \ell \in \mathbb{Z}$ .
  - (vi) Deduce that the map  $f : \mathbb{Z}_n^\times \rightarrow \text{Aut}(\mathbb{Z}_n)$  defined by  $f(k) = f_k$ ,  $\forall k \in \mathbb{Z}_n^\times$ , is an isomorphism of  $\mathbb{Z}_n^\times := U_n$  onto the automorphism group  $\text{Aut}(\mathbb{Z}_n)$ .
  - (vii) Conclude that  $\text{Aut}(\mathbb{Z}_n)$  is an abelian group of order  $\phi(n)$ , where  $\phi$  denotes the Euler phi function.
- Q27. Fix an integer  $n \geq 3$ . Show that the multiplicative group  $G := (\mathbb{Z}/2^n\mathbb{Z})^\times$  has two distinct subgroups of order 2. Conclude that  $G$  is not cyclic.
- Q28. Let  $G$  be a finite group of order  $n$ . Let  $k \in \mathbb{N}$  with  $\gcd(n, k) = 1$ . Use Lagrange's theorem and Cauchy's theorem to show that the map  $f : G \rightarrow G$  defined by  $f(a) = a^k$ ,  $\forall a \in G$ , is surjective.
- Q29. Let  $m, n \geq 2$  be two integers. Find all group homomorphism  $f : \mathbb{Z}_m \rightarrow \mathbb{Z}_n$ .
- Q30. Let  $G$  be a group. Show that there is a one-to-one correspondence between the set of all group homomorphisms from  $\mathbb{Z}_m$  into  $G$  with the set of all solutions of the equations  $x^m = e_G$  in  $G$ .
- Q31. Find the number of group homomorphisms from  $\mathbb{Z}_n$  into  $\mathbb{Z}_m \times \mathbb{Z}_k$ .
- Q32. Find the number of all group homomorphisms from  $S_3$  into  $\mathbb{Z}_n \times \mathbb{Z}_m$ . (*Hint: Use abelianization of  $S_3$ .*)
- Q33. Let  $G$  be a group and  $H$  an abelian subgroup of  $G$ . Show that the subgroup  $\langle H, Z(G) \rangle$  is abelian. Give an example of a group  $G$  and an abelian subgroup  $H$  of  $G$  such that the subgroup  $\langle H, C_G(H) \rangle$  is not abelian, where  $C_G(H) = \{a \in G : a^{-1}ha = h, \forall h \in H\}$  is the centralizer of  $H$  in  $G$ .
- Q34. Show that the subgroup generated by any two distinct elements of order 2 in  $S_3$  is  $S_3$ .
- Q35. Show that any finitely generated subgroup of  $(\mathbb{Q}, +)$  is cyclic. Conclude that  $\mathbb{Q}$  is not finitely generated.
- Q36. Show that the subgroup of  $(\mathbb{Q}^*, \cdot)$  generated by the subset  $\{1/p \in \mathbb{Q}^+ : p \text{ is a prime number}\}$  is  $\mathbb{Q}^+$ , the multiplicative group of positive rational numbers.
- Q37. Show that any group of order 4 is isomorphic to either  $\mathbb{Z}_4$  or  $K_4$ .

Q38. Show that any group of order 6 is isomorphic to either  $\mathbb{Z}_6$  or  $S_3$ .

Q39. Let  $p > 0$  be a prime number, and let

$$G = \{z \in \mathbb{C}^* : z^{p^n} = 1, \text{ for some } n \in \mathbb{N} \cup \{0\}\}.$$

Prove the following.

- (i)  $G$  is a subgroup of  $\mathbb{C}^*$ .
  - (ii) The map  $F_p : G \rightarrow G$  given by  $z \mapsto z^p$ , is a surjective group homomorphism.
  - (iii) Find  $\text{Ker}(F_p)$ .
  - (iv) Show that  $G$  is isomorphic to a *proper quotient group* (i.e., quotient by a non-trivial normal subgroup) of itself.
- Q40. Let  $G$  be the additive group  $(\mathbb{R}, +)$ . Show that  $G$  is isomorphic to the product group  $G \times G$ . (Hint: Note that both  $\mathbb{R}$  and  $\mathbb{R} \times \mathbb{R}$  are  $\mathbb{Q}$ -vector spaces). Show that this fails for  $G = (\mathbb{Z}, +)$ .
- Q41. Let  $G$  be a finite group and let  $S(G)$  be the permutation group on  $G$ . Let  $\pi : G \rightarrow S(G)$  be the *left regular representation* of  $G$  (i.e.,  $\pi$  is the group homomorphism defined by sending  $a \in G$  to the permutation  $\sigma_a \in S(G)$  that sends  $b \in G$  to  $ab \in G$ ).
- (i) If  $a \in G$  with  $\text{ord}(a) = n$  and  $|G| = mn$ , show that  $\pi(a)$  is a product of  $m$  number of  $n$ -cycles.
  - (ii) Deduce that  $\pi(a)$  is an odd permutation if and only if  $\text{ord}(a)$  is even and  $|G|/\text{ord}(a)$  is odd.
  - (iii) If  $\pi(G)$  contains an odd permutation, show that  $G$  has a subgroup of index 2.
- Q42. If  $G$  is a finite group of order  $2n$ , where  $n$  is odd, show that  $G$  has a subgroup of index 2. (Hint: Use Cauchy's theorem and the previous exercise).
- Q43. Let  $G$  be finite group of order  $n$ , where  $n$  is not a prime number. If  $G$  has a subgroup of order  $r$ , for each positive integer  $r$  that divides  $n$ , show that  $G$  is not a simple group.
- Q44. Let  $G$  be a group. A subgroup  $H$  of  $G$  is said to be a *characteristic subgroup* of  $G$  if  $f(H) \subseteq H$ , for all  $f \in \text{Aut}(G)$ . Prove the following.
- (i) Characteristic subgroups are normal.
  - (ii) If  $H$  is the unique subgroup of a given finite order in  $G$ , then  $H$  is a characteristic subgroup of  $G$ .
  - (iii) If  $K$  is a characteristic subgroup of  $H$  and  $H$  is normal in  $G$ , show that  $K$  is normal in  $G$ .
- Q45. Compute the conjugacy class and the stabilizer of  $\sigma := \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 3 & 2 & 5 & 7 & 1 & 6 & 4 \end{pmatrix} \in S_7$ .
- Q46. Let  $H$  be a subgroup of  $G$  with finite index  $[G : H] = n$ . Show that there is a normal subgroup  $K$  of  $G$  with  $K \subseteq H$  and  $[G : K] \leq n!$ .
- Q47. Show that every non-abelian group of order 6 has a non-normal subgroup of order 2. (Hint: Produce an injective group homomorphism  $G \rightarrow S_3$ ). Use this to show that, upto isomorphism, there are only two groups of order 6, namely  $S_3$  and  $\mathbb{Z}_6$ .
- Q48. Let  $P$  be a Sylow  $p$ -subgroup of a finite group  $G$ , and let  $H$  be a subgroup of  $G$ . Prove the following statements.
- (i)  $aPa^{-1} \cap H$  is a Sylow  $p$ -subgroup of  $H$ , for some  $a \in G$ .
  - (ii) If  $P$  is normal in  $G$  then  $H \cap P$  is the unique Sylow  $p$ -subgroup of  $H$ .

- Q49. Let  $G$  be a finite group of order  $pq$ , where  $p, q$  are prime numbers with  $p \leq q$  and  $p \nmid (q-1)$ . Show that  $G$  is abelian. If  $p < q$  and  $p \nmid (q-1)$ , what can you say about  $G$ ?
- Q50. Let  $p > 0$  be a prime number. Let  $P$  be a non-trivial  $p$ -subgroup of  $S_p$ . Show that  $|N_{S_p}(P)| = p(p-1)$ .

## 2.16 Applications of Sylow's Theorems

**Lemma 2.16.1.** *Let  $G$  be a group of order  $pn$ , where  $p$  is a prime number and  $p > n$ . Then  $G$  has a normal subgroup of order  $p$ , and hence is not simple.*

*Proof.* By Cauchy's theorem (Theorem 2.12.14)  $G$  has a subgroup, say  $H$ , of order  $p$ . Let  $K$  be a subgroup of  $G$  of order  $p$ . If  $H \neq K$ , then  $H \cap K = \{e\}$ , and hence the cardinality of the subset  $HK \subseteq G$  is

$$|HK| = \frac{|H| \cdot |K|}{|H \cap K|} = p^2 > pn = |G|,$$

which is a contradiction. Therefore,  $H$  is the unique subgroup of  $G$  of order  $p$ , and hence it is normal in  $G$ . This completes the proof.  $\square$

**Exercise 2.16.2.** Let  $G$  be a finite group and let  $H$  be a proper subgroup of  $G$  of index  $[G : H] = n$ . If  $|G|$  does not divide  $n!$ , show that  $H$  contains a non-trivial normal subgroup of  $G$ , and hence  $G$  is not simple.

*Proof.* Let  $\mathcal{L}_H = \{aH : a \in G\}$  be the set of all left cosets of  $H$  in  $G$ . Note that  $|\mathcal{L}_H| = [G : H] = n$ . Then the left  $G$ -action

$$\sigma : G \times \mathcal{L}_H \rightarrow \mathcal{L}_H, \quad (g, aH) \mapsto gaH,$$

on  $\mathcal{L}_H$  induces a group homomorphism

$$f_\sigma : G \rightarrow S(\mathcal{L}_H) \cong S_n, \quad g \mapsto \sigma(g, -),$$

with  $\text{Ker}(f_\sigma) \subseteq H$  (see Corollary 2.11.17). Since  $|G/\text{Ker}(f_\sigma)|$  divides  $|S_n| = n!$  by Lagrange's theorem (Theorem 2.7.5) and  $|G|$  does not divide  $n!$  by assumption, we have  $\text{Ker}(f_\sigma) \neq \{e\}$ . Since  $H$  is a proper subgroup of  $G$  containing  $\text{Ker}(f_\sigma)$ , it follows that  $G$  is not simple.  $\square$

**Exercise 2.16.3.** Show that any group of order 12 is not simple. (Hint: Use Exercise 2.16.2 with  $H \in \text{Syl}_2(G)$ ).

**Lemma 2.16.4.** *Let  $G$  be a group of order  $p^2q$ , where  $p$  and  $q$  are distinct prime numbers. Then  $G$  is not simple.*

*Proof.* Let  $P \in \text{Syl}_p(G)$  and  $Q \in \text{Syl}_q(G)$ . Suppose that  $p > q$ . Since  $n_p \mid q$  and  $n_p = kp + 1$ , for some integer  $k \geq 0$ , we must have  $n_p = 1$ . So  $P$  is normal in  $G$ .

Suppose that  $p < q$ . If  $n_q = 1$ , then  $Q$  is normal in  $G$ . Assume that  $n_q > 1$ . Then  $n_q = kq + 1$ , for some integer  $k \geq 1$ . Since  $n_q \mid p^2$  and  $n_q > 1$ , either  $n_q = p$  or  $p^2$ . If  $n_q = p$ , then  $p = n_q = 1 + kq > q$ , which contradicts our assumption that  $p < q$ . So  $n_q = p^2$ . Then  $p^2 = n_q = 1 + kq$  implies that  $q$  divides  $p^2 - 1 = (p+1)(p-1)$ . Since  $p < q$ , so  $q \nmid (p-1)$ , and hence  $q \mid (p+1)$ . Since  $p < q$ , this forces  $p = 2$  and  $q = 3$ . Then  $|G| = p^2q = 12$ , and so  $G$  is not simple by Exercise 2.16.3.  $\square$

**Exercise 2.16.5.** Let  $G$  be a group of order 56. Show that  $G$  is not simple.

*Proof.* Given that  $|G| = 56 = 2^3 \times 7$ . Let  $n_2 = |\text{Syl}_2(G)|$  and  $n_7 = |\text{Syl}_7(G)|$ . Then by Sylow's third theorem (Theorem 2.14.12),  $n_2 \mid 7$  and  $n_2 = 1 + 7m$ , for some integer  $m \geq 0$ , and  $n_7 \mid 8$  and

$n_7 = 1 + 7k$ , for some integer  $k \geq 0$ . Then  $n_2 \in \{1, 7\}$  and  $n_7 \in \{1, 8\}$ . If  $n_2 = 1$  or if  $n_7 = 1$ , then  $G$  has a normal Sylow 2-subgroup or a normal Sylow 7-subgroup, and hence  $G$  is not simple. Assume that  $n_2 \neq 1$  and  $n_7 \neq 1$ . Let  $\text{Syl}_2(G) = \{A_1, \dots, A_7\}$  and  $\text{Syl}_7(G) = \{B_1, \dots, B_8\}$ . Since  $|B_i| = 7$ , a prime number, all  $B_i$  are cyclic groups and each of them have 6 elements of order 7. So  $G$  contains  $6 \times 8 = 48$  elements of order 7. Now each  $A_i$  has 8 elements. Since  $B_i \cap B_j$  is a subgroup of both  $B_i$  and  $B_j$ , by Lagrange's theorem we have  $|B_i \cap B_j| \leq 4$ , for  $i \neq j$ . Then  $B_i \cup B_j$  contains at least  $(8+8) - 4 = 12$  elements that has order different from 7. So  $G$  contains at least  $48 + 12 = 60$  elements, which contradicts the fact that  $|G| = 56$ . Therefore, we must have either  $n_2 = 1$  or  $n_7 = 1$ . This completes the proof.  $\square$

Applying the above mentioned results to groups of composite order  $< 60$  we record them into the following Table 2.16.5.1.

$ G $	Reference	$ G $	Reference
$4 = 2^2$	Corollary 2.12.24 & Lemma 2.13.3	$6 = 3 \times 2$	Lemma 2.16.1
$8 = 2^3$	Corollary 2.12.11 & Lemma 2.13.3	$9 = 3^2$	Corollary 2.12.24 & Lemma 2.13.3
$10 = 5 \times 2$	Lemma 2.16.1	$12 = 2^2 \times 3$	Exercise 2.16.3
$14 = 7 \times 2$	Lemma 2.16.1	$15 = 5 \times 3$	Lemma 2.16.1
$16 = 2^4$	Corollary 2.12.11	$18 = 2 \times 9$	Exercise 2.11.21
$20 = 5 \times 4$	Lemma 2.16.1	$21 = 7 \times 3$	Lemma 2.16.1
$22 = 11 \times 2$	Lemma 2.16.1	$24 = 2^3 \times 3$	Exercise 2.16.2 with $H \in \text{Syl}_2(G)$ .
$25 = 5^2$	Corollary 2.12.24 & Lemma 2.13.3	$26 = 13 \times 2$	Lemma 2.16.1
$27 = 3^3$	Corollary 2.12.11 & Lemma 2.13.3	$28 = 7 \times 4$	Lemma 2.16.1
$30 = 2 \times 15$	Exercise 2.11.21	$32 = 2^5$	Corollary 2.12.11 & Lemma 2.13.3
$33 = 11 \times 3$	Lemma 2.16.1	$34 = 17 \times 2$	Lemma 2.16.1
$35 = 7 \times 5$	Lemma 2.16.1	$36 = 3^2 \times 4$	Exercise 2.16.2
$38 = 19 \times 2$	Lemma 2.16.1	$39 = 13 \times 3$	Lemma 2.16.1
$40 = 5 \times 8$	$n_5 = 1$ & Corollary 2.14.15	$42 = 7 \times 6$	Lemma 2.16.1
$44 = 11 \times 4$	Lemma 2.16.1	$45 = 3^2 \times 5$	Lemma 2.16.4
$46 = 23 \times 2$	Lemma 2.16.1	$48 = 2^4 \times 3$	Exercise 2.16.2 with $H \in \text{Syl}_2(G)$
$49 = 7^2$	Corollary 2.12.24 & Lemma 2.13.3	$50 = 2 \times 25$	Exercise 2.11.21
$51 = 17 \times 3$	Lemma 2.16.1	$52 = 13 \times 4$	Lemma 2.16.1
$54 = 2 \times 27$	Exercise 2.11.21	$55 = 11 \times 5$	Lemma 2.16.1
56	Exercise 2.16.5	$57 = 19 \times 3$	Lemma 2.16.1
$58 = 29 \times 2$	Lemma 2.16.1		

TABLE 2.16.5.1: Non-simple groups of order  $< 60$

Thus, we have the following.

**Theorem 2.16.6.** *Let  $G$  be a finite group of order  $n$ , where  $2 \leq n < 60$  and  $n$  is not a prime number. Then  $G$  is not simple.*

In Theorem 2.13.12 we have shown that  $A_5$  is a simple group of order 60. Now we show that, upto isomorphism,  $A_5$  is the only simple group of order 60. We break this problem into a set of small exercises.

**Exercise 2.16.7.** Let  $H$  be a subgroup of  $G$  with  $[G : H] = 2$ .

- (i) Show that  $a^2 \in H$ ,  $\forall a \in G$ .
- (ii) Show that  $H$  contains all elements of  $G$  of odd order.

*Proof.* (i) Since  $H$  is normal in  $G$  by Proposition 2.8.7, the quotient group  $G/H$  has order 2, and so given any element  $a \in G$ , it follows from Corollary 2.7.7 that  $a^2H = (aH)^2 = H$  in  $G/H$ , and hence  $a^2 \in H$ .

(ii) Let  $a \in G$  with  $\text{ord}(a) = 2n + 1$ , for some integer  $n \geq 0$ . Then in the quotient group  $G/H$  we have  $H = a^{2n+1}H = (a^2H)^n aH = H \cdot aH = aH$  by part (i), and hence  $a \in H$ .  $\square$

**Exercise 2.16.8.** Fix an integer  $n \geq 3$ . If  $H$  is a subgroup of  $S_n$  with  $[S_n : H] = 2$ , then  $H = A_n$ .

*Proof.* Clearly the index of  $A_n$  in  $S_n$  is 2. Let  $H$  be a subgroup of  $S_n$  with  $[S_n : H] = 2$ . Then  $H$  is normal in  $S_n$  by Proposition 2.8.7. Let  $\sigma \in S_n$  be any 3-cycle. Since  $\text{ord}(\sigma) = 3$ , an odd number, we have  $\sigma \in H$  by Exercise 2.16.7. Since  $A_n$  is generated by 3-cycles in  $S_n$  by Exercise 2.5.34, we have  $A_n \subseteq H$ . Since both  $A_n$  and  $H$  have the same cardinality, namely  $n!/2$ , we conclude that  $H = A_n$ .  $\square$

**Exercise 2.16.9.** Let  $G$  be a finite simple group. Let  $p > 0$  be a prime number such that  $p \mid |G|$ . If the number of all Sylow  $p$ -subgroups of  $G$  is  $n \geq 2$ , show that  $G$  is isomorphic to a subgroup of  $S_n$ .

*Proof.* Let  $X_p$  be the set of all Sylow  $p$ -subgroups of  $G$ . Since conjugate of a Sylow  $p$ -subgroup of  $G$  is again a Sylow  $p$ -subgroup of  $G$  (see Exercise 2.14.8), the conjugation action of  $G$  on  $X_p$  gives rise to a group homomorphism  $\varphi : G \rightarrow S(X_p) \cong S_n$  defined by sending  $g \in G$  to the permutation

$$\varphi_g : X_p \rightarrow X_p, \quad P \mapsto gPg^{-1}.$$

If  $\text{Ker}(\varphi) = G$ , then  $\varphi_g = \text{Id}_{X_p}$ ,  $\forall g \in G$ , and so given any  $P \in X_p$  we have  $P = gPg^{-1}$ ,  $\forall g \in G$ . Since any two Sylow  $p$ -subgroups are conjugates, we must have  $|X_p| = 1$ , which contradicts our assumption that  $|X_p| = n \geq 2$ . Since  $\text{Ker}(\varphi)$  is a normal subgroup of  $G$  and  $G$  is simple, we must have  $\text{Ker}(\varphi) = \{e\}$ . Therefore,  $\varphi : G \rightarrow S_n$  is an injective group homomorphism. Hence the result follows.  $\square$

**Exercise 2.16.10.** Show that  $A_5$  has a subgroup isomorphic to  $A_4$ .

*Proof.* Define a map  $\varphi : A_4 \rightarrow S_5$  by sending  $\sigma \in A_4$  to  $\varphi(\sigma) = \tilde{\sigma}$ , where

$$\tilde{\sigma}(i) = \begin{cases} \sigma(i), & \text{if } 1 \leq i \leq 4, \\ i, & \text{if } i = 5. \end{cases}$$

Clearly  $\tilde{\sigma} \in S_5$ . Moreover, from decomposition of  $\sigma$  into product of transpositions it follows immediately that  $\tilde{\sigma}$  is even, and hence in  $A_5$ . It is easy to check that  $\varphi$  is an injective group homomorphism, and hence  $\varphi$  identifies  $A_4$  as a subgroup of  $A_5$ .  $\square$

**Exercise 2.16.11.** Let  $G$  be a simple group of order 60. Show that  $G$  has a subgroup of order 12.

*Proof.* Since  $|G| = 60 = 2^2 \times 3 \times 5$ , by Sylow's third theorem (Theorem 2.14.12)  $n_2 \mid 15$  and  $n_2 = 1 + 2k$ , for some integer  $k \geq 0$ . Then  $n_2 \in \{1, 3, 5, 15\}$ . Since  $G$  is simple,  $n_2 \neq 1$  by Corollary 2.14.15. Since  $|G| = 60 > 3!$ , it follows from Exercise 2.16.9 that  $n_2 \neq 3$ . If  $n_2 = 5$ , then again by Exercise 2.16.9  $G$  is isomorphic to a subgroup of  $S_5$ . Since the only index 2 subgroup of  $S_5$  is  $A_5$ , we have  $G \cong A_5$ . Then  $G$  has a subgroup of order 12 by Exercise 2.16.10.

Suppose that  $n_2 = 15$ . Let  $\{P_1, \dots, P_{15}\}$  be the set of all Sylow 2-subgroups of  $G$ . Note that each  $P_i$  has order 4. Since  $|G| = 60 = 2^2 \times 3 \times 5$ , by Sylow's third theorem (Theorem 2.14.12)

we have  $n_5 \mid 12$  and  $n_5 = 1 + 5k$ , for some integer  $k \geq 0$ . Then  $n_5 \in \{1, 6\}$ . Since  $G$  is simple,  $n_5 \neq 1$ , and so  $n_5 = 6$ . Let  $\{Q_1, \dots, Q_6\}$  be the set of all Sylow 5-subgroups of  $G$ . Since each  $Q_i$  has order 5 and  $Q_i \cap Q_j = \{e\}$ , for  $i \neq j$ , we see that  $G$  has  $(5 - 1) \times 6 = 24$  elements of order 5.

If  $B_i \cap B_j = \{e\}$ , for all  $i \neq j$ , then  $\bigcup_{i=1}^{15} P_i$  contains  $(4 - 1) \times 15 + 1 = 46$  elements of order different from 5. Thus we get at least  $24 + 46 = 70$  elements in  $G$  which contradicts our assumption that  $|G| = 60$ . Therefore, there exists at least a pair of distinct elements  $i, j \in I_{15}$  such that  $P_i \cap P_j \neq \{e\}$ . Then  $|P_i \cap P_j| = 2$  by Lagrange's theorem, and hence  $P_i \cap P_j$ , being an index 2 subgroup of both  $P_i$  and  $P_j$ , is normal in both  $P_i$  and  $P_j$  (see Proposition 2.8.7). Therefore,  $P_i, P_j \subseteq N_G(P_i \cap P_j)$ , and so  $P_i P_j \subseteq N(P_i \cap P_j)$ . Therefore,  $|P_i \cap P_j| \geq |P_i P_j| = \frac{4 \times 4}{2} = 8$ . Since  $N(P_i \cap P_j)$  is a subgroup of  $G$ , by Lagrange's theorem (Theorem 2.7.5) we see that  $|N(P_i \cap P_j)| \in \{12, 20, 30, 60\}$ . Since  $G$  is simple and  $P_i \cap P_j$  is a non-trivial proper normal subgroup of  $G$ ,  $|N(P_i \cap P_j)| \neq 60$ . Since  $G$  is simple and any index 2 subgroup of  $G$  is normal in  $G$  (see Proposition 2.8.7),  $|N(P_i \cap P_j)| \neq 30$ . Again  $|N(P_i \cap P_j)| \neq 20$  by Exercise 2.16.2. Therefore,  $|N(P_i \cap P_j)| = 12$ , as required.  $\square$

**Theorem 2.16.12.** *Any simple group of order 60 is isomorphic to  $A_5$ .*

*Proof.* Let  $G$  be a simple group of order 60. Then  $G$  has a subgroup, say  $H$ , of order 12 by Exercise 2.16.11. Since  $[G : H] = 5$ , the natural left  $G$ -action on the set  $X = \{aH : a \in G\}$  gives rise to a group homomorphism  $\varphi : G \rightarrow S(X) \cong S_5$  with  $\text{Ker}(\varphi) \subseteq H$ . Since  $H$  is a proper subgroup of  $G$  and  $G$  is simple, we must have  $\text{Ker}(\varphi) = \{e\}$ . Since  $|G| = 60$ , it follows that  $G$  is isomorphic to an index 2 subgroup of  $S_5$ , and hence is isomorphic to  $A_5$  by Exercise 2.16.8. This completes the proof.  $\square$

### 2.16.1 Quaternion group $Q_8$

Consider the set  $Q_8 := \{1, -1, i, -i, j, -j, k, -k\}$ , together with the binary operation  $\cdot : Q_8 \times Q_8 \rightarrow Q_8$  given by the following law:

$$\begin{aligned} 1 \cdot a &= a \cdot 1 = a, \quad \forall a \in Q_8. \\ (-1) \cdot (-1) &= 1, \text{ and } (-1) \cdot a = a \cdot (-1) = -a, \quad \forall a \in Q_8, \\ i \cdot i &= j \cdot j = k \cdot k = -1, \\ i \cdot j &= k, \quad j \cdot i = -k, \\ j \cdot k &= i, \quad k \cdot j = -i, \\ k \cdot i &= j, \quad i \cdot k = -j. \end{aligned}$$

Verify that the set  $Q_8$  together with the binary operation as defined above forms a group, known as the *quaternion group*.

**Exercise 2.16.13.** Show that the quaternion group  $Q_8$  has following presentations:

- (i)  $Q_8 = \langle i, j, k, \ell : i^2 = j^2 = k^2 = ijk = \ell \text{ and } \ell^2 = e \rangle$ .
- (ii)  $Q_8 = \langle x, y : x^4 = 1, x^2 = y^2, y^{-1}xy = x^{-1} \rangle$ .

**Exercise 2.16.14.** (i) Show that  $Q_8$  is a non-abelian group.

(ii) Find all subgroups of  $Q_8$  and draw the associated lattice diagram (see Definition 2.4.13).

(iii) Show that all subgroups of  $Q_8$  are normal.



**Exercise 2.16.15.** Consider the presentation of  $Q_8 = \langle x, y : x^4 = 1, x^2 = y^2, y^{-1}xy = x^{-1} \rangle$ . Show that the map  $\varphi : Q_8 \rightarrow \text{GL}_2(\mathbb{C})$  defined on the generators of  $Q_8$  by

$$\varphi(x) = \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix} \quad \text{and} \quad \varphi(y) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

extends to an injective group homomorphism that identifies  $Q_8$  as a subgroup of  $\text{GL}_2(\mathbb{C})$ .

**Exercise 2.16.16.** Show that the subgroup of  $\text{SL}_2(\mathbb{Z}_3)$  generated by the matrices

$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

is isomorphic to the quaternion group  $Q_8$ .

**Exercise 2.16.17.** Show that the subgroup of  $\text{SL}_2(\mathbb{Z}_5)$  generated by

$$A = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

is isomorphic to the quaternion group  $Q_8$ .

**\*Exercise 2.16.18.** (i) Show that  $G := \text{SL}_2(\mathbb{Z}_5)$  is a finite group of order 120.

(ii) Show that the subgroup  $H$  of  $G$  generated by

$$A = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

is isomorphic to the quaternion group  $Q_8$ .

(iii) Let  $C := \begin{pmatrix} 2 & 2 \\ 1 & -1 \end{pmatrix} \in \text{SL}_2(\mathbb{Z}_5)$ . Show that  $C \in N_G(H)$  and the subgroup  $K = \langle A, B, C \rangle$  of  $G$  has index 5 in  $G$ .

(iv) Show that the conjugation action of  $G$  on the set of all left cosets  $\mathcal{L}_{K,G} = \{aK : a \in G\}$  is transitive, and the associated group homomorphism  $\varphi : G \rightarrow S(\mathcal{L}_K) \cong S_5$  has kernel  $\text{Ker}(\varphi) = Z(\text{SL}_2(\mathbb{Z}_5))$ .

(v) Use Corollary 2.11.14 to conclude that this gives rise to an injective group homomorphism  $\tilde{\varphi} : \text{PSL}_2(\mathbb{Z}_5) \rightarrow S_5$ .

(vi) Show that the image of  $\tilde{\varphi}$  has index 2 in  $S_5$ , and hence  $\text{PSL}_2(\mathbb{Z}_5)$  is isomorphic to  $A_5$ .

## Projective spaces\*

Let  $k$  be a field. Given an integer  $n \geq 1$ , consider the  $(n+1)$ -dimensional  $k$ -vector space  $V = k^{n+1}$ . On the set of all non-zero vectors  $V \setminus \{0\}$ , we define a relation  $\sim$  as follows: given  $v, w \in V \setminus \{0\}$ , we define

$$v \sim w \quad \text{if} \quad \exists \lambda \in k \setminus \{0\} \quad \text{such that} \quad w = \lambda v.$$

Verify that  $\sim$  is an equivalence relation on  $V \setminus \{0\}$ . We denote by  $\mathbb{P}^n(k) = (V \setminus \{0\}) / \sim$ , the set of all  $\sim$ -equivalence classes of non-zero vectors in  $V$ , and call it the *projective  $n$ -space over  $k$* . Note that an element of  $\mathbb{P}^n(k)$  is an one-dimensional  $k$ -linear subspace (a straight-line in  $V = k^{n+1}$  passing through the origin), i.e.,

$$\ell_v := \{\lambda v : \lambda \in k\} \subset V,$$

for some  $v \in V \setminus \{0\}$ . Given non-zero vectors  $v, w \in V \setminus \{0\}$ , we have  $\ell_v = \ell_w$  if and only if  $w = \lambda v$ , for some  $\lambda \in k \setminus \{0\}$ .

**Exercise 2.16.19.** (i) Let  $\mathbb{F}_q$  be a finite field of order  $q = p^n$ , where  $p > 0$  is a prime number and  $n \in \mathbb{N}$ . Show that the projective line  $\mathbb{P}^1(\mathbb{F}_q)$  has  $q + 1$  elements. Find the cardinality of the projective  $n$ -space  $\mathbb{P}^n(\mathbb{F}_q)$ , for all  $n \in \mathbb{N}$ .

(ii) Let  $k$  be a field. Show that the natural map  $\sigma : \mathrm{GL}_{n+1}(k) \times \mathbb{P}^n(k) \rightarrow \mathbb{P}^n(k)$  given by

$$\sigma(A, \ell_v) = \ell_{Av}, \quad \forall (A, \ell_v) \in \mathrm{GL}_{n+1}(k) \times \mathbb{P}^n(k),$$

is well-defined and is a left  $\mathrm{GL}_{n+1}(k)$ -action on  $\mathbb{P}^n(k)$ .

(iii) Show that  $\sigma$  is not faithful (c.f. Definition 2.11.7).

(iv) Show that  $\sigma$  gives rise to  $\mathrm{PGL}_{n+1}(k)$ -action, and hence a  $\mathrm{PSL}_{n+1}(k)$ -action, on  $\mathbb{P}^n(k)$ .

**\*Exercise 2.16.20.** (i) Show that the natural  $\mathrm{PSL}_2(\mathbb{Z}_5)$ -action on the projective line  $\mathbb{P}^1(\mathbb{Z}_5)$  gives rise to a group homomorphism  $f : \mathrm{SL}_2(\mathbb{Z}_5) \rightarrow S_6$ .

(ii) Show that the image of  $f$  is isomorphic to  $S_5$ .

(iii) Show that  $f(A) \in S_5$  is even, for all  $A \in \mathrm{PSL}_2(\mathbb{Z}_5)$ .

(iv) Conclude that  $\mathrm{PSL}_2(\mathbb{Z}_5) \cong A_5$ , and hence is simple.

## 2.17 Finitely Generated Abelian Groups

See page 42–44 of the Algebra book by S. Lang [Lan02].

**Lemma 2.17.1.** Let  $A$  be a finite abelian  $p$ -group. Let  $a_1 \in A$  be such that  $\mathrm{ord}(a_1) = p^{r_1} \geq \mathrm{ord}(a)$ ,  $\forall a \in A$ . Let  $A_1 = \langle a_1 \rangle$  and let  $\pi : A \rightarrow A/A_1$  be the quotient map. Then for given  $w \in A/A_1$ , there exists  $a \in A$  such that  $\mathrm{ord}(a) = \mathrm{ord}(w)$  and  $\pi(a) = w$  in  $A/A_1$ .

**Lemma 2.17.2.** Let  $A$  be a finite abelian  $p$ -group, where  $p$  is a prime number. Then there exists a unique finite sequence of integers  $(\alpha_1, \dots, \alpha_k)$  satisfying  $\alpha_1 \geq \dots \geq \alpha_k \geq 1$  and  $\alpha_1 + \dots + \alpha_k = |A|$  such that  $A \cong \mathbb{Z}_{p^{\alpha_1}} \times \dots \times \mathbb{Z}_{p^{\alpha_k}}$ .

*Proof.* We prove this by induction on  $|A| = p^n$ . Let  $a_1 \in A$  be of maximal order; i.e.,  $\mathrm{ord}(a_1) \geq \mathrm{ord}(a)$ ,  $\forall a \in A$ . Let  $A_1 = \langle a_1 \rangle$ . Suppose that  $\mathrm{ord}(a_1) = p^{r_1}$ , for some integer  $r_1 \leq n$ . Now we need the following lemma. □

**Theorem 2.17.3.** Let  $G$  be a finite abelian group of order  $n > 1$ . Let  $n = p_1^{\alpha_1} \dots p_k^{\alpha_k}$  be the unique factorization of  $n$  into product of distinct prime powers. Then we have the following.

(i)  $G \cong A_1 \times \dots \times A_k$ , where  $A_i$  is an abelian group of order  $p_i^{\alpha_i}$ , for all  $i \in \{1, \dots, k\}$ .

(ii) For each  $A \in \{A_1, \dots, A_k\}$ , with  $|A| = p^\alpha$ , we have

$$A \cong \mathbb{Z}_{p^{\beta_1}} \times \dots \times \mathbb{Z}_{p^{\beta_t}},$$

with  $\beta_1 \geq \beta_2 \geq \dots \geq \beta_t \geq 1$  and  $\beta_1 + \dots + \beta_t = \alpha$  (where  $t$  and  $\beta_1, \dots, \beta_t$  depends on  $i$ ).

(iii) The decomposition in (i) and (ii) is unique; i.e., if  $G \cong B_1 \times \dots \times B_k$  with  $|B_i| = p_i^{\alpha_i}$ , for all  $i \in \{1, \dots, k\}$ , then  $B_i \cong A_i$ , for all  $i$ .

*Proof.* Since  $G$  is abelian, any subgroup of  $G$  is normal. Then by Sylow's second theorem (Theorem 2.14.11)  $G$  has a unique Sylow  $p_i$ -subgroup, say  $A_i$ , for each  $i \in \{1, \dots, k\}$ . Since  $|A_i| = p_i^{\alpha_i}$ , for all  $i$ , and  $p_1, \dots, p_k$  are distinct prime numbers, we have  $A_i \cap (\prod_{j \neq i} A_j) = \{e\}$ , for all  $i$ . Since

$|\prod_{i=1}^k A_i| = n = |G|$ , we have  $G = A_1 \cdots A_k$ . Then  $G \cong A_1 \times \cdots \times A_k$  by Theorem 2.10.18. This proves the first part.

For part (ii), we use induction on  $|A|$ . Let  $E$  be an elementary abelian  $p$ -group (i.e.,  $a^p = e$ ,  $\forall a \in E$ ). Then by Lemma ?? we have  $\square$

Recall that a group  $G$  is said to be *finitely generated* if there is a finite subset  $S \subseteq G$  such that  $\langle S \rangle = G$ . For example, finite groups, cyclic groups are finitely generated.

**Definition 2.17.4.** Given an integer  $r \geq 1$ , let  $\mathbb{Z}^r$  be the direct product of  $r$  copies of  $\mathbb{Z}$ . A group  $G$  is said to be a *free abelian group of rank  $r$*  if  $G$  is isomorphic to  $\mathbb{Z}^r$ .

## 2.18 Free Group

## 2.19 Solvable & Nilpotent Groups

## 2.20 Semi-direct product

Let  $H$  and  $K$  be groups. We say that  $K$  *acts on  $H$  by automorphisms* if there is a group homomorphism  $f : K \rightarrow \text{Aut}(H)$ , where  $\text{Aut}(H)$  is the group of all automorphisms of  $H$ . To simplify the notation, we denote by  $f_k$  the automorphism  $f(k) \in \text{Aut}(H)$ .

On the Cartesian product  $H \times K$  of  $H$  with  $K$ , we define a binary operation by setting

$$(2.20.1) \quad (h_1, k_1) \cdot (h_2, k_2) := (h_1 f_{k_1}(h_2), k_1 k_2), \quad \forall (h_1, k_1), (h_2, k_2) \in H \times K.$$

Note that if  $f$  is the trivial homomorphism, then  $f_k(h) = h$ ,  $\forall h \in H, k \in K$ , and in that case the above binary operation become the component-wise binary operation of the direct product group  $H \times K$ . Given  $(h_1, k_1), (h_2, k_2), (h_3, k_3) \in H \times K$ , we have

$$\begin{aligned} ((h_1, k_1)(h_2, k_2))(h_3, k_3) &= (h_1 f_{k_1}(h_2), k_1 k_2)(h_3, k_3) \\ &= (h_1 f_{k_1}(h_2) f_{k_1 k_2}(h_3), (k_1 k_2) k_3) \\ &= (h_1 f_{k_1}(h_2) f_{k_1}(f_{k_2}(h_3)), k_1(k_2 k_3)) \\ &= (h_1 f_{k_1}(h_2 f_{k_2}(h_3)), k_1(k_2 k_3)) \\ &= (h_1, k_1)(h_2 f_{k_2}(h_3), k_2 k_3) \\ &= (h_1, k_1)((h_2, k_2)(h_3, k_3)). \end{aligned}$$

Therefore, the binary operation on  $H \times K$  defined in (2.20.1) is associative. Note that given  $(h, k) \in H \times K$ , we have

$$\begin{aligned} (h, k)(e_H, e_K) &= (h f_k(e_H), k e_K) = (h e_H, k) = (h, k), \\ \text{and } (e_H, e_K)(h, k) &= (e_H f_{e_K}(h), e_K k) = (h, k), \end{aligned}$$

where  $e_H \in H$  and  $e_K \in K$  are the neutral elements of  $H$  and  $K$ , respectively. Finally, given  $(h, k) \in H \times K$ , we have

$$\begin{aligned}
 (h, k)(f_{k^{-1}}(h^{-1}), k^{-1}) &= (hf_k(f_{k^{-1}}(h^{-1})), kk^{-1}) \\
 &= (h(f_k \circ f_k^{-1})(h^{-1}), e_K) \\
 &= (hh^{-1}, e_K) \\
 &= (e_H, e_K), \\
 \text{and } (f_{k^{-1}}(h^{-1}), k^{-1})(h, k) &= (f_{k^{-1}}(h^{-1})f_{k^{-1}}(h), k^{-1}k) \\
 &= (f_{k^{-1}}(h^{-1}h), e_K) \\
 &= (f_{k^{-1}}(e_H), e_K) \\
 &= (e_H, e_K).
 \end{aligned}$$

Therefore,  $(h, k)^{-1} = (f_{k^{-1}}(h^{-1}), k^{-1})$ ,  $\forall (h, k) \in H \times K$ . Therefore, the binary operation (2.20.1) on  $H \times K$  makes it a group, called the *semidirect product of  $H$  with  $K$  along  $f$* , and is denoted by  $H \rtimes_f K$  or simply by  $H \rtimes K$ , if there is no confusion about  $f$ .

## 2.21 Linear Groups

# Bibliography

- [DF04] David S. Dummit and Richard M. Foote, Abstract algebra. Third edition. *John Wiley & Sons, Inc., Hoboken, NJ*, 2004. xii+932 pp.
- [MMS97] D. S. Malik, J. M. Mordeson and M. K. Sen, Fundamentals of Abstract Algebra, *International series in pure and applied mathematics*, McGraw-Hill, 1997.
- [Lan02] Serge Lang, Algebra. *Grad. Texts in Math.*, **211**. Springer-Verlag, New York, 2002, xvi+914 pp.
- [Art91] Michael Artin, Algebra. *Prentice Hall, Inc., Englewood Cliffs, NJ*, 1991, xviii+618 pp.