

Slow Dynamics and High Variability in Clustered Networks

Reproduction of Litwin-Kumar and Doiron (2012)

Arjun Puri*

Johan Kirikal*

Shuo Wang*

February 5, 2026

Abstract

This report reproduces the key findings of Litwin-Kumar and Doiron (2012) on slow dynamics in clustered cortical networks. We implement a network of leaky-integrate-and-fire neurons with increased connection probabilities based on neurons cluster assignment. We examine the effects of clustering on the network's firing patterns through measures like Fano factor, correlation coefficients, and firing rate distributions.

1. Introduction

Cortical networks are often modeled as unstructured, randomly connected networks with balanced excitation and inhibition. This balance can place the network in an irregular, asynchronous regime in which neurons exhibit high trial-to-trial variability in spike timing. In this regime, excitation and inhibition are large but cancel in the mean. Thus, spikes are often fluctuation-driven, resulting in high spike-time variability.

However, cortical networks also exhibit slower, population-level changes in firing rate over longer timescales (hundreds of milliseconds to seconds). Baseline, unstructured, balanced excitation-inhibition networks do not typically capture these slower fluctuations in firing rate.

In this study, we find that introducing clustered excitatory connectivity creates slow dynamics. Clusters transition between states of high and low firing, such that neurons exhibit both high spike-time variability and slow firing-rate fluctuations. Introducing stimuli can bias the network toward particular activity states, reducing variability.

2. Results

We present our reproduction of the key findings from the original study.

*Berlin Center for Computational Neuroscience, alphabetical order

2.1. Metastable Dynamics in Clustered Networks

An excitatory-inhibitory (E/I) network with clustered E→E connectivity produces metastable “high” firing activity states within clusters that are stable yet quick to transition. These high activity states persist for hundreds of milliseconds to seconds before switching, leading to slow rate dynamics and elevated trial-to-trial spike count variability (super-Poisson) compared to unclustered networks.

2.1.1 Timescales

Clustered networks exhibit substantially longer correlation times than unstructured networks. The autocovariance function decays much more slowly as a function of lag in clustered networks compared to their unstructured counterparts. Similarly, the cross-covariance for within-cluster neuron pairs decays slowly, indicating a shared correlated rate change among neurons belonging to the same cluster.

2.1.2 Variability

Neurons in the clustered network exhibit Fano factors greater than one (“super-Poisson” spike count variability), which increase with the size of the counting window. This pattern illustrates the presence of slow firing rate variability in clustered networks.




Figure 1. Distribution of Fano factors across neurons for uniform and clustered networks. The clustered network exhibits a broader distribution with a higher mean (dashed lines), indicating greater spike count variability.




Figure 2. Mean Fano factor as a function of counting window size. In the clustered network, the Fano factor increases with window size, reflecting slow rate fluctuations. The uniform network remains near the Poisson baseline ($FF = 1$).

2.1.3 Correlations

Pairwise spike count correlations remain near zero on average for both network types. However, in the clustered network, within-cluster neuron pairs show a heavier positive tail in the distribution of correlation coefficients. This observation is consistent with the notion that clusters undergo additional correlated firing rate transitions—switching between high and low activity states—when examined across temporal windows spanning the simulation. Furthermore, this positive tail becomes increasingly pronounced as the counting window size increases, as expected given that within-cluster neurons share these slow rate fluctuations.




Figure 3. Distribution of pairwise spike count correlations for all neuron pairs (left) and within-cluster pairs only (right). While correlations remain centered near zero on average, within-cluster pairs in the clustered network show a heavy positive tail that increases with counting window size (50 ms vs. 100 ms).

2.1.4 Mean Rate

Despite the switching between high and low firing states, the clustered network exhibits a low mean firing rate similar to that of the unstructured network. This is likely due to additional suppression within the network during high firing activity states. The high activity epochs in clustered networks do not alter the overall mean firing rate; however, the firing rate distribution broadens somewhat, likely reflecting the alternation between high and low activity states.




Figure 4. Distribution of firing rates for excitatory neurons in uniform and clustered networks. Both networks maintain low mean firing rates, though the clustered network shows a broader distribution reflecting the alternation between high and low activity states.

2.2. Critical Cluster Strength




Figure 5. Mean Fano factor as a function of cluster strength (R_{EE}). As within-cluster connection strength increases, the network transitions from Poisson-like variability to super-Poisson variability.

3. Discussion

Our reproduction confirms that clustered connectivity generates slow dynamics through metastable attractor states.

3.1. Interpretation

The clustering mechanism provides a parsimonious explanation for slow cortical fluctuations.

3.2. Limitations

The model uses simplified neurons and imposed rather than emergent clustering.

3.3. Future Directions

Promising extensions include plasticity mechanisms and functional implications.

4. Extensions

Beyond reproduction, we explore extensions to the clustered network model.

4.1. Extension 1

Description of the first extension and its motivation.

4.2. Extension 2

Description of the second extension and its motivation.

5. Model

We implement a recurrent spiking network with clustered excitatory connectivity using Brian2.

5.1. Network Architecture

The network consists of excitatory and inhibitory leaky integrate-and-fire neurons with structured connectivity.

5.2. Neuron Model

Membrane potential dynamics follow a leaky integrate-and-fire model with exponential synaptic filtering.

5.3. Connectivity

Connection probability between excitatory neurons depends on cluster membership, with enhanced within-cluster connectivity.

5.4. Parameters

Model parameters are chosen to match those of the original study.

References