

Elementary Number Theory: Divisibility Theory in the Integers

Arjun Vardhan

†

Created: 5th April 2022

Last updated: 7th April 2022

1 Division Algorithm

- **Division Algorithm:** Let $a, b \in \mathbb{Z}$, $b > 0$. Then there exist unique integers r, q such that $a = qb + r$, $0 \leq r < b$. *Proof:* Let $S = \{a - xb : x \in \mathbb{Z}, a - xb \geq 0\}$. Since $b \geq 1$, $|a|b \geq |a|$, and so $a - (-|a|)b = a + |a|b \geq a + |a| \geq 0$. Thus S is nonempty. By the well ordering principle, S must have a least element r . By the definition of S , there exists $q \in \mathbb{Z}$ such that $r = a - qb$, $r \geq 0$. Suppose $r \geq b$. Then $a - (q+1)b = (a - qb) - b = r - b \geq 0$. Thus $a - (q+1)b \in S$, but since r is the least element of S , this is a contradiction. So $r < b$. Now, suppose that $a = qb + r = q'b + r'$, where $0 \leq r < b$, $0 \leq r' < b$. Then $r' - r = b(q - q')$ and so $|r - r'| = b|q - q'|$. On adding the inequalities $-b < -r \leq 0$ and $0 \leq r' < b$, we get $-b < r' - r < b$, or $|r' - r| < b$. Thus $b|q - q'| < b$, implying that $0 \leq |q - q'| < 1$. So $q - q' = 0$ and thus $r - r' = 0$. Thus q and r are unique. ■
- **Corollary:** Let $a, b \in \mathbb{Z}$, $b \neq 0$. Then there exist unique integers r, q such that $a = qb + r$, $0 \leq r < |b|$. *Proof:* Let $b < 0$. Then $|b| > 0$, and by the division algorithm there exist unique integers q' and r such that $a = q'|b| + r$. Since $|b| = -b$, let $q = -q'$ to get $a = qb + r$, with $0 \leq r < |b|$. ■

2 Greatest Common Divisor

- Let $a, b \in \mathbb{Z}$, $a \neq 0$. b is said to be divisible by a , denoted $a \mid b$ if there exists $c \in \mathbb{Z}$ such that $b = ac$.
- **Let $a, b, c \in \mathbb{Z}$. Then:**
 1. $a \mid 0$, $1 \mid a$, and $a \mid a$. *Proof:* $0 = 0 \times a$, $1 = 1 \times a$ and $a = 1 \times a$. ■
 2. $a \mid 1$ if and only if $a = \pm 1$. *Proof:* Suppose $a \mid 1$. Then $1 = na$ for some $n \in \mathbb{Z}$. Let $|a| > 1$. Since $n \neq 0$, $|na| > 1$, which is a contradiction. So $|a| = 1$, and thus $a = \pm 1$. Conversely, suppose $a = +1$. Then $1 = 1 \times 1 = (-1) \times (-1)$. ■
 3. If $a \mid b$ and $c \mid d$, then $ac \mid bd$. *Proof:* There exist integers m, n such that $b = am$ and $d = cn$. Then $ac(mn) = bd$. ■
 4. If $a \mid b$ and $b \mid c$, then $a \mid c$. *Proof:* There exist integers m, n such that $b = am$ and $c = bn$. Then $c = a(mn)$. ■
 5. $a \mid b$ and $b \mid a$ if and only if $a = \pm b$. *Proof:* Suppose $a \mid b$ and $b \mid a$. There exist integers m, n such that $a = bm$ and $b = an$. Thus $a = amn$, implying $mn = 1$. So $m = n = \pm 1$ and thus $a = \pm b$. The converse is obvious. ■
 6. If $a \mid b$ and $b \neq 0$, then $|a| \leq |b|$. *Proof:* There exists an integer c such that $b = ac$. Since $b \neq 0$, $c \neq 0$. So $|b| = |a||c|$. Since $c \neq 0$, $|c| \geq 1$ and thus $|b| = |a||c| \geq |a|$. ■
 7. If $a \mid b$ and $a \mid c$, then $a \mid (bx + cy)$ for any $x, y \in \mathbb{Z}$. *Proof:* There exist integers r, s such that $b = ar$ and $c = as$. Given integers x, y , $bx + cy = arx + asy = a(rx + sy)$. So $a \mid (bx + cy)$ for all $x, y \in \mathbb{Z}$. ■
- Let $a, b \in \mathbb{Z}$, $|a| + |b| \neq 0$. The greatest common divisor of a and b , denoted $\gcd(a, b)$, is the positive integer d satisfying: $d \mid a$, $d \mid b$, and if $c \mid a$ and $c \mid b$, then $c \leq d$.

- **Given $a, b \in \mathbb{Z}$, $|a| + |b| \neq 0$, there exist $x, y \in \mathbb{Z}$ such that $\gcd(a, b) = ax + by$.** *Proof:* Consider the set $S = \{au + bv : u, v \in \mathbb{Z}, au + bv > 0\}$. S is nonempty as $|a| = (au + b \times 0) \in S$, where $u = \pm 1$. By the well ordering principle, S must contain a least element d . By the definition of S , there exist integers x, y such that $d = ax + by$. Using the division algorithm, we obtain $q, r \in \mathbb{Z}$ such that $a = qd + r$, $0 \leq r < d$. Then $r = a - qd = a - q(ax + by) = a(1 - qx) + b(-qy)$. If $r > 0$, then $r \in S$, but this contradicts d being the least element of S . Thus $r = 0$, and $d \mid a$. By the same reasoning, $d \mid b$. Let c be a positive common divisor of a and b . Then $c \mid (ax + by) = d$, and so $c = |c| \leq |d| = d$. Thus $d = \gcd(a, b)$. ■
- **If $a, b \in \mathbb{Z}$, then the set $T = \{ax + by : x, y \in \mathbb{Z}\}$ is the set of all multiples of $d = \gcd(a, b)$.** *Proof:* Since $d \mid a$ and $d \mid b$, $d \mid (ax + by)$ for all $x, y \in \mathbb{Z}$. Conversely, there exist $x_0, y_0 \in \mathbb{Z}$ such that $d = ax_0 + by_0$. Given $n \in \mathbb{Z}$, $nd = anx_0 + bny_0 \in T$. Thus T is the set of all multiples of d . ■
- Let $a, b \in \mathbb{Z}$, $|a| + |b| \neq 0$. a and b are said to be relatively prime or coprime if $\gcd(a, b) = 1$.
-

3 Euclidean Algorithm

- **Euclidean Algorithm:** Let $a, b \in \mathbb{Z}$. Since $\gcd(|a|, |b|) = \gcd(a, b)$, we can assume that $a \geq b > 0$. Applying the division algorithm, we get $a = q_1b + r_1$, $0 \leq r_1 < b$. If $r_1 = 0$, then $b \mid a$ and $\gcd(a, b) = b$. Otherwise, apply the division algorithm again on b and r_1 to get $b = q_2r_1 + r_2$, $0 \leq r_2 < r_1$. If $r_2 = 0$, we are done. Otherwise divide r_1 by r_2 and so on. The last nonzero remainder obtained in this way is $\gcd(a, b)$.
- **If $a = qb + r$, then $\gcd(a, b) = \gcd(b, r)$.** *Proof:* If $d = \gcd(a, b)$, then $d \mid (a - qb)$ or $d \mid r$. Thus d is a common divisor of b and r . Suppose c is also a common divisor of b and r . Then $c \mid qb + r = a$, so $c \mid a$ and thus $c \leq d$. Therefore, $\gcd(b, r) = d$. This provides the justification for the Euclidean Algorithm. ■
-

4 The Diophantine Equation $ax + by = c$