

Linear Algebra: Systems of Linear Equations

Arjun Vardhan

†

Created: 11th July 2022

Last updated: 11th July 2022

1 Introduction

- Consider a system of m linear equations with n unknowns: $\sum_{k=1}^n a_{1,k}x_k = b_1, \sum_{k=1}^n a_{2,k}x_k = b_2, \dots, \sum_{k=1}^n a_{m,k}x_k = b_m$. Taking the coefficients from the equations, we can form a matrix $A = \begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} \\ \dots & \dots & \dots & \dots \\ a_{m,1} & a_{m,2} & \dots & a_{m,n} \end{pmatrix}$. Then the linear system can be written in the form of a matrix equation $Ax = b$, where $x = (x_1, x_2, \dots, x_n)^T$ and $b = (b_1, b_2, \dots, b_m)^T$.
- A here is the coefficient matrix of the system. If we join the coefficient matrix to the vector b , we get the augmented matrix $\begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} & b_1 \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} & b_2 \\ \dots & \dots & \dots & \dots & \dots \\ a_{m,1} & a_{m,2} & \dots & a_{m,n} & b_m \end{pmatrix}$, which contains all the information necessary to solve the system.

2 Solution of a Linear System, Echelon Form and Reduced Echelon Form

•

3 Analyzing the Pivots

•

4 Finding A^{-1} by Row Reduction

•

5 Dimension

•

6 Solution Set of a Linear System

•

7 Fundamental Subspaces of a Matrix, Rank

-

8 Representation of a Linear Transformation in Arbitrary Bases

-