

Algebra I: Subgroups

Arjun Vardhan

†

Created: 4th February 2022

Last updated: 27th March 2022

1 Definition

- Let G be a group. $H \subseteq G$ is a subgroup of G if $H \neq \phi$ and if $x, y \in H \implies x^{-1}, xy \in H$. We denote this relation by $H \leq G$, or $H < G$ if the containment is proper.
- Subgroups are just subsets of a group that are also groups themselves with the same operations.
- **Subgroup Criterion:** $H \subseteq G$ is a subgroup if and only if $H \neq \phi$ and for all $x, y \in H$, $xy^{-1} \in H$. *Proof:* If $H \leq G$, then $H \neq \phi$ and $x, y \in H \implies xy^{-1} \in H$. Conversely, suppose that H satisfies the two conditions. Then $x \in H \implies xx^{-1} = e \in H$. And thus $e, x \in H \implies ex^{-1} = x^{-1} \in H$. Suppose $x, y \in H$. Then, $y^{-1} \in H \implies xy \in H$. ■

2 Centralizers, Normalizers, Stabilizers and Kernels

- Let $A \subseteq G$, $A \neq \phi$. Let $C_G(A) = \{g \in G : gag^{-1} = a, \forall a \in A\}$. $C_G(A)$ is called the centralizer of A in G . Since $gag^{-1} = a$ if and only if $ga = ag$, $C_G(A)$ is the set of all elements in G that commute with all elements in A .
- $C_G(A) \leq G$. *Proof:* Let $a \in A$. $ea = ae$ so $e \in C_G(A)$ and thus $C_G(A) \neq \phi$. Suppose $x, y \in C_G(A)$. Then $xax^{-1} = y^{-1}ay = a$ for all $a \in C_G(A) \implies xy^{-1}ayx^{-1} = a \implies xy^{-1} \in C_G(A)$. ■
- The center of G , denoted $Z(G)$ is the set of all elements that commute with all elements of G . So $Z(G) = C_G(G)$. $Z(G) = G$ if and only if G is abelian.
- Let $A \subseteq G$, $A \neq \phi$. Let $gAg^{-1} = \{gag^{-1} : a \in A\}$. The normalizer of A in G , is the set $N_G(A) = \{g \in G : gAg^{-1} = A\}$. If $g \in C_G(A)$, then $gag^{-1} = a$ for all $a \in A$, so $C_G(A) \subseteq N_G(A)$.
-

3 Cyclic Groups and Subgroups

4 Subgroups Generated by Subsets

5 Lattice of Subgroups