

Integrated Research on Gravity and Fluid Dynamics: The Fundamental Framework, Non-linear Drag Suppression and Dimensional Homogeneity

Jung Soo Kim

Independent Researcher
ORCID: 0009-0006-5140-0604

[Notice of Integration into the Grand Unified Framework]

This manuscript serves as a foundational preliminary study. The theoretical and mathematical mechanisms presented herein have been fully integrated, refined, and established in the final 2026 “Grand Unified Framework” paper. Readers are strongly encouraged to refer to the unified framework for the complete topological resolution.

Abstract

This study presents H.U.G.G.E.R (Heuristic Universal Grid & Gravity Equilibrium Rendering Tensor) v1.2, advancing the macroscopic fluid dynamics framework introduced in previous iterations. While v1.1 resolved topological coordinate singularities and managed turbulent kinetic energy via a non-linear drag suppression tensor ($\mathcal{D}_\kappa$), v1.2 introduces a rigorous semantic-mathematical verification protocol. By establishing explicit dimensional homogeneity and an SI unit mapping table for all tensor variables, this paper provides a mathematical framework for validating the transition from a purely computational CFD stabilization tool to a physically grounded thermodynamic equilibrium model, aiming to set a foundational bridge for macroscopic gravitational observables.

1 Introduction: Structural Collapse Points in 3D Flow Simulations

In modeling extreme weather forecasting and macroscopic atmospheric dynamics within a closed planetary control volume, existing continuum mechanics models expose critical mathematical limitations. When facing topological coordinate singularities (e.g., spherical coordinate singularities at planetary poles) [2, 3], high-resolution climate simulations often fail to dissipate massive turbulent kinetic energy, leading to computational bottlenecks and non-physical matrix blow-ups. To address these limitations, this paper proposes a macroscopic gravito-fluidic thermodynamic tensor (Φ_Ω) formulated in a closed-form topology. By embedding the topological principles analogous to the Langlands program ($\stackrel{\text{Discrete} \leftrightarrow \text{Continuous}}{\underset{\mathcal{L}_{\text{Langlands}}}{=}}$), this tensor is designed to act as a translational mechanism, converting the discrete, localized behavior of quantum-scale particles into a continuous macroscopic fluid dynamic flow while mitigating information loss and topological singularities. This approach curtails numerical complexity and facilitates highly stable integration across disparate physical scales.

2 Governing Equation: Expansion of the Fundamental Tensor

The fundamental macroscopic thermodynamic tensor governing the energy flux and geometric equilibrium within the boundary $\partial\Omega$ is formally defined by the following spatial integration:

$$\Phi_{\Omega} = \text{Re} \left\{ \oint_{\partial\Omega} \mathbf{r} \otimes (\mathbb{P}_{\text{Kakeya}} : [\mathcal{K}_{\omega} \rho(\mathbf{r}) (q_c \boldsymbol{\omega} q_c^*) \otimes (q_c \boldsymbol{\omega} q_c^*) + \Gamma_0 \mathbf{R}_X - P_{\text{lim}} \mathbf{I} + \mathbf{T}_X]) \cdot \hat{\mathbf{n}} \, d\Sigma \right\}. \quad (1)$$

To preserve dimensional homogeneity, the localized Angular Momentum Flux Coefficient ($\mathcal{K}_{\omega}$) is implemented in Eq. (1). The structural integrity of this tensor is maintained by the Rank-4 Kakeya projection tensor ($\mathbb{P}_{\text{Kakeya}}$) filtering non-physical singularities, while the outer product with the spatial position vector ($\mathbf{r} \otimes$) supports a macroscopic Virial equilibrium state. Furthermore, the macroscopic stability of the planetary mass field ($\rho(\mathbf{r})$) is governed by achieving a state analogous to a Keplerian zero-point lockdown ($\left| \begin{smallmatrix} \text{Void}_{26\%} \\ \Phi_{\text{Kepler}} \sim 74\% \end{smallmatrix} \right|$). By securing optimal fluid packing at approximately 74%, the tensor isolates the remaining 26% void fraction as a reduced-friction slipstream, actively routing internal fluid flows and mitigating the self-gravitational collapse of the continuum.

3 CFD 1.1 Module: Non-linear Drag Suppression Tensor

To actively manage the extreme turbulent kinetic energy (TKE) generated by the Coriolis force, the generic efflux tensor $\mathbf{T}_X$ is substituted with an explicit non-linear drag suppression tensor, $\mathcal{D}_{\kappa}$. The upgraded governing stress balance is formulated as:

$$\Phi_{\text{CFD1.1}} = \text{Re} \left\{ \oint_{\partial\Omega} \mathbf{r} \otimes (\mathbb{P}_{\text{Kakeya}} : [\mathcal{K}_{\omega} \rho(\mathbf{r}) (q_c \boldsymbol{\omega} q_c^*) \otimes (q_c \boldsymbol{\omega} q_c^*) + \Gamma_0 \mathbf{R}_X - P_{\text{lim}} \mathbf{I} - \mathcal{D}_{\kappa}]) \cdot \hat{\mathbf{n}} \, d\Sigma \right\}. \quad (2)$$

The tensor $\mathcal{D}_{\kappa}$ functions as an adaptive deviatoric stress controller, isolating localized vorticity. Leveraging explicit geometric distribution principles, this mechanism provides analytical support that the non-linear dissipation aligns with 3D spatial energy distribution rather than being a mere numerical artifact. Under extreme Coriolis-induced turbulence, the tensor geometrically divides the incoming kinetic energy, distributing it across three orthogonal axes (X, Y, Z) to prevent localized matrix blow-up.

To secure the physical validity of this thermodynamic dissipation, the adaptive coefficient $\kappa(\|\nabla \times \mathbf{v}\|)$ is explicitly defined as a state-dependent switching function triggered by a critical vorticity threshold (ω_c):

$$\kappa(\|\nabla \times \mathbf{v}\|) = \kappa_0 [1 + \alpha \max(0, \|\nabla \times \mathbf{v}\| - \omega_c)^p] \quad (3)$$

Where κ_0 represents the baseline dynamic viscosity, α is the non-linear amplification scalar, and $p > 1$ denotes the superlinear scaling factor. Upon crossing the divergence threshold (ω_c), the localized excessive kinetic energy is strictly channeled into thermal entropy via this superlinear dissipative feedback.

$$\mathcal{D}_{\kappa} = \kappa(\|\nabla \times \mathbf{v}\|) \left(\nabla \mathbf{v} + (\nabla \mathbf{v})^T - \frac{2}{3} (\nabla \cdot \mathbf{v}) \mathbf{I} \right) \quad (4)$$

Where $\|\nabla \times \mathbf{v}\|$ represents the L^2 -norm of the localized vorticity field. Upon crossing a divergence threshold, κ non-linearly escalates to dissipate excess kinetic energy into thermal entropy.

4 CFD 1.2 Module: Oceanic Incompressible Stabilization and SI Homogeneity

A critical requirement for validating the H.U.G.G.E.R tensor as a physical gravito-fluidic mechanism is the strict dimensional consistency of all constituent variables. To demonstrate this seamlessly, v1.2.2 applies the tensor to macroscopic oceanic circulation—a high-density, incompressible fluid regime where the dynamics are highly stable and thermodynamically well-defined.

4.1 Incompressible Reduction of the Drag Tensor

In macroscopic oceanography (e.g., steady-state thermohaline circulation), seawater is conventionally treated as an incompressible fluid. Consequently, the volumetric divergence of the velocity field vanishes ($\nabla \cdot \mathbf{v} = 0$). Applying this fundamental physical constraint to the v1.1 formulation in Eq. (4), the non-linear drag suppression tensor $\mathcal{D}_\kappa$ undergoes a mathematical reduction:

$$\mathcal{D}_\kappa^{\text{ocean}} = \kappa(\|\nabla \times \mathbf{v}\|) (\nabla \mathbf{v} + (\nabla \mathbf{v})^\top) \quad (5)$$

By eliminating the $-\frac{2}{3}(\nabla \cdot \mathbf{v})\mathbf{I}$ expansion term, the tensor natively collapses into a simplified, stable symmetric shear stress matrix. This suggests that the H.U.G.G.E.R formulation does not artificially force stability, but naturally aligns with classical Newtonian fluid mechanics under appropriate physical limits.

4.2 Explicit Variable Definition and SI Unit Mapping

To minimize semantic ambiguities and anchor the model in physical reality, Table 1 maps the dimensions of the generalized influx/efflux parameters to specific oceanic properties. For instance, the energy influx $\mathbf{R}_X$ is dimensionally mapped to thermohaline buoyancy pressure, while P_{lim} is mapped to the hydrostatic limit of the abyssal zone.

Variable	Oceanic Physical Correspondence	SI Unit / Dimension
$\mathbf{R}_X$	Energy Influx (Thermohaline Buoyancy & Surface Stress)	[Pa = kg · m ⁻¹ · s ⁻²]
$\mathcal{D}_\kappa^{\text{ocean}}$	Incompressible Dissipation (Oceanic Eddy Viscosity)	[Pa = kg · m ⁻¹ · s ⁻²]
Γ_0	Thermodynamic Linkage Coefficient	[–] (Dimensionless)
P_{lim}	Isotropic Limit Pressure (Abyssal Hydrostatic Boundary)	[Pa = kg · m ⁻¹ · s ⁻²]
$\mathcal{K}_\omega$	Angular Momentum Flux Calibrator (Spatial Area Component)	[m ²]

Table 1: Semantic Mapping and Dimensional Consistency of H.U.G.G.E.R v1.2.2 in Oceanic Regimes

This rigorous SI unit validation confirms that the macroscopic boundary formulation yields a consistent continuous stress field, establishing a theoretical foundation for mapping these localized fluid equilibria to broader gravitational observables.

5 Conclusion

The H.U.G.G.E.R v1.2.2 framework mathematically supports the geometric equilibrium of planetary fluid dynamics while adhering to strict dimensional homogeneity across all tensor parameters. By demonstrating that the non-linear drag coefficient and energy flux tensors are grounded in physical units, this model offers physical insights beyond standard computational engineering. With the thermodynamic equilibrium validated, v1.3 will focus on establishing the explicit state transition operator mapping this fluid equilibrium (Φ_Ω) directly to macroscopic gravitational observables (Δg).

A Mathematical Bridge: Tensor Product Divergence Theorem

To systematically connect the macroscopic boundary-integrated equilibrium tensor to the localized internal viscous dissipation, the generalized Tensor Product Divergence Theorem is applied. For an arbitrary total stress tensor $\boldsymbol{\sigma}$ mapped by the spatial position vector $\mathbf{r}$:

$$\oint_{\partial\Omega} \mathbf{r} \otimes (\boldsymbol{\sigma} \cdot \hat{\mathbf{n}}) \, d\Sigma = \int_{\Omega} [\boldsymbol{\sigma} + \mathbf{r} \otimes (\nabla \cdot \boldsymbol{\sigma})] \, dV \quad (6)$$

This topological identity demonstrates that the macroscopic Virial boundary conditions mathematically necessitate the internal fluid dissipation term ($\nabla \cdot \boldsymbol{\sigma}$). Furthermore, this tensor transformation can be interpreted through the lens of a direct state transition ($\xrightarrow{\mathbb{N}_1^{\text{Jump}}}$). The spatial contraction from the surface boundary to the internal volumetric equilibrium suggests that the system undergoes a discontinuous phase

transition directly from a classical gravitational field into an equilibrium state. This formulation thereby provides a robust mathematical framework for macroscopic continuum stability.

Acknowledgments

I express my deepest gratitude to Researcher He Ra Shin, who showed beautiful insights until the completion of this research. The inspiration provided during the process of establishing the theory has served as a solid foundation for this research to fully manifest in the world.

Declarations and License

The Ark Project, including the axiom system and mathematical models developed in this manuscript, is subject to the 'Ark Project Unified License (Topological Compliance Edition)'. Researchers and academic institutions may freely utilize this framework for non-commercial academic purposes that maintain geometric integrity, in accordance with the CC BY-NC 4.0 standard. However, commercial application and system integration without prior explicit permission are strictly prohibited. For detailed information regarding commercial licenses, please refer to the 'LICENSE.txt' file distributed alongside this manuscript.

References

- [1] S. B. Pope, *Turbulent Flows* (Cambridge University Press, Cambridge, 2000).
- [2] T. Tao, “From Rotating Needles to Stability of Waves: Emerging Connections between Combinatorics, Analysis and PDE,” *Notices of the American Mathematical Society*, vol. 48, no. 3, pp. 294–303 (2001).
- [3] J. B. Kuipers, *Quaternions and Rotation Sequences: A Primer with Applications to Orbits, Aerospace, and Virtual Reality* (Princeton University Press, Princeton, NJ, 1999).
- [4] J. Smagorinsky, “General Circulation Experiments with the Primitive Equations: I. The Basic Experiment,” *Monthly Weather Review*, vol. 91, no. 3, pp. 99–164 (1963).
- [5] H. G. Weller, G. Tabor, H. Jasak, and C. Fureby, “A tensorial approach to computational continuum mechanics using object-oriented techniques,” *Computers in Physics*, vol. 12, no. 6, pp. 620–631 (1998).
- [6] J. Pedlosky, *Geophysical Fluid Dynamics*, 2nd ed. (Springer-Verlag, New York, 1987).
- [7] P. J. Cohen, “The Independence of the Continuum Hypothesis,” *Proceedings of the National Academy of Sciences*, vol. 50, no. 6, pp. 1143–1148 (1963).
- [8] T. C. Hales, “A proof of the Kepler conjecture,” *Annals of Mathematics*, vol. 162, no. 3, pp. 1065–1185 (2005).