

# Physics Presentation

## Electromagnetism and Special Relativity

Ahan Nag  
20211084

Batch Number - 3  
17<sup>th</sup> June 2022

### 1 The (easy) problem

Two infinite parallel line charges, with line charge density  $\lambda$  each, are separated by a distance  $d$ . They are constrained to move in opposite directions with velocity  $v$  relative to the ground frame. What is the ratio of the magnetic to the electric force (per unit length) exerted by one charge onto the other?




Figure 1: The problem setup (in  $\mathcal{S}$ )

### Solution

Let us consider the upper line charge. The moving charges constitute a current  $I = \lambda v$ . This current, in accordance with Ampère's law, generates a magnetic field

$$B = \frac{\mu_0 I}{2\pi d} = \frac{\mu_0 \lambda v}{2\pi d}$$

at a distance  $d$  from the wire. By the right-hand rule, this field acts outwards from the plane of the charges, for points below the line charge. This field results in a force on the other line charge. Using

$$\mathbf{F}_B = q(\mathbf{v} \times \mathbf{B})$$

the force due to the magnetic field on a segment of length  $l$  (and charge  $\lambda l$ ) is given by

$$F_B = qv \frac{\mu_0 \lambda v}{2\pi d} = \lambda l v \frac{\mu_0 \lambda v}{2\pi d} = \frac{\mu_0 \lambda^2 v^2 l}{2\pi d}$$

which points downward, away from the top charge.

The electric field generated by the top line charge is given by Gauss's law.

$$E = \frac{\lambda}{2\pi\epsilon_0 d}$$

Using

$$\mathbf{F}_E = q\mathbf{E}$$

the force due to the electric field on a segment of length  $l$  is given by

$$F_E = q \frac{\lambda}{2\pi\epsilon_0 d} = \lambda l \frac{\lambda}{2\pi\epsilon_0 d} = \frac{\lambda^2 l}{2\pi\epsilon_0 d}$$

This force, naturally, also points downward, away from the top charge.

Therefore, the ratio of the magnetic force and electric force is given by

$$\frac{F_B}{F_E} = \frac{\frac{\mu_0 \lambda^2 v^2 l}{2\pi d}}{\frac{\lambda^2 l}{2\pi\epsilon_0 d}} = \epsilon_0 \mu_0 v^2 = \frac{v^2}{c^2}$$

where  $c$  is the speed of light. The equality

$$\frac{1}{c^2} = \epsilon_0 \mu_0$$

arises by decoupling Maxwell's equations and obtaining two wave equations for  $\mathbf{E}$  and  $\mathbf{B}$ , from which the speed of propagation of electromagnetic waves is obtained.

The total force per length  $l$  acting on either line charge is

$$F = F_E + F_B = \left(1 + \frac{v^2}{c^2}\right) F_E = \left(1 + \frac{v^2}{c^2}\right) \frac{\lambda^2 l}{2\pi\epsilon_0 d}$$

as  $F_E$  and  $F_B$  act in the same direction.

## 2 The (not so easy) twist

The ratio

$$\frac{F_B}{F_E} = \frac{v^2}{c^2}$$

is reminiscent of the Lorentz factor

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

in special relativity. It can be shown that even though the magnetic force vanishes in the frame of either line charge, the total force per length  $l$  acting on either charge is invariant and is attributable to the electric field alone.

In the frame of the top charge ( $\bar{\mathcal{S}}$ ), the resultant velocity of the bottom charge is given by the Einstein velocity addition rule.

$$\bar{v} = \frac{v_1 - v'}{1 - \frac{v_1 v'}{c^2}} = \frac{2v}{1 + \frac{v^2}{c^2}}$$




Figure 2: The problem setup in  $\bar{\mathcal{S}}$

Here,  $v_1$  is the velocity of the bottom charge in the ground frame ( $\mathcal{S}$ ) and  $v'$  is the velocity of  $\bar{\mathcal{S}}$  with respect to  $\mathcal{S}$  (here,  $v_1 = v$  and  $v' = -v$ ).

In  $\bar{\mathcal{S}}$ , the bottom charge will appear length-contracted by a factor of

$$\bar{\gamma} = \frac{1}{\sqrt{1 - \frac{\bar{v}^2}{c^2}}}$$

Due to the fact that charge is invariant across inertial frames, the charge density on the bottom charge will undergo a corresponding increase, with

$$\begin{aligned} \bar{\lambda}_b &= \bar{\gamma} \lambda \\ &= \frac{1}{\sqrt{1 - \frac{\bar{v}^2}{c^2}}} \lambda \\ &= \frac{c}{\sqrt{c^2 - \bar{v}^2}} \lambda \\ &= \frac{c}{\sqrt{c^2 - \frac{4v^2}{\left(1 + \frac{v^2}{c^2}\right)^2}}} \lambda \\ &= \frac{c \left(1 + \frac{v^2}{c^2}\right)}{\sqrt{c^2 \left(1 + \frac{v^2}{c^2}\right)^2 - 4v^2}} \lambda \end{aligned}$$

Also, the top charge is not moving in  $\bar{\mathcal{S}}$ , but is moving in  $\mathcal{S}$  with velocity  $v$ . Hence, it will appear length-dilated in  $\bar{\mathcal{S}}$ , and the charge density will decrease to

$$\bar{\lambda}_t = \frac{1}{\gamma} \lambda = \sqrt{1 - \frac{v^2}{c^2}} \lambda$$

Therefore, the force per length  $l$  between the charges in  $\bar{\mathcal{S}}$  is given by

$$\bar{F} = \bar{\lambda}_b l \frac{\bar{\lambda}_t}{2\pi\epsilon_0 d} = \frac{\lambda^2 l}{2\pi\epsilon_0 d} \frac{c \left(1 + \frac{v^2}{c^2}\right) \sqrt{1 - \frac{v^2}{c^2}}}{\sqrt{c^2 \left(1 + \frac{v^2}{c^2}\right)^2 - 4v^2}}$$

The separation  $d$  remains constant as the contraction takes place only in the direction of relative motion. As in  $\mathcal{S}$ , the force acts away from the top charge.

In  $\mathcal{S}$ , therefore, the force is transformed as

$$\begin{aligned} F &= \frac{1}{\gamma} \bar{F} \\ &= \frac{\lambda^2 l}{2\pi\epsilon_0 d} \frac{c \left(1 + \frac{v^2}{c^2}\right) \left(1 - \frac{v^2}{c^2}\right)}{\sqrt{c^2 \left(1 + \frac{v^2}{c^2}\right)^2 - 4v^2}} \\ &= \frac{\lambda^2 l}{2\pi\epsilon_0 d} \frac{c \left(1 + \frac{v^2}{c^2}\right) \left(1 - \frac{v^2}{c^2}\right)}{\sqrt{c^2 \left(1 + \frac{v^4}{c^4} + \frac{2v^2}{c^2}\right) - 4v^2}} \\ &= \frac{\lambda^2 l}{2\pi\epsilon_0 d} \frac{c \left(1 + \frac{v^2}{c^2}\right) \left(1 - \frac{v^2}{c^2}\right)}{c \sqrt{1 + \frac{v^4}{c^4} - \frac{2v^2}{c^2}}} \\ &= \frac{\lambda^2 l}{2\pi\epsilon_0 d} \frac{\left(1 + \frac{v^2}{c^2}\right) \left(1 - \frac{v^2}{c^2}\right)}{\sqrt{\left(1 - \frac{v^2}{c^2}\right)^2}} \\ &= \frac{\lambda^2 l}{2\pi\epsilon_0 d} \left(1 + \frac{v^2}{c^2}\right) \end{aligned}$$

This force is exactly the same as the total force  $F$  obtained by considering magnetic and electric forces separately. Thus, what is perceived as a magnetic force in one frame is actually the consequence of the electric force in another frame, and both electric and magnetic forces are components of a single **electromagnetic force**.

### 3 Appendices

#### Derivation of the Einstein velocity addition rule

The Lorentz transformations between frames  $\mathcal{S}$  and  $\bar{\mathcal{S}}$ , where  $\bar{\mathcal{S}}$  moves at a velocity  $v'$  with respect

to  $\mathcal{S}$ , parallel to the x-axis, are given by

$$\begin{aligned}\bar{x} &= \gamma(x - v't) \\ \bar{y} &= y \\ \bar{z} &= z \\ \bar{t} &= \gamma\left(t - \frac{v'}{c^2}x\right)\end{aligned}$$

Let a particle move a distance  $dx$  in  $\mathcal{S}$  in time  $dt$ . Its velocity  $v_1$  in  $\mathcal{S}$  is given by

$$v_1 = \frac{dx}{dt}$$

In  $\bar{\mathcal{S}}$ , it has moved a distance

$$d\bar{x} = \gamma(dx - v'dt)$$

in a time

$$d\bar{t} = \gamma\left(dt - \frac{v'}{c^2}dx\right)$$

according to the Lorentz transformations. Therefore its velocity in  $\bar{\mathcal{S}}$  is given by

$$\bar{v} = \frac{d\bar{x}}{d\bar{t}} = \frac{\gamma(dx - v'dt)}{\gamma\left(dt - \frac{v'}{c^2}dx\right)} = \frac{\frac{dx}{dt} - v'}{1 - \frac{v'}{c^2}\frac{dx}{dt}} = \frac{v_1 - v'}{1 - \frac{v_1 v'}{c^2}}$$

This is the Einstein velocity addition rule.

### Derivation of the force transformation rule

The transformation equation for forces perpendicular to  $v'$  can be obtained as follows.

$$\bar{F}_\perp = \frac{d\bar{p}_\perp}{d\bar{t}} = \frac{dp_\perp}{\gamma\left(dt - \frac{v'}{c^2}dx\right)} = \frac{\frac{dp_\perp}{dt}}{\gamma\left(1 - \frac{v'}{c^2}\frac{dx}{dt}\right)} = \frac{F_\perp}{\gamma\left(1 - \frac{v'v_\parallel}{c^2}\right)}$$

Here,  $d\bar{p}_\perp$  and  $dp_\perp$  are the momenta in the respective frames. The momenta do not transform as they are perpendicular to the direction of relative motion. Now, if  $v_\parallel$  is zero, then

$$\bar{F}_\perp = \frac{1}{\gamma}F_\perp$$

In the problem,  $v_\parallel$  is zero in  $\bar{\mathcal{S}}$  (as the top charge is stationary) instead of  $\mathcal{S}$ , and hence, the perpendicular forces transform as

$$F = \frac{1}{\gamma}\bar{F}$$

## 4 References

- Griffiths, D. J. (1999). *Introduction to Electrodynamics*, Chapter 12, pp. 495-564
- Feynman, Richard P. (1964). *The Feynman Lectures on Physics, Volume II*, Chapter 13-6