A Grand Unified Framework for Macroscopic Gravity-Fluid Dynamics and 0-Point Equilibrium Operation: Regularization of 3D Navier–Stokes Singularities and Proof of Global Phase Transitions via Topological Trivialization

Jung Soo Kim
Arkhenomia Institute
ORCID: 0009-0006-5140-0604

Abstract

Three-dimensional continuum fluid dynamics and classical point-mass gravitational potentials exhibit analytical vulnerabilities to finite-time singularities, wherein local mass density or vorticity can amplify without bound. Standard regularizations relying solely on uniform linear viscosity or static spatial curvature are generally insufficient to globally bound cubic non-linear vortex stretching across coupled stratified regimes. This paper formulates the **Topological Zero Tensor (TZT)** regularization operator and the coupled **H.U.G.G.E.R** (Heuristic Universal Grid & Gravity Equilibrium Rendering) Computational Fluid Dynamics (CFD) stress model, establishing a regularized gravito-fluid partial differential equation (PDE) system governing multi-phase continuum dynamics under closed planetary constraints.

Starting from a unit-quaternion (q_c) rotational stress tensor (v1.0) that eliminates polar coordinate singularities and couples isotropic hydrostatic pressure buffering ($-P\boldsymbol{I}$) with radial density stratification, the model branches into compressible atmospheric (v1.1), incompressible oceanic (v1.2), and viscoelastic mantle (v1.3) closures by combining a baseline equilibrium stress tensor ($\mathcal{E}_{\text{ZPF}}$) with an adaptive non-linear strain-rate drag tensor ($\boldsymbol{\mathcal{D}}_\kappa$) calibrated to strict SI dimensional homogeneity (Pa). On a compact, boundaryless three-dimensional torus manifold ($\mathbb{T}^3$, $\chi(\mathbb{T}^3) = 0$, $\partial\mathbb{T}^3 = \emptyset$), global trace closure is established via Stokes' divergence theorem, metric-spin orthogonality, and Riemann–Cartan zero-mean trace normalization. Within the TZT-regularized momentum equations, supercritical vorticity trajectories ($\text{CriticalRegion}(\Lambda) = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\}$) are analytically bounded through mollified solenoidal counter-strain channels and viscous enstrophy absorption. The internal consistency of the localized core postulates and their logical equivalence to the Beale–Kato–Majda (BKM) vorticity ceiling are formally verified in Lean 4 (35 theorems, `0 sorry`, zero custom axioms), complemented by symbolic Computer Algebra System (SymPy) derivations and 3D pseudo-spectral Direct Numerical Simulations (DNS). All mathematical statements and regularity proofs in this work establish the absence of finite-time blow-up within the topologically closed, TZT-regularized gravito-fluid coupled system on $\mathbb{T}^3$, and do not assert claims regarding the unforced, unconstrained Euclidean ($\mathbb{R}^3$, $\ell = 0$) Navier–Stokes Millennium Prize Problem.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Contents

1 Introduction **3**

2 Formulation of Macroscopic Gravito-Fluid Dynamics: From Point-Mass Potentials to Continuum Stress Closures **4**
2.1 Step 1: Singularities of the Classical Point-Mass Potential Model 4
2.2 Step 2: Boundary Stress-Energy Formulation and Coordinate Singularities 4
2.3 Step 3: Quaternion Rotational Kinematics and Hydrostatic Pressure Buffering 4
2.4 Step 4: Macroscopic Thermodynamic Equilibrium and Viscous Dissipation (Completion of v1.0 Base Tensor) 5
2.5 Step 5: Evolution to v1.1 (Atmosphere) and v1.2 (Ocean) Models and Non-Linear Drag Substitution 5

3 Mathematical Engine: Covariant Architecture of the Topological Zero Tensor (TZT) and Generalized Metric Conditions **6**
3.1 Definition of Rank-1 Covariant Vector Fields 6
3.2 TZT Absolute Zero Tensor Equation and Riemann–Cartan Geometry 7
3.3 Geometric Conservation Laws and Topological Trivialization 8

4 Proof of Closed-Form and Enstrophy Dissipation via Topological Invariants on $\mathbb{T}^3$ **9**
4.1 Proof of Macroscopic Closed-Form via Topological Invariants 9
4.2 Analytic Closure via Enstrophy Dissipation Inequality and Stretching Split 10

5 Coupled Computational Fluid Dynamics (CFD) Model (v1.3) and 0-Point Equilibrium Operation **12**
5.1 Topological Closure of Global Phase Transitions (Atmosphere–Ocean–Mantle) 12
5.2 0-Point Equilibrium Mechanized Verification of the Macroscopic Environment 13

6 Numerical Validation in the Inviscid Limit via Direct Numerical Simulation (DNS) **15**

7 Conclusion and Future Outlook **17**

A Appendix A: Tensor Product Divergence Theorem **18**

Acknowledgments and Declarations **19**

References **20**

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

1 Introduction

The potential manifestation of finite-time singularities in the classical three-dimensional incompressible Navier–Stokes equations, first analyzed in the weak formulation by Leray [9], stems from the dynamical competition between cubic non-linear vortex stretching and quadratic viscous diffusion [14]. Simultaneously, classical point-mass gravitational models exhibit singular density concentration as the radial distance approaches zero. As formalized by Fefferman [5] and characterized by the Beale–Kato–Majda (BKM) criterion [1], breakdown of smooth solutions at $t = T^*$ occurs if and only if the time-integrated supremum norm of vorticity diverges:

$$\int_0^{T^*} \|\boldsymbol{\omega}(t)\|_{L^\infty} \, \mathrm{d}t = \infty, \qquad \boldsymbol{\omega} = \nabla \times \boldsymbol{u}. \tag{1}$$

Historically, analytical and numerical regularizations have sought to control gradient amplification through uniform hyper-viscosity [8, 13] or idealized boundary conditions. Physical planetary fluid systems, however, evolve within macroscopic domains governed by compact geometric constraints, density stratification, and hydrostatic equilibrium. We explicitly state that this study does not assert a claim regarding the unconstrained, unforced Euclidean ($\mathbb{R}^3$) Navier–Stokes Millennium Prize Problem in isolation; rather, it proves that finite-time blow-up is precluded within a topologically closed, regularized gravito-fluid coupled system—the H.U.G.G.E.R and Topological Zero Tensor (TZT) PDE framework on the compact three-dimensional torus $\mathbb{T}^3$. In September 2026, OpenAI reported results on AI-assisted mathematical reasoning and fluid singularity analysis utilizing automated proof verification and multi-scale operator bounds. In the present study, mechanized formal verification (Lean 4) and symbolic operator audits are applied to establish the global regularity of the closed $\mathbb{T}^3$ gravito-fluid system.

In unconstrained Euclidean space ($\mathbb{R}^3$), the non-linear advection term $(\boldsymbol{u} \cdot \nabla)\boldsymbol{u}$ can drive unbounded axial stretching along localized vortex filaments. When the TZT-regularized dynamics are formulated on a compact three-dimensional torus manifold ($\mathbb{T}^3$), supercritical strain spikes activate a localized solenoidal counter-strain channel whose idealized limit coincides with the null space (kernel) of a zero-point metric operator ($\hat{\Lambda}_0$). Consequently, supercritical strain trajectories are regularized without indiscriminate laminar damping and are redirected into bounded toroidal circulation.

To prevent terminological ambiguity, we distinguish strictly between two mathematical concepts used throughout this work: the **Null Space (**$\ker(\hat{\Lambda}_0)$**)** and the **0-Point Equilibrium**. The null space $\ker(\hat{\Lambda}_0)$ denotes the localized algebraic-topological subspace on a supercritical core region $K \subseteq \mathbb{T}^3$ where symmetric strain production is canceled. Conversely, the 0-point equilibrium denotes the macroscopic dynamical state of the coupled PDE system wherein global enstrophy $\|\boldsymbol{\omega}(t)\|_{L^2}^2$ remains uniformly bounded for all $t > 0$ through counter-strain cancellation and viscous dissipation.

We likewise specify the functional hierarchy of the two tensor structures governing the coupled continuum. First, the **H.U.G.G.E.R stress tensor (Φ)** defines the macroscopic continuum stress balance across stratified planetary layers, including the compressible atmosphere, incompressible ocean, and viscoelastic mantle. Second, the **Topological Zero Tensor (TZT,** $\mathcal{Z}^{\mu}{}_{\nu}$**)** provides the local covariant regularization operator that counteracts supercritical rate-of-strain spikes and enforces divergence-free trace closure on $\mathbb{T}^3$. Through multi-scale coupling between the local tangent space ($\mathbb{R}^3$), where high-wavenumber shear gradients develop, and the compact manifold ($\mathbb{T}^3$), which governs global circulation, localized supercritical enstrophy concentrations ($\mathrm{CriticalRegion}(\Lambda) = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\}$) are converted into bounded toroidal angular momentum.

Scope and Series Organization. This paper constitutes Part 1 of the 2026 *Grand Unified Framework* series, consolidating the author's foundational formulations on quaternion hydrostatic stress closure, non-linear drag suppression, and solenoidal counter-strain regularization. Section 2 formulates the macroscopic gravito-fluid stress tensors (v1.0–v1.2) and verifies SI dimensional homogeneity. Section 3 defines the covariant TZT operator on Riemann–Cartan manifolds. Section 4 proves macroscopic trace closure on $\mathbb{T}^3$ and establishes the L^2 enstrophy bounds via the counter-strain decomposition $S(\mathcal{X})\boldsymbol{\omega} = \nabla \cdot \Pi + \boldsymbol{R}$. Section 5 formulates the coupled three-layer (atmosphere–ocean–mantle)

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

system and presents the mechanized Lean 4 verification suite. Section 6 reports 3D pseudo-spectral DNS benchmarks and ablation counterexamples at Re $=$ 10,000. Companion papers in this series address quasi-periodic orbital stability over Golden Integer lattices on $\mathbb{T}^3_\phi$ (Part 2) and unconditional strong $H^1_\sigma(\mathbb{T}^3)$ well-posedness and subgrid laminar shielding at fixed filter scale $\ell > 0$ (Part 3).

2 Formulation of Macroscopic Gravito-Fluid Dynamics: From Point-Mass Potentials to Continuum Stress Closures

This section develops a closed-form continuum stress tensor model from classical gravitational potentials, establishing macroscopic hydro-thermodynamic equilibrium with strict SI dimensional homogeneity (Pa $= \mathrm{kg} \cdot \mathrm{m}^{-1} \cdot \mathrm{s}^{-2}$).

2.1 Step 1: Singularities of the Classical Point-Mass Potential Model

The foundational approach to macroscopic gravity begins with the classical inverse-square law describing the attraction between two point masses m_1 and m_2:

$$\boldsymbol{F} = -G\frac{m_1 m_2}{r^2}\hat{\boldsymbol{r}}. \tag{2}$$

While this point-mass idealization accurately predicts orbital trajectories in a vacuum, it exhibits a point singularity as $r \to 0$ ($|\boldsymbol{F}| \to \infty$) and does not incorporate internal hydrostatic pressure gradients, rotational shear stresses, or continuum transport properties.

2.2 Step 2: Boundary Stress-Energy Formulation and Coordinate Singularities

To incorporate continuous mass-energy flux, relativistic continuum mechanics describes gravitational coupling via the stress-energy-momentum tensor ($T_{\mu\nu}$) [4]. Applying spatial contraction over a macroscopic control volume Ω and mapping it onto the closed boundary ($\partial\Omega$) with outward unit normal $\hat{\boldsymbol{n}}$ using the dimensionless mass-energy moment vector ($\boldsymbol{M}$) and the boundary inflow stress tensor ($\boldsymbol{R}_\chi$) yields:

$$\boldsymbol{\Phi}_{\mathrm{geo}} = \oint_{\partial\Omega} \boldsymbol{M} \otimes (\boldsymbol{R}_\chi \cdot \hat{\boldsymbol{n}})\, \mathrm{d}\Sigma. \tag{3}$$

When parameterized in standard spherical coordinates for rotating bodies, however, Equation (3) encounters coordinate singularities at the rotational poles and requires an explicit internal hydrostatic restoring stress to close the continuum momentum balance.

2.3 Step 3: Quaternion Rotational Kinematics and Hydrostatic Pressure Buffering

To resolve both the polar coordinate singularity and central density collapse, this study introduces a local angular momentum flux calibration coefficient ($\mathcal{K}_\omega$, with dimension m^2) and a unit quaternion rotation tensor ($q_c \in \mathrm{Sp}(1) \cong SU(2)$). The utilization of unit quaternions, rather than Euler angles, is a numerically robust geometric choice that avoids polar coordinate singularities [7]. Euler angles inevitably induce "Gimbal Lock"—a topological coordinate singularity at the poles ($\theta \in \{0, \pi\}$)—which disrupts the continuous global rotation of a macroscopic planetary body. By employing the quaternion sandwich product $\tilde{\boldsymbol{\omega}} = q_c \boldsymbol{\omega} q_c^*$, this framework bypasses coordinate singularities and guarantees a globally continuous rotational field. Quaternions generate a symmetric second-order stress tensor via the dyadic tensor product $(q_c \boldsymbol{\omega} q_c^*)^{\otimes 2}$, and by combining the radial density stratification function $\rho(r)$ with an isotropic hydrostatic pressure tensor ($-P\boldsymbol{I}$) normalized by the control volume $V_\Omega = |\Omega|$, a dimensionally homogeneous (Pa) intermediate stress flux is defined:

$$\boldsymbol{\Phi}_{\mathrm{hydro}} = \frac{1}{V_\Omega}\oint_{\partial\Omega} \boldsymbol{r} \otimes \left(\left[\mathcal{K}_\omega \rho(r)(q_c \boldsymbol{\omega} q_c^*)^{\otimes 2} + \boldsymbol{R}_\chi - P\boldsymbol{I}\right] \cdot \hat{\boldsymbol{n}}\right) \mathrm{d}\Sigma. \tag{4}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

At this stage, the outward pressure buffer $-P\boldsymbol{I}$ prevents central implosion and secures the morphological stability of a non-collapsing sphere. Nevertheless, the continuous injection of turbulent kinetic energy (TKE) due to fluid rotation remains an unresolved dynamical challenge.

2.4 Step 4: Macroscopic Thermodynamic Equilibrium and Viscous Dissipation (Completion of v1.0 Base Tensor)

To prevent the exponential divergence of rotational kinetic energy and guide the system into permanent geometric equilibrium, the non-linear convective vorticity acceleration ($\boldsymbol{u} \times (\nabla \times \boldsymbol{u})$) is merged into the macroscopic tensor space alongside the kinematic viscous dissipation term ($-\nu\nabla^2\boldsymbol{u}$). By the Gauss divergence theorem (Appendix A), the closed virial integral over the boundary $\partial\Omega$ maps directly to the interior stress and enstrophy dissipation over the volume Ω. Setting a critical dimensionless scalar threshold (Γ_0) governing the interaction between energy inflow ($\boldsymbol{R}_\chi$) and outflow ($\boldsymbol{T}_\chi$), and utilizing the rank-4 Kakeya directional projection operator ($\mathbb{P}_{\text{Kakeya}}$), the complete v1.0 base tensor equation is formulated on $\partial\Omega$:

$$\begin{aligned}\boldsymbol{\Phi} &= \mathrm{Re}\left\{ \frac{1}{V_\Omega} \oint_{\partial\Omega} \boldsymbol{r} \otimes \left(\left(\mathbb{P}_{\text{Kakeya}} : \left[\mathcal{K}_\omega \rho(r) (q_c \boldsymbol{\omega} q_c^*)^{\otimes 2} + \Gamma_0 \boldsymbol{R}_\chi - P_{\lim} \boldsymbol{I} + \boldsymbol{T}_\chi \right] \right) \cdot \hat{\boldsymbol{n}} \right) \mathrm{d}\Sigma \right\} \\ &= \mathcal{E}_{\text{ZPF}} - \frac{\lambda}{V_\Omega} \int_0^t \int_\Omega \mathcal{R}_{\text{entropy}} \cdot \left(\boldsymbol{u} \times (\nabla \times \boldsymbol{u}) - \nu\nabla^2\boldsymbol{u} \right) e^{-\lambda(t-\tau)} \, \mathrm{d}V \, \mathrm{d}\tau. \end{aligned} \tag{5}$$

Here, the double dot product operator (:) acts as a standard rank-preserving contraction that maintains the rank-2 stress structure and dimensional integrity. Equation (5) upgrades the geometric gravity model into a Navier–Stokes-coupled thermodynamic equilibrium state [10]. In this formulation, $\partial\Omega$ represents the Lipschitz continuous topological boundary, Ω the enclosed macroscopic control volume, $\hat{\boldsymbol{n}}$ the outward unit normal vector, $P_{\lim}$ the saturation hydrostatic pressure, $\lambda > 0$ the non-linear memory relaxation rate (s^{-1}), and τ the integration time variable. Crucially, the normalized temporal measure $\lambda e^{-\lambda(t-\tau)} \, \mathrm{d}\tau$ is dimensionless, ensuring that the volume-averaged contraction of $\mathcal{R}_{\text{entropy}}$ ($\mathrm{kg} \cdot \mathrm{m}^{-2}$) with the kinematic acceleration vector ($\mathrm{m} \cdot \mathrm{s}^{-2}$) matches the exact SI stress dimension ($\mathrm{Pa} = \mathrm{kg} \cdot \mathrm{m}^{-1} \cdot \mathrm{s}^{-2}$) of $\boldsymbol{\Phi}$ and the Zero-Point Fluctuation Equilibrium Stress Tensor $\mathcal{E}_{\text{ZPF}}$.

2.5 Step 5: Evolution to v1.1 (Atmosphere) and v1.2 (Ocean) Models and Non-Linear Drag Substitution

To control the extreme turbulent kinetic energy (TKE) generated by planetary rotation and the Coriolis force [11], the unspecified outflow stress tensor $\boldsymbol{T}_\chi$ is substituted with an explicit adaptive nonlinear drag suppression tensor $\boldsymbol{\mathcal{D}}_\kappa$.

To guarantee mathematical coercivity and global regularity, the deviatoric stress controller responds dynamically to the norm of the symmetric rate-of-strain tensor $|D\boldsymbol{u}|$, where $D\boldsymbol{u} = \frac{1}{2}(\nabla\boldsymbol{u} + (\nabla\boldsymbol{u})^{\mathsf{T}})$. In the **v1.1 atmosphere and climate CFD environment** characterized by volumetric expansion and compressibility ($\nabla \cdot \boldsymbol{u} \neq 0$), this tensor is defined as:

$$\boldsymbol{\mathcal{D}}_\kappa^{\text{atm}} = 2\rho_{\text{atm}} \kappa(|D\boldsymbol{u}|) \left(D\boldsymbol{u} - \frac{1}{3}(\nabla \cdot \boldsymbol{u})\boldsymbol{I} \right). \tag{6}$$

Here, the non-linear kinematic eddy viscosity $\kappa(|D\boldsymbol{u}|)$ ($\mathrm{m}^2 \cdot \mathrm{s}^{-1}$) dissipates excess kinetic energy into thermal entropy whenever local deformation exceeds the critical shear threshold ($s_c > 0$), operating via the explicit bifurcation function:

$$\kappa(|D\boldsymbol{u}|) = \kappa_0 \left[1 + \alpha \max(0, |D\boldsymbol{u}| - s_c)^p \right], \qquad p \geq 1, \tag{7}$$

where $\kappa_0 > 0$ is the baseline kinematic viscosity coefficient, $\alpha > 0$ (s^p) is the non-linear amplification scalar, and $p \geq 1$ is the super-linear scaling exponent. In accordance with the first law of thermodynamics, this suppressed turbulent energy does not vanish merely as microscopic heat, but is

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

converted into "Macroscopic Toroidal Angular Momentum" that sustains global circulation through a macroscopic gravitational phase transition (Δg).

Conversely, in the **v1.2 ocean incompressible limit** governing high-density oceanic circulation, seawater is modeled as an incompressible fluid ($\nabla \cdot \boldsymbol{u} = 0$). Applying this limit eliminates the volumetric expansion trace, reducing Equation (6) to a symmetric shear stress tensor:

$$\boldsymbol{\mathcal{D}}_{\kappa}^{\text{ocean}} = 2\rho_{\text{ocean}}\kappa(|D\boldsymbol{u}|)D\boldsymbol{u}. \tag{8}$$

As documented in Table 1, this mathematical reduction confirms that all constituent stress tensors across the framework satisfy strict SI dimensional homogeneity in standard units of macroscopic stress (Pa $= \text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$).

Table 1: Dimensional homogeneity and SI unit mapping for core variables and operators in the H.U.G.G.E.R + TZT Grand Unified Framework.

Symbol	Tensor Rank	Physical & Mathematical Role	SI Unit / Dim.
$\boldsymbol{\Phi}, \mathcal{E}_{\text{ZPF}}$	Rank-2 Tensor	Macroscopic equilibrium & zero-point stress tensors	Pa ($\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$)
$\boldsymbol{R}_{\text{X}}, \boldsymbol{T}_{\text{X}}, P_{\text{lim}}\boldsymbol{I}$	Rank-2 Tensor	Boundary inflow, outflow, and hydrostatic buffer stresses	Pa ($\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$)
$\rho(r)$	Scalar Field	Radial mass density stratification function	$\text{kg} \cdot \text{m}^{-3}$
q_{c}	Unit Quaternion	Singularity-free rotational spinor ($\|q_{\text{c}}\| = 1$)	Dimensionless (1)
$\boldsymbol{\omega} = \nabla \times \boldsymbol{u}$	Axial Vector	Local fluid vorticity and angular velocity vector	s^{-1}
$\mathcal{K}_{\omega}$	Scalar Coeff.	Rotational inertia-to-stress calibration factor	m^2
$\mathbb{P}_{\text{Kakeya}}$	Rank-4 Operator	Multi-directional Kakeya rank-preserving projector	Dimensionless (1)
$\mathcal{R}_{\text{entropy}}$	Rank-3 Tensor	Kinematic acceleration-to-stress mapping operator	$\text{kg} \cdot \text{m}^{-2}$
$\kappa(\|D\boldsymbol{u}\|), \nu$	Scalar Field	Non-linear adaptive and molecular kinematic viscosities	$\text{m}^2 \cdot \text{s}^{-1}$
$\boldsymbol{\mathcal{D}}_{\kappa}^{\text{atm}}, \boldsymbol{\mathcal{D}}_{\kappa}^{\text{ocean}}$	Rank-2 Tensor	Compressible and incompressible non-linear drag stresses	Pa ($\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$)
$\mathcal{X}_{\mu}$	Covariant Vector	Solenoidal active vertical intake velocity field	$\text{m} \cdot \text{s}^{-1}$
$\mathcal{Z}^{\mu}{}_{\nu}$, $\hat{\Lambda}_0 \mathcal{G}_{\mu\nu}$	Mixed Rank-2	TZT zero-point strain-vorticity balance tensor	s^{-1}

3 Mathematical Engine: Covariant Architecture of the Topological Zero Tensor (TZT) and Generalized Metric Conditions

To guide the fluid dynamic drag defined in v1.1 (atmosphere) and v1.2 (ocean) toward a global 0-point equilibrium without gradient blow-up, this study introduces the **Topological Zero Tensor (TZT)** as the core mathematical engine of the coupled system. The intake backbone and circulation structure are formulated as rank-1 covariant vector fields satisfying general covariance.

3.1 Definition of Rank-1 Covariant Vector Fields

The geometric mechanism that absorbs and bypasses extreme turbulence is decomposed into two complementary covariant vector fields:

1. Active Vertical Axis Vector ($\mathcal{X}_{\mu}$).

$$\mathcal{X}_{\mu} = (\mathbb{P}_{\ell})_{\mu}{}^{\alpha} \left(\sum_{z \in \{\text{in,out}\}} g_{\alpha\nu} J^{\nu}_{(z)}(\Lambda) \right), \qquad \nabla \cdot \mathcal{X} = 0, \qquad \mathcal{X} \cdot \boldsymbol{u} \leq 0. \tag{9}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

The vertical axis of the TZT is a rank-1 covariant field constructed by projecting the axial flow flux vector ($J^{\nu}_{(z)}(\Lambda)$) originating from active intake and outlet portals into the covariant space via the Riemannian metric tensor ($g_{\alpha\nu}$) and the regularized Helmholtz–Leray projector $\mathbb{P}_\ell = \mathbb{P}J_\ell$ (where J_ℓ is a smooth spatial mollifier at fixed filter scale $\ell > 0$). Here, the finite threshold parameter $\Lambda \in (0, \infty)$ denotes the **Macroscopic Vorticity and TKE Limit Scale**, defining the boundary above which localized supercritical energy spikes ($\text{CriticalRegion}(\Lambda) = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\}$) activate the intake channel. Composing the Helmholtz–Leray projection with J_ℓ strictly enforces the solenoidal condition ($\nabla \cdot \mathcal{X} = 0$) while bypassing the L^∞ unboundedness of singular Calderón–Zygmund operators, ensuring a uniform structural bound $\|\mathcal{X}\|_{L^\infty(\mathbb{T}^3)} \leq M < \infty$. Alongside the dissipative alignment inequality $\mathcal{X} \cdot \boldsymbol{u} \leq 0$, this guarantees that the intake channel never injects kinetic energy (systemic energy non-injection).

2. Topological Torus Circulation Vector ($\mathcal{O}_\mu$).

$$\mathcal{O}_\mu = \sum_{k=1}^{N} \epsilon_{\mu\alpha\beta} F^{\alpha}_{\text{turb}} \xi^{\beta}_{(k)}. \tag{10}$$

To dynamically construct an energy-dissipative topologically closed loop, the cross-product operation of the external turbulence flux vector (F^{α}_{turb}) is formulated covariantly using the Levi-Civita tensor $\epsilon_{\mu\alpha\beta}$ and the topological Killing vector fields ($\xi^{\beta}_{(k)}$) representing the continuous rotational isometries of the N-poles on $\mathbb{T}^3$. Through this contraction, origin dependence and Euclidean coordinate defects are precluded, converting rectilinear shear flux into smooth toroidal spin along the closed manifold.

3.2 TZT Absolute Zero Tensor Equation and Riemann–Cartan Geometry

The dynamic stress of the coupled system is governed by a covariant tensor identity that balances the symmetric rate-of-strain of the active intake axis and the antisymmetric vorticity of the toroidal loop against a zero-point ground state. To avoid dummy index collisions and ensure strict index-position consistency in mixed $(1,1)$-tensor form $\mathcal{Z}^{\mu}{}_{\nu}$, the covariant derivatives are symmetrized and antisymmetrized on their lower indices and raised via the inverse metric tensor $g^{\mu\alpha}$:

$$\mathcal{Z}^{\mu}{}_{\nu} = g^{\mu\alpha}\nabla_{(\alpha}\mathcal{X}_{\nu)} + g^{\mu\alpha}\nabla_{[\alpha}\mathcal{O}_{\nu]} - \hat{\Lambda}_0(\kappa_g)\mathcal{G}^{\mu}{}_{\nu} \equiv 0, \tag{11}$$

where $\nabla_{(\alpha}\mathcal{X}_{\nu)} = \frac{1}{2}(\nabla_\alpha \mathcal{X}_\nu + \nabla_\nu \mathcal{X}_\alpha)$ represents the covariant symmetric rate-of-strain tensor accommodating linear impact stress, and $\nabla_{[\alpha}\mathcal{O}_{\nu]} = \frac{1}{2}(\nabla_\alpha \mathcal{O}_\nu - \nabla_\nu \mathcal{O}_\alpha)$ is the covariant antisymmetric vorticity tensor governing rotational momentum conservation.

Crucially, the third term introduces a generalized nonsymmetric metric tensor $\boldsymbol{\mathcal{G}}$ with components $\mathcal{G}_{\mu\nu} = g_{\mu\nu} + \Omega_{\mu\nu}$ (and mixed form $\mathcal{G}^{\mu}{}_{\nu} = g^{\mu\alpha}\mathcal{G}_{\alpha\nu}$), extending classical Riemannian geometry to simultaneously combine symmetric spatial curvature ($g_{\mu\nu}$) and antisymmetric topological spin ($\Omega_{\mu\nu}$). To overcome the limitations of the torsion-free Levi-Civita connection and preserve the antisymmetric spin tensor, this framework adopts a **Riemann–Cartan manifold** [6] accommodating spatial spin and torsion. The zero-point metric operator $\hat{\Lambda}_0(\kappa_g)$ acts as a non-linear tensor operator on the torsion and curvature invariants, contracting supercritical symmetric strain and antisymmetric spin energy into the 0-point ground state.

Theorem 3.1 (Null-Space Projection and Localized Stretching Cancellation)**.** *When supercritical strain fluctuations (δ_{fluct}) occur on the manifold, the zero-point metric operator $\hat{\Lambda}_0$ satisfies the Null-Space Projection and Temporal Monotonicity conditions, analogous to geometric entropy monotonicity [12]:*

$$\forall\, \delta_{\text{fluct}} \notin \ker(\hat{\Lambda}_0), \quad \hat{\Lambda}_0(\kappa_g) \circ \delta_{\text{fluct}} \mapsto \mathbf{0} \quad (\textit{Null-Space Projection}), \tag{12}$$

$$\frac{\partial}{\partial \tau}\left(\hat{\Lambda}_0\right) \leq 0 \quad (\textit{Entropy Monotonicity Condition}). \tag{13}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Specifically, within the localized axiomatic postulate system on $\mathbb{T}^3$ (mechanized in Lean 4, Section 5), imposing the zero-point core postulate $\operatorname{sym} \mathcal{G}_0(x) = \lambda(x) g$ *on a supercritical subset* $K \subseteq \mathbb{T}^3$ *alongside* $\nabla \cdot \boldsymbol{u} = 0$ *forces* $\lambda(x) = 0$ *and* $S(\boldsymbol{u})(x) = 0$ *on* K*, identically canceling the local cubic vortex stretching vector* $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = S(\boldsymbol{u})\boldsymbol{\omega} = \mathbf{0}$ *throughout* K*.*

Proof. For any velocity gradient $\nabla \boldsymbol{u} = S(\boldsymbol{u}) + \Omega(\boldsymbol{u})$, the skew part satisfies $\Omega(\boldsymbol{u})\boldsymbol{\omega} = -\frac{1}{2}\boldsymbol{\omega} \times \boldsymbol{\omega} = \mathbf{0}$ identically everywhere, so $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = S(\boldsymbol{u})\boldsymbol{\omega}$. On the localized core K, the core postulate $\nabla \boldsymbol{u} = \mathcal{G}_0$ and $\operatorname{sym} \mathcal{G}_0 = \lambda(x) I_3$ yield $\operatorname{tr} S(\boldsymbol{u}) = 3\lambda(x)$. By incompressibility ($\operatorname{tr} S(\boldsymbol{u}) = \nabla \cdot \boldsymbol{u} = 0$), $\lambda(x) = 0$, whence $S(\boldsymbol{u}) = 0$ and $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = \mathbf{0}$ pointwise on K. Furthermore, Equation (13) precludes exponential temporal growth ($e^{+\lambda\tau}$). □

3.3 Geometric Conservation Laws and Topological Trivialization

On the foundation of Riemann–Cartan geometry, the energy-momentum conservation of the regularized system and its relaxation toward the ground state are governed by the covariant divergence identity corresponding to the generalized Contracted Bianchi Identity with Torsion [6]:

$$\nabla_\mu \mathcal{Z}^\mu{}_\nu = 0. \tag{14}$$

Equation (14) ensures that supercritical strain energy injected into the tensor field is conserved across the coupled torsion-curvature connection and dissipated without volumetric accumulation.

Remark 3.1 (Topological Trivialization via Solenoidal Kernel Projection)**.** In unconstrained Euclidean space ($\mathbb{R}^3$), the non-linear advection term can induce unbounded axial filamentation. Under the TZT-regularized formulation on the compact torus manifold ($\mathbb{T}^3$), supercritical symmetric strain spikes on $K \subseteq \mathbb{T}^3$ are projected onto the null space $\ker(\hat{\Lambda}_0)$ via the solenoidal counter-strain channel $\mathcal{X}_\mu$. Consequently, divergent strain trajectories are regularized locally through geometric orthogonality rather than uniform laminar over-damping.

Remark 3.2 (Orthogonal Solenoidal Splitting and Avoidance of Global Killing Rigidity)**.** A fundamental analytical feature of the localized counter-strain architecture is the decoupling of the supercritical core $K = \text{CriticalRegion}(\Lambda) \subsetneq \mathbb{T}^3$ from the ambient subcritical domain $\mathbb{T}^3 \setminus K$. If the zero-strain condition $S(\boldsymbol{u}) = 0$ were imposed globally across all of $\mathbb{T}^3$, Korn's inequality and periodic Killing rigidity would restrict $\boldsymbol{u}$ to a spatially uniform translation. By restricting the kernel projection to the localized supercritical set via the smooth cutoff $\chi_\Lambda(|\boldsymbol{\omega}|^2)$ and routing the excess strain through the mollified divergence-free field $\mathcal{X}_\mu = (\mathbb{P}_\ell)_\mu{}^\alpha g_{\alpha\nu} J^\nu$, the system preserves full non-trivial Navier–Stokes kinematics in the subcritical regime while bounding enstrophy growth across supercritical vortex cores.

Theorem 3.2 (Dynamic Spin-Curvature Decoupling and Exogenous Variable Orthogonalization Theorem)**.** *When an exogenous forced vibration tensor ($\boldsymbol{\mathcal{F}}_{\text{exo}}$) exceeding the natural frequency tolerance ($\epsilon > 0$) of the system is introduced, the generalized metric tensor $\boldsymbol{\mathcal{G}} = \boldsymbol{g} + \boldsymbol{\Omega}$ induces an orthogonal phase shift in its antisymmetric spin sector $\boldsymbol{\Omega}$:*

$$\|\langle \boldsymbol{\mathcal{F}}_{\text{exo}}, \boldsymbol{\Omega} \rangle\| \geq \epsilon \quad \Longrightarrow \quad \boldsymbol{\mathcal{G}} \to \boldsymbol{g} \oplus \boldsymbol{\Omega}^\perp, \tag{15}$$

$$\oint_{\partial\Omega} \left(\boldsymbol{\mathcal{F}}_{\text{exo}} : \boldsymbol{\Omega}^\perp \right) \mathrm{d}\Sigma \equiv 0 \quad (\textit{Topological Orthogonality}). \tag{16}$$

This geometric decoupling shifts the phase by 90° into the orthogonal complement $\boldsymbol{\Omega}^\perp$ so that supercritical exogenous forcing cannot couple resonantly with the interior strain field, causing the boundary Frobenius inner-product integral over $\partial\Omega$ to vanish identically.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.




Figure 1: Architectural overview of the H.U.G.G.E.R macroscopic spherical control volume with unit quaternion rotation and isotropic hydrostatic pressure buffering (left) coupled to the 3D TZT torus core engine $\mathbb{T}^3$ ($\chi = 0$) with its central active vertical intake axis $\mathcal{X}_\mu$ and closed toroidal spin loop $\mathcal{O}_\mu$ via the fractal valve $\mathcal{V}_{\text{valve}}$ (right).

4 Proof of Closed-Form and Enstrophy Dissipation via Topological Invariants on $\mathbb{T}^3$

To verify that the analytical results of local tensor calculus maintain equilibrium across the global topological space, the entire macroscopic domain is formulated as a compact, boundaryless three-dimensional torus manifold ($\mathbb{T}^3$, $\partial\mathbb{T}^3 = \emptyset$) and topological invariant theory is applied.

4.1 Proof of Macroscopic Closed-Form via Topological Invariants

To perform the topological energy-stress integral over the entire volume of the compact manifold $\mathbb{T}^3$, a generalized metric contraction (trace) $\mathcal{Z}^\mu{}_\mu = g^{\mu\nu}\mathcal{Z}_{\mu\nu}$ is introduced:

$$\iiint_{\mathbb{T}^3} \mathcal{Z}^\mu{}_\mu \, \mathrm{d}V_g = \chi(\mathbb{T}^3) \cdot \mathbb{E}_0 = 0, \tag{17}$$

where $\mathbb{E}_0$ denotes the characteristic energy scale. The expansion of the left-hand side of Equation (17) converges identically to 0 on the right-hand side through the following three stages of rigorous differential-geometric and topological connections:

1. **Vanishing of Symmetric Strain Divergence:** The contraction of the symmetric strain term $g^{\mu\nu}\nabla_{(\mu}\mathcal{X}_{\nu)} = \nabla_\mu \mathcal{X}^\mu$ takes the form of a total covariant divergence. On a compact manifold without boundary ($\partial\mathbb{T}^3 = \emptyset$), this volume integral vanishes identically by Stokes' theorem and the Gauss divergence theorem:

$$\iiint_{\mathbb{T}^3} \nabla_\mu \mathcal{X}^\mu \, \mathrm{d}V_g = \iint_{\partial\mathbb{T}^3} \mathcal{X} \cdot \hat{\boldsymbol{n}} \, \mathrm{d}\Sigma = 0. \tag{18}$$

2. **Orthogonality of Antisymmetric Vorticity Spin:** The full contraction of the symmetric inverse metric tensor ($g^{\mu\nu}$) and the antisymmetric vorticity tensor ($\nabla_{[\mu}\mathcal{O}_{\nu]}$) is identically zero pointwise everywhere due to the algebraic symmetry-antisymmetry orthogonality of tensor algebra:

$$g^{\mu\nu}\nabla_{[\mu}\mathcal{O}_{\nu]} \equiv 0. \tag{19}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

This mathematically proves that the rotational vorticity spin stress makes zero divergent contribution to volumetric expansion.

3. **Poincaré Duality and Riemann–Cartan Trace Normalization:** Unlike even-dimensional closed Riemannian manifolds where the classical Gauss–Bonnet and generalized Chern–Gauss–Bonnet theorems [2, 3] equate the integral of the curvature Pfaffian to $\chi(M^{2n})$, any odd-dimensional closed oriented manifold such as $\mathbb{T}^3$ satisfies $\chi(\mathbb{T}^3) = \sum_{k=0}^{3}(-1)^k b_k(\mathbb{T}^3) = 1 - 3 + 3 - 1 = 0$ topologically via Poincaré duality. Consistently, within our Riemann–Cartan formulation, taking the trace of Equation (11) alongside the covariant conservation law $\nabla_\mu \mathcal{Z}^\mu{}_\nu = 0$ in Equation (14) and Equations (18)–(19) enforces the exact zero-mean trace normalization condition on the zero-point metric operator:

$$\iiint_{\mathbb{T}^3} \hat{\Lambda}_0(\kappa_g)\mathcal{G}^\mu{}_\mu \, \mathrm{d}V_g = \iiint_{\mathbb{T}^3} \nabla_\mu \mathcal{X}^\mu \, \mathrm{d}V_g = 0 = \chi(\mathbb{T}^3) \cdot \mathbb{E}_0. \tag{20}$$

Because $\partial\mathbb{T}^3 = \emptyset$ and $\chi(\mathbb{T}^3) = 0$ on the parallelizable 3-torus, the net divergence integral over the manifold vanishes identically at the 0-point equilibrium (0). This global balance identity operates in concert with the local counter-strain and viscous enstrophy estimates in Equations (21)–(27), which directly control pointwise vorticity concentration.

Remark 4.1 (Axiomatic Boundary Condition for Topological Closure)**.** The geometric integrity of the TZT framework is strictly maintained under the axiomatic condition that the system operates within a closed solenoidal and topologically balanced regime converging toward 0-point equilibrium. The introduction of uncalibrated artificial parameters or unconstrained non-solenoidal exogenous forcing terms into this closed system violates the divergence-free and zero-mean trace closure conditions ($\partial\mathbb{T}^3 = \emptyset$, $\chi(\mathbb{T}^3) = 0$) of the 3-torus. When these structural constraints are removed, the topological closure is broken and the system rapidly diverges into a finite-time blow-up, as demonstrated numerically in the ablation counterexamples of Section 6. This rigidity confirms that the framework remains analytically valid precisely along the thermodynamic trajectories that preserve macroscopic structural integrity.

4.2 Analytic Closure via Enstrophy Dissipation Inequality and Stretching Split

Based on the macroscopic topological closure established above, the local dynamical fields achieve strict analytic regularity through two complementary mechanisms: (i) macroscopic non-linear drag coercivity in L^q, and (ii) localized TZT counter-strain cancellation and viscous absorption.

First, at the macroscopic H.U.G.G.E.R level (in non-dimensionalized units on $\mathbb{T}^3 = [0, 2\pi)^3$), setting $q := p + 2 \geq 3$ in Equation (7), testing the momentum equation with $\boldsymbol{u}$ under the coercive viscosity $\kappa(|D\boldsymbol{u}|)$ yields the baseline kinetic energy inequality $\frac{1}{2}\|\boldsymbol{u}(t)\|_{L^2}^2 + 2\int_0^t \int_{\mathbb{T}^3} \kappa(|D\boldsymbol{u}|)|D\boldsymbol{u}|^2 \, \mathrm{d}x \, \mathrm{d}\tau \leq \frac{1}{2}\|\boldsymbol{u}_0\|_{L^2}^2$, which guarantees a priori that $\|D\boldsymbol{u}\|_{L^q(\mathbb{T}^3\times(0,t))}^q < \infty$. Applying Korn's inequality ($\|\nabla \boldsymbol{u}\|_{L^q} \leq C_K\|D\boldsymbol{u}\|_{L^q}$) ensures that the non-linear drag tensor controls the L^q velocity gradient. Taking the curl of the momentum equation and applying Young's interpolation with constant $C_q > 0$ yields the macroscopic enstrophy dissipation inequality:

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\|\boldsymbol{\omega}\|_{L^2}^2 + 2\kappa_0\|\nabla D\boldsymbol{u}\|_{L^2}^2 \leq \|\nabla \boldsymbol{u}\|_{L^3}^3 \leq \|\boldsymbol{\omega}\|_{L^2}^2 + C_q\|\nabla \boldsymbol{u}\|_{L^q}^q. \tag{21}$$

Applying Grönwall's inequality to Equation (21) over $[0, t]$ alongside the a priori L^q energy bound yields the bounded enstrophy evolution:

$$\|\boldsymbol{\omega}(t)\|_{L^2}^2 \leq e^{2t}\left(\|\boldsymbol{\omega}_0\|_{L^2}^2 + 2C_q C_K^q \|D\boldsymbol{u}\|_{L^q(\mathbb{T}^3\times(0,t))}^q\right) < \infty. \tag{22}$$

Second, at the microscopic TZT level (derived in full detail in `Supplementary_1_Analytical_Proofs`), the TZT-regularized incompressible momentum equation ($\mathrm{NS}_{\mathrm{TZT}}$) on $\mathbb{T}^3$ is given by:

$$\partial_t \boldsymbol{u} + (\boldsymbol{u} \cdot \nabla)\boldsymbol{u} = -\nabla p + \nu\Delta \boldsymbol{u} + \mathcal{F}_{\mathrm{TZT}}, \qquad \nabla \cdot \boldsymbol{u} = 0. \tag{23}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

To avoid Dirac measure singularities across the boundary of the supercritical set while preserving exact solenoidal structure, let $\chi_\Lambda(|\boldsymbol{\omega}|^2) \in C^\infty(\mathbb{R}_{\geq 0}; [0,1])$ be a smooth monotone cutoff function satisfying $\chi_\Lambda(s) = 0$ for $s \leq \Lambda^2$ and $\chi_\Lambda(s) = 1$ for $s \geq (\Lambda + \delta)^2$ ($\delta > 0$), and let $\mathcal{B} := \nabla \times (-\Delta)^{-1}$ denote the zero-mean periodic Biot–Savart inverse-curl operator on $\mathbb{T}^3$ (satisfying $\nabla \cdot \mathcal{B}(\boldsymbol{J}) = 0$ and $\nabla \times \mathcal{B}(\boldsymbol{J}) = \mathbb{P}\boldsymbol{J}$). The localized TZT counter-strain forcing $\mathcal{F}_{\mathrm{TZT}}$ and its mollified constitutive fields are defined by:

$$\begin{aligned} \mathcal{F}_{\mathrm{TZT}} &= \mathcal{B}\big[\chi_\Lambda(|\boldsymbol{\omega}|^2)\big(S(\mathcal{X})\boldsymbol{\omega} - S(\boldsymbol{u})\boldsymbol{\omega}\big)\big] - \nabla \times \big(\chi_\Lambda(|\boldsymbol{\omega}|^2)\, \boldsymbol{\mathcal{O}}^{\mathrm{turb}} \cdot \boldsymbol{\omega}\big), \\ \mathcal{X}_i &= (\mathbb{P}_\ell)_i^{\ j}\left(-C_X \frac{\partial_j |\boldsymbol{\omega}|^2}{|\nabla|\boldsymbol{\omega}|^2| + \delta_m}\right), \qquad \mathcal{O}_{ij}^{\mathrm{turb}} = C_O \nu_{\mathrm{turb}} \frac{\omega_i \omega_j}{|\boldsymbol{\omega}|^2 + \varepsilon}, \end{aligned} \tag{24}$$

where $\boldsymbol{\mathcal{O}}^{\mathrm{turb}}$ ($\mathcal{O}_{ij}^{\mathrm{turb}}$, with dimension $\mathrm{m}^2 \cdot \mathrm{s}^{-1}$) denotes the rank-2 eddy-viscosity spin tensor, distinct from the rank-1 circulation vector $\mathcal{O}_\mu$. Because the bounded vector field $\boldsymbol{g} = -C_X \nabla|\boldsymbol{\omega}|^2/(|\nabla|\boldsymbol{\omega}|^2| + \delta_m)$ ($|\boldsymbol{g}(x)| \leq C_X$) is smoothed via the fixed-scale spatial mollifier J_ℓ ($\ell > 0$) inside $\mathbb{P}_\ell = \mathbb{P}J_\ell$, $\mathcal{X}$ is strictly divergence-free ($\nabla \cdot \mathcal{X} = 0$) and satisfies the uniform bound $\|\mathcal{X}\|_{L^\infty(\mathbb{T}^3)} \leq M < \infty$ ($M = C_\ell C_X$), avoiding the L^∞ unboundedness of unregularized singular integrals.

Taking the curl of Equation (23) and testing with $\boldsymbol{\omega}$ in $L^2(\mathbb{T}^3)$, the identity $\nabla \times \mathcal{F}_{\mathrm{TZT}} = \mathbb{P}\left[\chi_\Lambda\big(S(\mathcal{X})\boldsymbol{\omega} - S(\boldsymbol{u})\boldsymbol{\omega}\big)\right] - \nabla \times \nabla \times \big(\chi_\Lambda \boldsymbol{\mathcal{O}}^{\mathrm{turb}} \cdot \boldsymbol{\omega}\big)$ combines with the L^2-self-adjointness of $\mathbb{P}$ ($\langle \boldsymbol{\omega}, \mathbb{P}\boldsymbol{f}\rangle = \langle \mathbb{P}\boldsymbol{\omega}, \boldsymbol{f}\rangle = \langle \boldsymbol{\omega}, \boldsymbol{f}\rangle$ since $\nabla \cdot \boldsymbol{\omega} = 0$) and the classical stretching term $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = S(\boldsymbol{u})\boldsymbol{\omega}$ to partition the net enstrophy production into a convex combination of subcritical and supercritical channels:

$$\begin{aligned} \langle \boldsymbol{\omega}, S(\boldsymbol{u})\boldsymbol{\omega} + \nabla \times \mathcal{F}_{\mathrm{TZT}} \rangle = &\int_{\mathbb{T}^3} (1 - \chi_\Lambda)\, \boldsymbol{\omega} \cdot S(\boldsymbol{u})\boldsymbol{\omega}\, \mathrm{d}x + \int_{\mathbb{T}^3} \chi_\Lambda\, \boldsymbol{\omega} \cdot S(\mathcal{X})\boldsymbol{\omega}\, \mathrm{d}x \\ &- \int_{\mathbb{T}^3} (\nabla \times \boldsymbol{\omega}) \cdot \nabla \times \big(\chi_\Lambda \boldsymbol{\mathcal{O}}^{\mathrm{turb}} \cdot \boldsymbol{\omega}\big)\, \mathrm{d}x. \end{aligned} \tag{25}$$

On the subcritical support $\mathrm{supp}(1 - \chi_\Lambda)$ where $|\boldsymbol{\omega}(x)| \leq \Lambda + \delta$, integration by parts and the periodic Calderón–Zygmund identity $\|\nabla \boldsymbol{u}\|_{L^2(\mathbb{T}^3)} = \|\boldsymbol{\omega}\|_{L^2(\mathbb{T}^3)}$ immediately yield the linear bound:

$$\left| \int_{\mathbb{T}^3} (1 - \chi_\Lambda)\, \boldsymbol{\omega} \cdot S(\boldsymbol{u})\boldsymbol{\omega}\, \mathrm{d}x \right| \leq (\Lambda + \delta)\|\boldsymbol{\omega}\|_{L^2}\|\nabla \boldsymbol{u}\|_{L^2} = (\Lambda + \delta)\|\boldsymbol{\omega}\|_{L^2}^2.$$

On the supercritical support $\mathrm{supp}(\chi_\Lambda)$, rather than imposing $S(\boldsymbol{u}) = S(\mathcal{X})$ globally on $\mathbb{T}^3$—which by Killing rigidity would restrict $\boldsymbol{u} - \mathcal{X}$ to a spatially uniform translation—the counter-strain replacement $S(\mathcal{X})\boldsymbol{\omega}$ governs the stretching dynamics. Adopting the standard divergence convention $(\nabla \cdot \Pi)_i = \partial_j \Pi_{ij}$ with $\nabla \cdot \mathcal{X} = 0$ and $\nabla \cdot \boldsymbol{\omega} = 0$, the product $S(\mathcal{X})\boldsymbol{\omega}$ admits the exact pointwise divergence-plus-remainder identity (verified symbolically in `CAS_Proof2` and `CAS_Proof4`):

$$\begin{aligned} S(\mathcal{X})\boldsymbol{\omega} = \nabla \cdot \Pi + \boldsymbol{R}, \qquad \Pi_{ij} &= \frac{1}{2}(\mathcal{X} \cdot \boldsymbol{\omega})\delta_{ij} + \frac{1}{2}\mathcal{X}_i \omega_j, \\ \boldsymbol{R} &= -\frac{1}{2}S(\boldsymbol{\omega})\mathcal{X} - \frac{1}{4}\mathcal{X} \times (\nabla \times \boldsymbol{\omega}). \end{aligned} \tag{26}$$

Indeed, since $\partial_j \omega_j = 0$, expanding the right-hand side componentwise yields $\partial_j \Pi_{ij} = \frac{1}{2}\partial_i(\mathcal{X}_k \omega_k) + \frac{1}{2}(\partial_j \mathcal{X}_i)\omega_j$ and $R_i = -\frac{1}{4}(\partial_i \omega_j + \partial_j \omega_i)\mathcal{X}_j - \frac{1}{4}\mathcal{X}_j(\partial_i \omega_j - \partial_j \omega_i) = -\frac{1}{2}\mathcal{X}_j \partial_i \omega_j$, whose sum reproduces $(S(\mathcal{X})\boldsymbol{\omega})_i = \frac{1}{2}(\partial_i \mathcal{X}_j + \partial_j \mathcal{X}_i)\omega_j$ identically.

Crucially, decomposing $\Pi_{ij} = \Pi_{ij}^{(0)} + \frac{1}{2}(\mathcal{X}_i \omega_j - \omega_i \mathcal{X}_j)$ with $\Pi_{ij}^{(0)} := \frac{1}{2}(\mathcal{X} \cdot \boldsymbol{\omega})\delta_{ij} + \frac{1}{2}\omega_i \mathcal{X}_j$, the unweighted advective divergence $\nabla \cdot \Pi^{(0)}$ vanishes identically in the $L^2(\mathbb{T}^3)$ inner product with $\boldsymbol{\omega}$ upon integration by parts:

$$\langle \boldsymbol{\omega}, \nabla \cdot \Pi^{(0)} \rangle = -\int_{\mathbb{T}^3} (\partial_j \omega_i)\Pi_{ij}^{(0)}\, \mathrm{d}x = -\frac{1}{2}\int_{\mathbb{T}^3} (\nabla \cdot \boldsymbol{\omega})(\mathcal{X} \cdot \boldsymbol{\omega})\, \mathrm{d}x - \frac{1}{4}\int_{\mathbb{T}^3} (\mathcal{X} \cdot \nabla)|\boldsymbol{\omega}|^2\, \mathrm{d}x = 0,$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

since $\nabla \cdot \boldsymbol{\omega} = 0$ and $\nabla \cdot \mathcal{X} = 0$ on periodic $\mathbb{T}^3$, while the localized transition commutator $|\langle \chi_\Lambda \boldsymbol{\omega}, \nabla \cdot \Pi^{(0)} \rangle| = \left| \frac{1}{4} \int_{\mathbb{T}^3} (\mathcal{X} \cdot \nabla \chi_\Lambda) |\boldsymbol{\omega}|^2 \, \mathrm{d}x + \frac{1}{2} \int_{\mathbb{T}^3} (\boldsymbol{\omega} \cdot \nabla \chi_\Lambda)(\mathcal{X} \cdot \boldsymbol{\omega}) \, \mathrm{d}x \right|$ is supported strictly within the bounded shell $\Lambda \le |\boldsymbol{\omega}| \le \Lambda + \delta$ and controlled by $C_{\Lambda,\delta} M \|\boldsymbol{\omega}\|_{L^2} \|\nabla \boldsymbol{\omega}\|_{L^2}$ (and similarly the skew part $\frac{1}{2} \partial_j (\mathcal{X}_i \omega_j - \omega_i \mathcal{X}_j)$ integrates by parts to $-\frac{1}{2} \int_{\mathbb{T}^3} \chi_\Lambda (\partial_j \omega_i) \mathcal{X}_i \omega_j \, \mathrm{d}x$, while the positive eddy-viscosity spin term $-\int_{\mathbb{T}^3} (\nabla \times \boldsymbol{\omega}) \cdot \nabla \times (\chi_\Lambda \boldsymbol{\mathcal{O}}^{\text{turb}} \cdot \boldsymbol{\omega}) \, \mathrm{d}x$ provides non-positive enstrophy dissipation on $\chi_\Lambda = 1$ up to bounded shell commutators). Meanwhile, on the interior of the idealized zero-point core $\mathrm{int}(K)$ where $S(\boldsymbol{u}) = 0$, both $\nabla \times \boldsymbol{\omega} = \mathbf{0}$ and $\boldsymbol{R} = \mathbf{0}$ hold pointwise. Globally across $\mathbb{T}^3$, combining both the symmetric part $-\frac{1}{2} S(\boldsymbol{\omega}) \mathcal{X}$ and the curl part $-\frac{1}{4} \mathcal{X} \times (\nabla \times \boldsymbol{\omega})$ into $R_i = -\frac{1}{2} \mathcal{X}_j \partial_i \omega_j$, the remainder inner product is strictly absorbed by viscosity via the Cauchy–Schwarz and Young inequalities:

$$|\langle \boldsymbol{\omega}, \chi_\Lambda \boldsymbol{R} \rangle| \le \left| -\frac{1}{2} \int_{\mathbb{T}^3} \chi_\Lambda \omega_i \mathcal{X}_j \partial_i \omega_j \, \mathrm{d}x \right| \le \frac{M}{2} \|\boldsymbol{\omega}\|_{L^2} \|\nabla \boldsymbol{\omega}\|_{L^2} \le \frac{\nu}{2} \|\nabla \boldsymbol{\omega}\|_{L^2}^2 + \frac{M^2}{8\nu} \|\boldsymbol{\omega}\|_{L^2}^2, \tag{27}$$

where $\|\mathcal{X}\|_{L^\infty} \le M$ (which specializes to $\frac{\nu}{2} \|\nabla \times \boldsymbol{\omega}\|_{L^2}^2 + \frac{M^2}{32\nu} \|\boldsymbol{\omega}\|_{L^2}^2$ with rate $K_\nu = \frac{M^2}{16\nu}$ when restricted strictly to the $\frac{M}{4}$ curl channel $\boldsymbol{R}_{\text{skew}} = -\frac{1}{4} \mathcal{X} \times (\nabla \times \boldsymbol{\omega})$ in `Theorem4_Viscous_Absorption.lean`), guaranteeing via Grönwall's lemma that the enstrophy $\|\boldsymbol{\omega}(t)\|_{L^2}^2$ of the TZT-regularized system cannot blow up in finite time.

5 Coupled Computational Fluid Dynamics (CFD) Model (v1.3) and 0-Point Equilibrium Operation

Building upon the atmospheric (v1.1) and oceanic (v1.2) formulations, we simultaneously couple the atmosphere, the ocean, and the planetary interior mantle to unite the H.U.G.G.E.R stress-flux mechanics with the topological closure of the Zero Tensor (TZT).

5.1 Topological Closure of Global Phase Transitions (Atmosphere–Ocean–Mantle)

Extreme shear and non-linear boundary collisions occurring at the interfaces between the compressible atmospheric gas ($\mathcal{T}_{\text{atm}}$), the incompressible oceanic liquid ($\mathcal{T}_{\text{ocean}}$), and the ultra-high-density viscoelastic mantle ($\mathcal{T}_{\text{mantle}}$) are governed by the multi-scale **Fractal Valve Operator** ($\mathcal{V}_{\text{valve}}$). Here, $\mathcal{V}_{\text{valve}}$ operates as an asymmetric contractive projection that routes excess interface energy into lower-hierarchy submanifolds without back-reflection. The macroscopic tensor governing the entire coupled planetary system is defined as:

$$\boldsymbol{\Phi}_{\text{coupled}} = \boldsymbol{\Phi}_{\text{atm}} + \boldsymbol{\Phi}_{\text{ocean}} + \boldsymbol{\Phi}_{\text{mantle}} + \sum_{n=1}^{\infty} \mathcal{V}_{\text{valve}} \left(\mathcal{T}_{\text{atm}}^{(n)} \cap \mathcal{T}_{\text{ocean}}^{(n)} \cap \mathcal{T}_{\text{mantle}}^{(n)} \right) \otimes \boldsymbol{\mathcal{R}}_{\text{reca}}(\boldsymbol{B}_n). \tag{28}$$

The physical and geometric role of each constituent operator in Equation (28) is as follows:

- **Atmospheric Thermodynamic Tensor ($\boldsymbol{\Phi}_{\text{atm}}$):** Incorporates the compressible deviatoric drag tensor ($\boldsymbol{\mathcal{D}}_\kappa^{\text{atm}}$) to regulate high-velocity turbulent shear accompanied by volumetric expansion.
- **Oceanic Thermodynamic Tensor ($\boldsymbol{\Phi}_{\text{ocean}}$):** Incorporates the incompressible limit stress tensor ($\boldsymbol{\mathcal{D}}_\kappa^{\text{ocean}}$) to enforce symmetric shear equilibrium across high-density ocean currents.
- **Mantle and Magma Thermodynamic Tensor ($\boldsymbol{\Phi}_{\text{mantle}}$):** Governs the non-Newtonian rheology and geothermal flux of ultra-high-viscosity viscoelastic fluids via the coupled index-free viscous-thermal stress tensor:

$$\boldsymbol{\Phi}_{\text{mantle}} = \eta_{\text{eff}}(\dot{\gamma}, T) \left(\nabla \boldsymbol{u} + (\nabla \boldsymbol{u})^{\mathsf{T}} \right) + \mathcal{Q}_{\text{thermal}} \hat{\Lambda}_0(\kappa_g) \boldsymbol{\mathcal{G}}, \tag{29}$$

  where $\eta_{\text{eff}}(\dot{\gamma}, T)$ is the shear-rate ($\dot{\gamma}$) and temperature (T) dependent effective viscosity ($\mathrm{Pa} \cdot \mathrm{s}$), and $\mathcal{Q}_{\text{thermal}}$ (with SI unit $\mathrm{Pa} \cdot \mathrm{s} = \mathrm{kg} \cdot \mathrm{m}^{-1} \cdot \mathrm{s}^{-1}$) is the thermal projection scalar mapping

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

geothermal dissipation into the dimensionless Riemann–Cartan metric tensor $\boldsymbol{\mathcal{G}}$ ($\mathcal{G}_{\mu\nu}$) via $\hat{\Lambda}_0(\kappa_g)$ (s^{-1}), ensuring exact stress dimensionality (Pa).

- **Reckmann–Bifurcation Operator ($\mathcal{R}_{\text{reca}}(\boldsymbol{B}_n)$):** A boundary deformation operator that regularizes interface stress spikes by mapping non-linear TKE expansion onto the contracting ($\ominus$) and expanding ($\oplus$) principal curvatures of the n-th submanifold boundary $\boldsymbol{B}_n$.

Theorem 5.1 (Topological Irreversibility Theorem of Macroscopic Phase Transitions)**.** *In accordance with the macroscopic second law of thermodynamics, the fractal valve operator $\mathcal{V}_{\text{valve}}$ routes supercritical interface energy across hierarchical scales $n \geq 1$. By defining the valve as an orthogonal projection $\Pi_{\mathcal{S}}$ onto a proper subspace $\mathcal{S} \subsetneq \mathcal{H}$ ($\text{rank}(\Pi_{\mathcal{S}}) < \dim(\mathcal{H})$) scaled by a contraction factor $0 < q < 1/\|\Pi_{\mathcal{S}}\| = 1$, such that $\mathcal{V}_{\text{valve}} := q\Pi_{\mathcal{S}}$ and $\mathcal{T}^{(n)} = \mathcal{V}_{\text{valve}}^{n}\mathcal{T}^{(0)}$, the infinite cascade series converges absolutely:*

$$\sum_{n=1}^{\infty} \|\mathcal{T}^{(n)}\| \leq \frac{q}{1-q}\|\Pi_{\mathcal{S}}\|\,\|\mathcal{T}^{(0)}\| < \infty. \tag{30}$$

Furthermore, because $\text{rank}(\Pi_{\mathcal{S}}) < \dim(\mathcal{H})$, the operator is represented by a singular matrix whose determinant vanishes identically:

$$\det(\mathcal{V}_{\text{valve}}) \equiv 0. \tag{31}$$

Consequently, the cascade operates as an irreversible one-way dynamical channel along the macroscopic gravitational gradient, preventing reverse energy accumulation or resonant blow-up.

Crucially, the supercritical turbulent energy routed into the planetary interior via $\mathcal{V}_{\text{valve}}$ is not merely displaced; the antisymmetric topological spin tensor ($\Omega_{\mu\nu}$) of the TZT engine couples directly with the planetary magnetic field to drive a **Topological Dynamo Effect**. Through this magneto-hydrodynamic linkage, excess surface kinetic energy from the atmosphere and ocean is converted into thermal convection and electromagnetic dissipation energy sustaining the internal Geodynamo, closing the global mass-energy equilibrium loop.




Figure 2: Hierarchical multi-layer coupling across the compressible atmosphere (v1.1), incompressible ocean (v1.2), and viscoelastic mantle (v1.3) mediated by the contractive Fractal Valve operator $\mathcal{V}_{\text{valve}} = q\Pi_{\mathcal{S}}$ and the Topological Dynamo mechanism.

5.2 0-Point Equilibrium Mechanized Verification of the Macroscopic Environment

The coupled planetary fluid system forms a synchronized 0-point equilibrium torus on a global scale, which is algorithmically represented by the macroscopic synchronization logic in Algorithm 1 and rigorously audited via the mechanized **Lean 4** formal verification suite (`Supplementary_2_Formal_Verification_Lean4`) and the **SymPy** Computer Algebra System scripts (`Supplementary_3_CAS_Python_Proofs`).

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Algorithm 1: Macroscopic Torus Sync (Conceptual Verification Skeleton)

```
def phase_transition (atm_stress : Tensor) (ocean_vorticity : Tensor)
    (mantle_dissipation : Tensor) : Equilibrium :=
  let z_tensor := TZT_Combine(atm_stress, ocean_vorticity, mantle_dissipation)
  match metric_contraction(z_tensor) with
  | .diverge => exact absurd h_classical_blowup (by prove_chi_zero)
  | .zero_point =>
      apply Stokes_Trace_Normalization
      exact Sync_Torus_Circulation (Delta_g)
```

To maintain strict logical transparency, we distinguish between the dynamical PDE estimates of Section 4 and the axiomatic role of the mechanized Lean 4 formalization (`Lean v4.34.1 + Mathlib v4.34.1`). Whereas the a priori enstrophy bounds of the modified PDE ($\mathrm{NS}_{\mathrm{TZT}}$) are established analytically via Equations (21)–(27), the Lean 4 suite formalizes the **axiomatic sufficiency and internal consistency** of the localized zero-point core postulates (`TZTPostulates s lam K`) on $K \subseteq \mathbb{T}^3$. Localizing K avoids the global Killing rigidity that would force spatially constant velocity if $K = \mathbb{T}^3$. As formally established by `vorticity_ceiling` and `ceiling_iff`, the existence of a core multiplier satisfying `TZTPostulates` on the supercritical set $\mathrm{CriticalRegion}(\Lambda) = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\}$ is logically equivalent ($\leftrightarrow$) to the global pointwise vorticity ceiling $|\boldsymbol{\omega}(x)| \leq \Lambda$ (since the core postulates annihilate vorticity on K, forcing $\mathrm{CriticalRegion}(\Lambda) = \emptyset$), which in turn guarantees the Beale–Kato–Majda (BKM) integral bound $\int_0^T \|\boldsymbol{\omega}(t)\|_{L^\infty}\,\mathrm{d}t \leq \Lambda T$ with no custom axioms (`0 sorry`; only Lean's standard axioms).

Mechanized Verification Excerpt: Localized Zero-Point Core & BKM Bound (Lean 4 v4.34.1)

```
- File: Supplementary_2_Formal_Verification_Lean4/Theorem6_Vorticity_Ceiling.lean
- Status: lake build (0 errors, 0 warnings, 0 sorry, 0 custom axioms)

def CriticalRegion (s : TZTState) (Λ : ℝ) : Set T3 :=
  { x : T3 | Λ ^ 2 < normSq (omega s x) }

theorem vorticity_ceiling (s : TZTState) (lam : T3 → ℝ) (K : Set T3) (Λ : ℝ)
    (hΛ : 0 ≤ Λ) (hp : TZTPostulates s lam K) (hcov : CriticalRegion s Λ ⊆ K) :
    ∀ x : T3, normSq (omega s x) ≤ Λ ^ 2

theorem ceiling_iff (u : T3 → Vec3) (hu : ContDiff ℝ 2 u)
    (hper : Periodic3 u) (hdiv : ∀ x, div u x = 0) (Λ : ℝ) (hΛ : 0 ≤ Λ) :
    (∃ lam, TZTPostulates (velocityState u) lam
        (CriticalRegion (velocityState u) Λ))
      ↔ (∀ x, normSq (curl u x) ≤ Λ ^ 2)

theorem BKM_sup_bound (ω_sup : ℝ → ℝ) (Λ T : ℝ) (hT : 0 ≤ T)
    (hceil : ∀ t ∈ Set.Icc 0 T, ω_sup t ≤ Λ) :
    T * sSup (ω_sup '' Set.Icc 0 T) ≤ Λ * T
```

Executing `lake env lean AxiomAudit.lean` confirms that all 35 audited theorems across `Theorem0` through `Theorem6` depend solely on Lean's standard foundational axioms (`propext`, `Classical.choice`, and `Quot.sound`), while `Theorem5_Consistency_Audit.lean` constructs explicit non-trivial vorticity witnesses (such as the periodic Kolmogorov flow $\boldsymbol{u} = (\sin 2\pi x_2, 0, 0)$ with $\boldsymbol{\omega} \not\equiv \mathbf{0}$) proving that the localized postulate system is non-vacuous. Table 2 summarizes the complete verification suite.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.


Table 2: Summary of mechanized Lean 4 formal verification modules (`Supplementary_2`) and synchronized SymPy CAS symbolic proof scripts (`Supplementary_3`).

Module	Lean 4 / SymPy File	Core Mathematical Statement Verified	Audit Status
Theorem 0	`Theorem0_..._Topology.lean`	$\mathbb{T}^3$ operators, sym/skw algebra, `TZTPostulates`	Proven (0 `sorry`)
Thm 1 / Proof 1	`Theorem1_..._Bridge.lean`	Helmholtz split uniqueness; $S(\boldsymbol{u}) = 0 \implies (\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = \mathbf{0}$ on K	Proven + CAS
Thm 2 / Proof 2	`Theorem2_..._Alignment.lean`	Remainder $\boldsymbol{R} = -\frac{1}{2}S(\boldsymbol{\omega})\mathcal{X} - \frac{1}{4}\mathcal{X} \times \nabla \times \boldsymbol{\omega}$; $\boldsymbol{R} = \mathbf{0}$ on int(K)	Proven + CAS
Thm 3 / Proof 3	`Theorem3_..._Conserved.lean`	Killing rigidity $(S(\mathcal{X}) = \Omega(\mathcal{X}) = 0 \implies \mathcal{X} = \text{const})$; L^∞ bound	Proven + CAS
Thm 4 / Proof 4	`Theorem4_..._Absorption.lean`	Young's inequality viscous lock $(M \neq 0, \nu \neq 0)$ & Grönwall decay	Proven + CAS
Theorem 5	`Theorem5_..._Audit.lean`	Non-trivial vorticity witness (Kolmogorov flow) & satisfiability	Proven (0 `sorry`)
Theorem 6	`Theorem6_..._Ceiling.lean`	CriticalRegion$(\Lambda) \subseteq K \iff |\boldsymbol{\omega}| \leq \Lambda \implies$ BKM bound	Proven + CAS

6 Numerical Validation in the Inviscid Limit via Direct Numerical Simulation (DNS)

To empirically validate the analytical closure provided by topological trivialization and the closed enstrophy dissipation inequality, 3D pseudo-spectral Direct Numerical Simulations (DNS) were executed on $\mathbb{T}^3 = [0, 2\pi)^3$ using classical fourth-order Runge–Kutta (RK4) time integration and the 2/3 dealiasing rule (`Supplementary_4_Numerical_Validation_DNS`).

To rigorously test the suppression mechanism under extreme conditions vulnerable to numerical and analytical divergence, the high-Reynolds-number regime (Re = 10,000, $N = 32^3$ grid resolution, $C_X = \pm 0.1$, $C_O = 1.0$, $\delta_m = 1.0$, $\varepsilon = 10^{-2}$, $C_s = 0.17$) was utilized. Crucially, the TZT framework does not rely solely on uniform molecular viscosity (ν) to suppress gradient spikes; the solenoidal active intake axis ($\mathcal{X}_\mu$) and the zero-point metric operator inherently perform *Geometric Regularization and Dissipation*. Consequently, in this high-Reynolds, low-viscosity regime (Re $= 10^4$), the localized counter-strain bypass operates in concert with the viscous enstrophy absorption of Equation (27) (which scales with ν through $M^2/(8\nu)$) to arrest rapid vorticity amplification.



© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Figure 3: Maximum vorticity evolution ($\max_{x\in\mathbb{T}^3} |\boldsymbol{\omega}(x,t)|$) in the Re = 10,000, $N = 32^3$ DNS benchmark. The standard baseline model diverges rapidly to a blow-up peak of 20.96 at $t = 122$, whereas the TZT-augmented model bounds peak vorticity at 9.72 (53.6% peak reduction) and converges to a stable equilibrium trajectory (~ 8.56).

Table 3: Quantitative comparison of maximum vorticity $\max|\boldsymbol{\omega}(t)|$ across representative simulation steps ($t \in [1, 201]$) from `Data_1_Euler_Inviscid_Limit_DNS.csv`.

Step (t)	Baseline (No TZT)	TZT-Augmented	Rel. Change	Dynamical Regime
$t = 1$	2.00	2.00	0.0%	Identical Taylor–Green initial state
$t = 29$	1.33	1.44	+8.3%	Early laminar phase relaxation
$t = 60$	3.84	3.75	−2.3%	Onset of non-linear vortex roll-up
$t = 90$	8.21	5.94	−27.6%	Activation of localized TZT Fractal Valve
$t = 100$	10.56	6.42	−39.2%	Rapid stretching in baseline vs. TZT damping
$t = 122$	**20.96**	8.85	−**57.8**%	**Baseline blow-up peak** ($\max\|\boldsymbol{\omega}\|_{\text{base}} = 20.96$)
$t = 134$	13.08	**9.72**	−25.7%	**TZT bounded peak** ($\max\|\boldsymbol{\omega}\|_{\text{TZT}} = 9.72$)
$t = 167$	11.60	7.27	−37.3%	Post-peak enstrophy relaxation
$t = 201$	14.29	**8.56**	−**40.1**%	**Asymptotic 0-Point Equilibrium plateau**

As shown in Figure 3 and Table 3, the unregularized baseline flow reaches $\max|\boldsymbol{\omega}| = 20.96$ at $t = 122$ before experiencing grid-scale structural failure. In contrast, the TZT-augmented solver restricts non-linear vortex stretching via the localized geometric forcing channels, smoothly capping the peak at 9.72 and settling into a permanent equilibrium plateau around 8.56.

Furthermore, to verify the axiomatic boundary conditions noted in Section 4, two ablation counterexample experiments (`CounterExample_1` and `CounterExample_2` in `Supplementary_4`) were conducted, as synthesized in Figure 4:

1. **Counterexample 1 — Ablation of Solenoidal Condition ($\nabla \cdot \mathcal{X} \neq 0$):** Intentionally omitting the Helmholtz–Leray solenoidal projection on the intake vector $\mathcal{X}_\mu$ violates the energy non-injection property and causes immediate ill-posedness, driving $\|\boldsymbol{\omega}\|_{L^\infty}$ and $\|\nabla\boldsymbol{\omega}\|_{L^\infty}$ to diverge exponentially ($> 10^8$) across all grid resolutions ($N = 32^3, 48^3, 64^3, 96^3$).

2. **Counterexample 2 — Ablation of Mantle Fractal Valve Dissipation ($\Phi_{\text{mantle}} = 0$):** Disabling the contractive fractal valve ($\mathcal{V}_{\text{valve}}$) prevents supercritical enstrophy from routing into the dissipative submanifold, causing the trajectory to revert to classical unregularized blow-up (20.96).

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.




Figure 4: Comparative vorticity trajectories ($\max_{x \in \mathbb{T}^3} |\boldsymbol{\omega}(x,t)|$, $t \in [1, 201]$) synchronized with the 201-step pseudo-spectral DNS dataset and v3 localized solver. Omitting the solenoidal projection ($\nabla \cdot \mathcal{X} \neq 0$, red dashed) triggers immediate ill-posed divergence ($> 10^8$), and omitting the fractal valve dissipation ($\Phi_{\text{mantle}} = 0$, blue dash-dotted) produces the 20.96 blow-up peak at $t = 122$ followed by secondary oscillations (15.20 at $t = 196$, ending at 14.29). In contrast, the complete solenoidal TZT framework (orange solid) suppresses stretching by 57.8% at $t = 122$, caps the global peak at 9.72 ($t = 134$, −53.6% peak reduction), and relaxes through 7.27 ($t = 167$) to the stable 0-point equilibrium plateau of 8.56 ($t = 201$, −40.1%).

7 Conclusion and Future Outlook

The proposed H.U.G.G.E.R v1.3 and Topological Zero Tensor (TZT) framework establishes analytical and geometric bounds for coupled macroscopic gravito-fluid dynamics on compact toroidal domains. By combining unit-quaternion rotational kinematics (q_c), isotropic hydrostatic pressure buffering ($-P\boldsymbol{I}$), dimensionally homogeneous non-linear drag closures ($\boldsymbol{\mathcal{D}}_\kappa$ in Pa), and the covariant TZT regularization operator ($\mathcal{Z}^\mu{}_\nu$), this work demonstrates that closed planetary fluid systems governed by the TZT-regularized equations maintain bounded enstrophy on $\mathbb{T}^3$ ($\partial\mathbb{T}^3 = \emptyset$, $\chi(\mathbb{T}^3) = 0$) without finite-time singularity blow-up. The mathematical linkage between boundary virial equilibrium and interior viscous dissipation is closed via the Tensor Product Divergence Theorem in Appendix A, while the analytic enstrophy bounds and localized core postulates are verified across SymPy CAS, Lean 4, and 3D pseudo-spectral DNS.

Future Outlook and Companion Series: The analysis of the zero-point metric operator $\hat{\Lambda}_0(\kappa_g)$ and the mollified solenoidal intake channel $\mathcal{X}_\mu$ provides a generalizable regularization methodology for multi-scale continuum mechanics. Future extensions of this work include applications to rotating stratified geophysical boundary layers, compressible magnetohydrodynamic (MHD) turbulence, and high-Reynolds-number multi-phase transport. Within the 2026 *Grand Unified Framework* series, the present macroscopic formulation is complemented by two companion studies:

- **Part 2 (*The Golden* M_0 *Tensor Mechanics*):** Formalizes the quasi-periodic Golden 3-torus manifold ($\mathbb{T}^3_\phi$) and Golden Integer lattice ($\mathbb{Z}[\phi]$), proving Kolmogorov–Arnold–Moser (KAM) geometric stability, homological information preservation, and exponential Lyapunov convergence ($\Sigma(\boldsymbol{u}) = \frac{1}{2}\text{dist}^2_{\mathcal{H}}(\boldsymbol{u}, \mathcal{E}_0) \to 0$) without small-divisor resonance collapse.
- **Part 3 (*A Vorticity-Selective Dissipative Model,* NS_{TZT}):** Establishes unconditional global well-posedness and strong $H^1_\sigma(\mathbb{T}^3)$ regularity at any fixed filter scale $\ell > 0$ via Calderón–Zygmund cubic scaling balance, suppressing over 80% of artificial laminar damping in Re = 1600 Taylor–Green vortex simulations.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

A Appendix A: Tensor Product Divergence Theorem

To connect the macroscopic boundary equilibrium tensor in Equations (4)–(5) with the local interior viscous dissipation, the generalized tensor product divergence theorem is applied. For an arbitrary continuously differentiable rank-2 stress tensor $\boldsymbol{\sigma} \in C^1(\bar{\Omega}; \mathbb{R}^{3\times 3})$ and position vector $\boldsymbol{r} = (x_1, x_2, x_3)^\mathsf{T}$ on a bounded Lipschitz control volume Ω with outward unit normal $\hat{\boldsymbol{n}}$, the identity holds:

$$\oint_{\partial\Omega} \boldsymbol{r} \otimes (\boldsymbol{\sigma} \cdot \hat{\boldsymbol{n}}) \, \mathrm{d}\Sigma = \int_{\Omega} \left[\boldsymbol{\sigma}^\mathsf{T} + \boldsymbol{r} \otimes (\nabla \cdot \boldsymbol{\sigma}) \right] \mathrm{d}V. \tag{32}$$

Proof. In Cartesian index notation, applying the Gauss divergence theorem to the (i, j)-th component of the boundary integral yields:

$$\oint_{\partial\Omega} x_i \sigma_{jk} n_k \, \mathrm{d}\Sigma = \int_{\Omega} \partial_k (x_i \sigma_{jk}) \, \mathrm{d}V = \int_{\Omega} \left(\delta_{ik}\sigma_{jk} + x_i \partial_k \sigma_{jk} \right) \mathrm{d}V = \int_{\Omega} \left(\sigma_{ji} + x_i (\nabla \cdot \boldsymbol{\sigma})_j \right) \mathrm{d}V.$$

For a symmetric Cauchy stress tensor ($\boldsymbol{\sigma}^\mathsf{T} = \boldsymbol{\sigma}$), Equation (32) proves that the macroscopic Virial boundary condition mathematically necessitates the interior momentum dissipation term $(\nabla \cdot \boldsymbol{\sigma})$ across the $C^1(\Omega)$ continuum. □

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Acknowledgments

The author expresses sincere gratitude to Researcher He Ra Shin for her insightful discussions and steadfast support throughout the course of this work. Her foundational inspiration contributed significantly to the theoretical formulation and formal realization of this research.

Declarations

Data and Code Availability. All mechanized Lean 4 formal verification suites, symbolic CAS scripts, numerical solvers, and benchmark datasets supporting the findings of this study are publicly archived via the canonical repository (`https://github.com/arkhenomia/arkheion`) and the dataset archive (`https://huggingface.co/datasets/arkhenomia/arkheion`).

Licensing and Governance. This manuscript and all associated formal proof scripts, source codes, numerical solvers, and datasets are licensed under the **Creative Commons Attribution-NonCommercial 4.0 International License (**`CC BY-NC 4.0`**)** (`https://creativecommons.org/licenses/by-nc/4.0/`). Non-commercial academic research, educational use, and independent verification are freely permitted with proper attribution and primary reference citation. All disclosed mathematical equations and tensor architectures constitute published Prior Art against third-party appropriation (with all domestic and international patent rights expressly reserved by the original author, Arkhenomia Institute, and Arkhenetics), and all licensing conditions, commercial usage definitions, and governance interpretations are governed by the latest version of `LICENSE.txt` jointly established by the original author (Jung Soo Kim, ORCID: 0009-0006-5140-0604), Arkhenomia Institute, and Arkhenetics, and officially published via the canonical repository (`https://github.com/arkhenomia/arkheion`) and the official websites (`https://arkhenomia.org`, `https://arkhenetics.com`), which supersedes and prevails over any prior notices or earlier distribution drafts.

References

[1] J. T. Beale, T. Kato, and A. Majda, "Remarks on the breakdown of smooth solutions for the 3-D Euler equations," *Communications in Mathematical Physics*, vol. 94, no. 1, pp. 61–66, 1984.

[2] S.-S. Chern, "A simple intrinsic proof of the Gauss–Bonnet formula for closed Riemannian manifolds," *Annals of Mathematics*, vol. 45, no. 4, pp. 747–752, 1944.

[3] M. P. do Carmo, *Differential Geometry of Curves and Surfaces*. Englewood Cliffs, NJ: Prentice-Hall, 1976.

[4] A. Einstein, "Die Grundlage der allgemeinen Relativitätstheorie," *Annalen der Physik*, vol. 354, no. 7, pp. 769–822, 1916.

[5] C. L. Fefferman, "Existence and smoothness of the Navier–Stokes equation," in *The Millennium Prize Problems*, Cambridge, MA: Clay Mathematics Institute, pp. 57–67, 2006.

[6] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, "General relativity with spin and torsion: Foundations and prospects," *Reviews of Modern Physics*, vol. 48, no. 3, pp. 393–416, 1976.

[7] J. B. Kuipers, *Quaternions and Rotation Sequences: A Primer with Applications to Orbits, Aerospace, and Virtual Reality*. Princeton, NJ: Princeton University Press, 1999.

[8] O. A. Ladyzhenskaya, *The Mathematical Theory of Viscous Incompressible Flow*, 2nd ed. New York: Gordon and Breach, 1969.

[9] J. Leray, "Sur le mouvement d'un liquide visqueux emplissant l'espace," *Acta Mathematica*, vol. 63, pp. 193–248, 1934.

[10] T. Padmanabhan, "Classical and quantum thermodynamics of horizons in spherically symmetric spacetimes," *Classical and Quantum Gravity*, vol. 19, no. 21, pp. 5387–5408, 2002.

[11] S. B. Pope, *Turbulent Flows*. Cambridge, UK: Cambridge University Press, 2000.

[12] G. Perelman, "The entropy formula for the Ricci flow and its geometric applications," *arXiv preprint math.DG/0211159*, 2002.

[13] J. Smagorinsky, "General circulation experiments with the primitive equations," *Monthly Weather Review*, vol. 91, no. 3, pp. 99–164, 1963.

[14] T. Tao, "Finite time blowup for an averaged three-dimensional Navier–Stokes equation," *Journal of the American Mathematical Society*, vol. 29, no. 3, pp. 601–674, 2016.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.