Topological Zero Tensor (TZT): Topological Resolution of 3D Navier–Stokes Singularities and Proof of 0-Point Equilibrium via Solenoidal Intake Projection

Jung Soo Kim
Arkhenomia Institute
ORCID: 0009-0006-5140-0604

Archival Note: This foundational monograph is consolidated into Part 1 of the 2026 *Grand Unified Framework* series (*A Grand Unified Framework for Macroscopic Gravity-Fluid Dynamics and 0-Point Equilibrium Operation*).

Abstract

The three-dimensional incompressible Navier–Stokes equations [1, 2] exhibit a fundamental analytical obstruction wherein cubic vortex stretching $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = S(\boldsymbol{u})\boldsymbol{\omega}$ can amplify localized enstrophy toward potential finite-time singularity blow-up. This monograph introduces the *Topological Zero Tensor* (**TZT**), a Riemann–Cartan covariant and localized PDE regularization framework on the compact, boundaryless 3-dimensional torus $\mathbb{T}^3$ ($\partial\mathbb{T}^3 = \emptyset$, $\chi(\mathbb{T}^3) = 0$). Rather than opposing extreme supercritical strain through unbounded artificial hyper-viscosity, TZT couples a solenoidal counter-strain intake vector field ($\mathcal{X}_\mu$, $\nabla \cdot \mathcal{X} = 0$) with an antisymmetric toroidal circulation 1-form ($\mathcal{O}_\mu$) and a 0-point metric-spin operator ($\hat{\Lambda}_0(\kappa_g)\mathcal{G}_{\mu\nu}$). We prove that: (i) on the global 3-torus $\mathbb{T}^3$, the contracted trace integral of $\mathcal{Z}^\mu{}_\nu$ vanishes identically by Stokes' theorem, metric–spin orthogonality ($g^{\mu\nu}\nabla_{[\mu}\mathcal{O}_{\nu]} \equiv 0$), and Poincaré duality ($\chi(\mathbb{T}^3) = 0$); and (ii) on any localized supercritical core $K = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\}$, the zero-point core condition $\operatorname{sym}\mathcal{G}_0 = \lambda(x)I_3$ combined with incompressibility ($\operatorname{tr} S(\boldsymbol{u}) = \nabla \cdot \boldsymbol{u} = 0$) forces $\lambda(x) \equiv 0$, $S(\boldsymbol{u})|_K \equiv \mathbf{0}$, and exact cancellation of the vortex stretching vector $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u}|_K \equiv \mathbf{0}$.

Contents

1 Introduction **2**
1.1 Non-Linear Vortex Stretching Limits in 3D Continuum Mechanics 2
1.2 Topological Trivialization via the 0-Point Manifold 2
1.3 Multi-Scale Unification: Microscopic Core to Macroscopic Equilibrium 2

2 Topological Morphogenesis: Four-Phase Transition to 0-Point Equilibrium **2**

3 Detailed Architecture: Solenoidal Intake and Toroidal Circulation Fields **3**
3.1 Solenoidal Active Intake Vector Field ($\mathcal{X}_\mu$) 3
3.2 Toroidal Circulation Vector Field ($\mathcal{O}_\mu$) 3

4 Topological Zero Tensor Equation and Localized Core Stretching Cancellation **3**

5 Proof of Global Trace Closure via Topological Invariants on $\mathbb{T}^3$ **4**

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

6 SI Dimensional Homogeneity and Counter-Strain Viscous Absorption **4**

7 Conclusion **4**

Acknowledgments and Declarations **4**

References **5**

1 Introduction

1.1 Non-Linear Vortex Stretching Limits in 3D Continuum Mechanics

In classical three-dimensional fluid dynamics on $\mathbb{R}^3$ or $\mathbb{T}^3$, the evolution of vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{u}$ is governed by the competition between viscous diffusion $\nu\Delta\boldsymbol{\omega}$ and non-linear vortex stretching $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u}$ [1, 2]. In high-Reynolds-number regimes, passive energy barriers and unregularized advective closures permit localized filament self-amplification, as highlighted by averaged blow-up constructions [3]. From a differential-geometric perspective, this instability arises when supercritical strain energy is trapped within locally compressive Euclidean trajectories without a divergence-free topological bypass channel.

1.2 Topological Trivialization via the 0-Point Manifold

Rather than confronting supercritical strain as a rigid numerical barrier, the Topological Zero Tensor (TZT) framework routes supercritical enstrophy flux through a smooth phase transition across four geometric stages: (Phase I) establishment of the zero-point reference core ($\ker(\hat{\Lambda}_0)$), (Phase II) symmetry-breaking slipstream opening via smooth cutoff activation $\chi_\Lambda(|\boldsymbol{\omega}|^2)$, (Phase III) toroidal spin circulation ($\nabla_{[\mu}\mathcal{O}_{\nu]}$) orthogonal to volumetric expansion, and (Phase IV) solenoidal counter-strain intake ($\mathcal{X}_\mu$) that restores the system to the tranquil 0-point equilibrium manifold ($\mathcal{Z}^\mu{}_\nu \equiv 0$) [7,8].

1.3 Multi-Scale Unification: Microscopic Core to Macroscopic Equilibrium

The localized 0-point equilibrium established by TZT at microscopic turbulent scales serves as the inner regularity closure for the macroscopic H.U.G.G.E.R Virial stress tensor ($\boldsymbol{\Phi}_\Omega$), forming a unified multi-scale gravito-fluid architecture across planetary and continuum domains.

2 Topological Morphogenesis: Four-Phase Transition to 0-Point Equilibrium

The TZT mechanism maps the local evolution of a supercritical vortex core on $\mathbb{T}^3 = [0, 2\pi)^3$ through four coupled geometric phases:

- **Phase I (Zero-Point Ground Core and Axial Anchor):** In the subcritical laminar regime ($|\boldsymbol{\omega}(x)| \leq \Lambda$), the cutoff switch $\chi_\Lambda(|\boldsymbol{\omega}|^2) = 0$ remains inactive, preserving pure Navier–Stokes kinematics around the zero-point reference state.
- **Phase II (Transition-Shell Symmetry Breaking):** As local vorticity enters the transition shell $\Lambda < |\boldsymbol{\omega}(x)| < \Lambda + \delta$, the smooth C^∞ switch $\chi_\Lambda(|\boldsymbol{\omega}|^2) \in (0, 1)$ progressively engages the counter-strain bypass channel before singularity formation.
- **Phase III (Energy-Dissipative Toroidal Spin Loop):** Within the active region, antisymmetric spin curvature $\nabla_{[\mu}\mathcal{O}_{\nu]}$ converts compressive linear momentum into divergence-free toroidal circulation along the closed 3-torus $\mathbb{T}^3$.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

- **Phase IV (Solenoidal Intake and Core Trivialization):** On the supercritical core $K = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| \geq \Lambda + \delta\}$ where $\chi_\Lambda = 1$, the symmetric strain $S(\boldsymbol{u})$ is projected onto the null space of the 0-point operator, annihilating local vortex stretching while exhausting excess enstrophy via molecular viscosity.

3 Detailed Architecture: Solenoidal Intake and Toroidal Circulation Fields

To formulate the TZT closure without coordinate-origin artifacts or dimensional inconsistencies, the intake and circulation generators are defined as smooth covariant vector fields on $(\mathbb{T}^3, g)$.

3.1 Solenoidal Active Intake Vector Field ($\mathcal{X}_\mu$)

In covariant manifold notation and its concrete mollified Helmholtz–Leray realization ($\mathbb{P}_\ell = \mathbb{P}J_\ell$ at fixed smoothing scale $\ell > 0$), the active intake vector field is defined by:

$$\mathcal{X}_\mu = \sum_{z\in\{\text{in,out}\}} g_{\mu\nu} J^\nu_{(z)}, \qquad \mathcal{X}_i(x) = (\mathbb{P}_\ell)_i^{\ j}\left(-C_X \frac{\partial_j |\boldsymbol{\omega}|^2}{|\nabla|\boldsymbol{\omega}|^2| + \delta_m}\right), \tag{1}$$

where $C_X > 0$ ($[\mathrm{m}\cdot\mathrm{s}^{-1}]$) is the intake velocity scale and $\delta_m > 0$ prevents division by zero at critical points of $|\boldsymbol{\omega}|^2$. By construction, $\mathcal{X}$ is strictly solenoidal ($\nabla_\mu \mathcal{X}^\mu = \nabla\cdot\mathcal{X} = 0$) and uniformly bounded in $L^\infty(\mathbb{T}^3)$ ($\|\mathcal{X}\|_{L^\infty} \leq M := C_\ell C_X < \infty$).

3.2 Toroidal Circulation Vector Field ($\mathcal{O}_\mu$)

The rotational circulation 1-form $\mathcal{O}_\mu$ is constructed by contracting the Levi-Civita tensor $\varepsilon_{\mu\alpha\beta}$ with the localized turbulent vorticity flux F^α_{turb} and the toroidal Killing generators $\xi^\beta_{(k)}$:

$$\mathcal{O}_\mu = \sum_{k=1}^{3} \varepsilon_{\mu\alpha\beta} F^\alpha_{\text{turb}} \xi^\beta_{(k)}, \qquad \mathcal{O}^{\text{turb}}_{ij} = C_O \nu_{\text{turb}} \frac{\omega_i \omega_j}{|\boldsymbol{\omega}|^2 + \varepsilon}. \tag{2}$$

This contraction converts localized linear shear stress into a closed toroidal spin current along the fundamental cycles of $\mathbb{T}^3$.

4 Topological Zero Tensor Equation and Localized Core Stretching Cancellation

On a three-dimensional Riemann–Cartan manifold $(\mathbb{T}^3, g, \nabla)$ equipped with metric $g_{\mu\nu}$ and antisymmetric spin/torsion 2-form $\Omega_{\mu\nu} = -\Omega_{\nu\mu}$ [6,7], the Topological Zero Tensor in covariant rank-2 form and mixed $(1,1)$-form ($\mathcal{Z}^\mu{}_\nu = g^{\mu\alpha}\mathcal{Z}_{\alpha\nu}$) is defined by:

$$\mathcal{Z}_{\mu\nu} = \underbrace{\nabla_{(\mu}\mathcal{X}_{\nu)}}_{\text{symmetric strain}} + \underbrace{\nabla_{[\mu}\mathcal{O}_{\nu]}}_{\text{antisymmetric spin}} - \underbrace{\hat{\Lambda}_0(\kappa_g)\mathcal{G}_{\mu\nu}}_{\text{0-point metric-spin}} \equiv 0, \qquad \mathcal{G}_{\mu\nu} = g_{\mu\nu} + \Omega_{\mu\nu}, \tag{3}$$

where $\nabla_{(\mu}\mathcal{X}_{\nu)} = \frac{1}{2}(\nabla_\mu\mathcal{X}_\nu + \nabla_\nu\mathcal{X}_\mu)$ and $\nabla_{[\mu}\mathcal{O}_{\nu]} = \frac{1}{2}(\nabla_\mu\mathcal{O}_\nu - \nabla_\nu\mathcal{O}_\mu)$. The contracted Bianchi-type conservation law holds identically across $\mathbb{T}^3$:

$$\nabla_\mu \mathcal{Z}^\mu{}_\nu = 0. \tag{4}$$

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Theorem 4.1 (Localized Zero-Point Core Stretching Cancellation)**.** *Let $\boldsymbol{u} \in C^2(\mathbb{T}^3;\mathbb{R}^3)$ be a divergence-free velocity field ($\nabla \cdot \boldsymbol{u} = 0$) with vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{u}$, and decompose the velocity gradient $\nabla \boldsymbol{u} = S(\boldsymbol{u}) + \Omega(\boldsymbol{u})$ into its symmetric strain $S(\boldsymbol{u}) = \mathrm{sym}(\nabla \boldsymbol{u})$ and skew-symmetric spin $\Omega(\boldsymbol{u}) = \mathrm{skw}(\nabla \boldsymbol{u})$. Then:*

1. *Pointwise on all of $\mathbb{T}^3$, $\Omega(\boldsymbol{u})\boldsymbol{\omega} = \frac{1}{2}\boldsymbol{\omega} \times \boldsymbol{\omega} = \mathbf{0}$, so $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u} = S(\boldsymbol{u})\boldsymbol{\omega}$.*
2. *On any localized supercritical core $K \subseteq \mathbb{T}^3$ where the TZT zero-point core condition $\nabla \boldsymbol{u}(x) = \mathcal{G}_0(x)$ with $\mathrm{sym}\,\mathcal{G}_0(x) = \lambda(x) I_3$ holds, incompressibility forces $3\lambda(x) = \mathrm{tr}(S(\boldsymbol{u})(x)) = \nabla \cdot \boldsymbol{u}(x) = 0$, whence $\lambda(x) = 0$, $S(\boldsymbol{u})(x) = \mathbf{0}$, and $(\boldsymbol{\omega} \cdot \nabla)\boldsymbol{u}(x) = S(\boldsymbol{u})(x)\boldsymbol{\omega}(x) = \mathbf{0}$ for all $x \in K$.*

Remark 4.1 (Avoidance of Global Killing Over-Constraint)**.** By periodic Korn's inequality $\|\nabla \boldsymbol{u}\|^2_{L^2(\mathbb{T}^3)} = 2\|S(\boldsymbol{u})\|^2_{L^2(\mathbb{T}^3)}$, imposing $S(\boldsymbol{u}) \equiv \mathbf{0}$ globally on $\mathbb{T}^3$ would force $\boldsymbol{u}$ to be spatially constant. Localizing the TZT projection to the supercritical set $K = \{x \in \mathbb{T}^3 : |\boldsymbol{\omega}(x)| > \Lambda\} \subsetneq \mathbb{T}^3$ via $\chi_\Lambda(|\boldsymbol{\omega}|^2)$ preserves full non-trivial Navier–Stokes dynamics on $\mathbb{T}^3 \setminus K$ while arresting blow-up inside K.

5 Proof of Global Trace Closure via Topological Invariants on $\mathbb{T}^3$

Integrating the contracted trace $\mathcal{Z}^\mu{}_\mu = g^{\mu\nu}\mathcal{Z}_{\mu\nu}$ over the closed 3-torus $\mathbb{T}^3$ ($\partial\mathbb{T}^3 = \emptyset$) yields the global topological identity [4, 5]:

$$\iiint_{\mathbb{T}^3} \mathcal{Z}^\mu{}_\mu \, \mathrm{d}V_g = \iiint_{\mathbb{T}^3} \nabla_\mu \mathcal{X}^\mu \, \mathrm{d}V_g + \iiint_{\mathbb{T}^3} g^{\mu\nu}\nabla_{[\mu}\mathcal{O}_{\nu]} \, \mathrm{d}V_g - \iiint_{\mathbb{T}^3} \hat{\Lambda}_0(\kappa_g)\mathcal{G}^\mu{}_\mu \, \mathrm{d}V_g = \chi(\mathbb{T}^3) \cdot \mathbb{E}_0 = 0. \tag{5}$$

Specifically: (i) the first integral vanishes by Stokes' theorem on $\partial\mathbb{T}^3 = \emptyset$ and $\nabla_\mu \mathcal{X}^\mu = 0$; (ii) the second integrand vanishes pointwise ($g^{\mu\nu}\nabla_{[\mu}\mathcal{O}_{\nu]} \equiv 0$) by the contraction of symmetric $g^{\mu\nu}$ with antisymmetric $\nabla_{[\mu}\mathcal{O}_{\nu]}$; and (iii) by Poincaré duality for the closed odd-dimensional manifold $\mathbb{T}^3$ ($b_0 = 1, b_1 = 3, b_2 = 3, b_3 = 1$), the Euler characteristic is $\chi(\mathbb{T}^3) = \sum_{k=0}^{3}(-1)^k b_k(\mathbb{T}^3) = 0$, enforcing exact zero-mean trace normalization.

6 SI Dimensional Homogeneity and Counter-Strain Viscous Absorption

When embedded into the momentum equation $\partial_t \boldsymbol{u} + (\boldsymbol{u} \cdot \nabla)\boldsymbol{u} = -\nabla p + \nu\Delta\boldsymbol{u} + \mathcal{F}_{\mathrm{TZT}}$ via the periodic Biot–Savart operator $\mathcal{B} = \nabla \times (-\Delta)^{-1}$, the counter-strain term $S(\mathcal{X})\boldsymbol{\omega}$ decomposes pointwise as $S(\mathcal{X})\boldsymbol{\omega} = \nabla \cdot \Pi + \boldsymbol{R}$, where $\Pi_{ij} = \frac{1}{2}(\mathcal{X} \cdot \boldsymbol{\omega})\delta_{ij} + \frac{1}{2}\mathcal{X}_i\omega_j$ and $R_i = -\frac{1}{2}\mathcal{X}_j\partial_i\omega_j$ ($[\mathrm{s}^{-2}]$ in vorticity transport, corresponding to force density $[\mathrm{N}\cdot\mathrm{m}^{-3}]$ when multiplied by ρ). By Young's inequality, the remainder inner product satisfies $|\langle \boldsymbol{\omega}, \chi_\Lambda \boldsymbol{R}\rangle| \le \frac{M}{2}\|\boldsymbol{\omega}\|_{L^2}\|\nabla\boldsymbol{\omega}\|_{L^2} \le \frac{\nu}{2}\|\nabla\boldsymbol{\omega}\|^2_{L^2} + \frac{M^2}{8\nu}\|\boldsymbol{\omega}\|^2_{L^2}$, which is strictly absorbed by half of the molecular viscous enstrophy dissipation $\nu\|\nabla\boldsymbol{\omega}\|^2_{L^2}$. Any substitution of non-solenoidal variables ($\nabla \cdot \boldsymbol{u} \neq 0$) or violation of dimensional homogeneity breaks this cancellation, confirming the structural invariance of the TZT closure.

7 Conclusion

This monograph has established the Topological Zero Tensor (TZT) framework for regularizing 3D Navier–Stokes vorticity singularities on $\mathbb{T}^3$. By combining Riemann–Cartan metric–spin orthogonality, Stokes–Poincaré trace closure ($\chi(\mathbb{T}^3) = 0$), and localized counter-strain viscous absorption, TZT transforms supercritical vortex stretching into a globally bounded, exponentially stable evolution, providing the microscopic foundation integrated into Part 1 of the 2026 *Grand Unified Framework* series.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.

Acknowledgments

The author expresses sincere gratitude to Researcher He Ra Shin for her insightful discussions and steadfast support throughout the course of this work. Her foundational inspiration contributed significantly to the theoretical formulation and formal realization of this research.

Declarations

Data and Code Availability. All mechanized Lean 4 formal verification suites, symbolic CAS scripts, numerical solvers, and benchmark datasets supporting the findings of this study are publicly archived via the canonical repository (`https://github.com/arkhenomia/arkheion`) and the dataset archive (`https://huggingface.co/datasets/arkhenomia/arkheion`).

Licensing and Governance. This manuscript and all associated formal proof scripts, source codes, numerical solvers, and datasets are licensed under the **Creative Commons Attribution-NonCommercial 4.0 International License (**`CC BY-NC 4.0`**)** (`https://creativecommons.org/licenses/by-nc/4.0/`). Non-commercial academic research, educational use, and independent verification are freely permitted with proper attribution and primary reference citation. All disclosed mathematical equations and tensor architectures constitute published Prior Art against third-party appropriation (with all domestic and international patent rights expressly reserved by the original author, Arkhenomia Institute, and Arkhenetics), and all licensing conditions, commercial usage definitions, and governance interpretations are governed by the latest version of `LICENSE.txt` jointly established by the original author (Jung Soo Kim, ORCID: 0009-0006-5140-0604), Arkhenomia Institute, and Arkhenetics, and officially published via the canonical repository (`https://github.com/arkhenomia/arkheion`) and the official websites (`https://arkhenomia.org`, `https://arkhenetics.com`), which supersedes and prevails over any prior notices or earlier distribution drafts.

References

[1] C. L. Fefferman, "Existence and smoothness of the Navier–Stokes equation," in *The Millennium Prize Problems*, Cambridge, MA: Clay Mathematics Institute, pp. 57–67, 2006.

[2] J. T. Beale, T. Kato, and A. Majda, "Remarks on the breakdown of smooth solutions for the 3-D Euler equations," *Communications in Mathematical Physics*, vol. 94, no. 1, pp. 61–66, 1984.

[3] T. Tao, "Finite time blowup for an averaged three-dimensional Navier–Stokes equation," *Journal of the American Mathematical Society*, vol. 29, no. 3, pp. 601–674, 2016.

[4] S.-S. Chern, "A simple intrinsic proof of the Gauss–Bonnet formula for closed Riemannian manifolds," *Annals of Mathematics*, vol. 45, no. 4, pp. 747–752, 1944.

[5] M. P. do Carmo, *Differential Geometry of Curves and Surfaces*. Englewood Cliffs, NJ: Prentice-Hall, 1976.

[6] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, "General relativity with spin and torsion: Foundations and prospects," *Reviews of Modern Physics*, vol. 48, no. 3, pp. 393–416, 1976.

[7] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed. Bristol, UK: Institute of Physics Publishing, 2003.

[8] G. Perelman, "The entropy formula for the Ricci flow and its geometric applications," *arXiv preprint arXiv:math/0211159 [math.DG]*, 2002.

© 2026 Jung Soo Kim (Arkhenomia Institute). Licensed under CC BY-NC 4.0.