

## Homework 4 – Algebraic Structures

---

Ch 7: 3.1, 3.3, 3.4, 4.7, 4.9, 5.3, 5.6, 5.7, 5.12, 6.4, 6.5

Blake Griffith

Pre-lect: 5.1

**Exercise (7.3.1).** *Prove the Fixed Point Theorem (7.3.2).*

*Proof.* Recall the counting formula

$$|G| = |G_s||O_s|$$

and since the orbits partition the set

$$|S| = |O_1| + \cdots + |O_k|$$

So for some element  $s \in S$ ,  $|O_s|$  must divide  $|G|$ , so it can be  $p$  or 1.  $|O_s|$  must also divide  $|S|$ . But since  $p$  does not divide  $|S|$ ,  $|O_s| = 1$ . Then by the counting theorem  $|G_s| = |G|$ . So  $s$  stabilizes the whole group.

□

**Exercise (7.3.3).** *A nonabelian group  $G$  has order  $p^3$ , where  $p$  is prime.*

1. *What are the possible orders of the center  $Z$ ?*
2. *Let  $x$  be an element of  $G$  that isn't in  $Z$ . What is the order of its centralizer  $Z(x)$ ?*
3. *What are the possible class equations for  $G$ ?*

*Proof.* 1. First we can rule out the case  $Z = 1$  using the class equation since it would imply  $p^3 = 1 + (\text{sum of } p \text{ and } p^2)$ . Which is impossible.

We can rule out  $p^3$  since it would imply the group is abelian.

We can rule out  $p^2$  since the centralizer for any element not in the center would have to be larger order than  $p^2$  and divide  $p^3$ , but not be equal to  $p^2$ .

So we are left with  $|Z| = p$ .

2. The centralizer must be larger than  $Z$  since it must contain  $x$ , but it cannot be the same size of the group  $p^3$ . And its order must divide the order of the group so we are left with  $p^2$ .

3. Applying the information above. We can only have  $1 \times p + p \times (p^2 - 1)$ .

□

**Exercise (7.3.4).** *Classify groups of order 8.*

*Proof.* For abelian groups we start with  $C_8$ , then by inspection we also have  $C_4 \times C_2$ , and  $C_2 \times C_2 \times C_2$ .

For non-abelian groups, note that  $8 = 2^3$ . So we can apply part 3 of the previous problem and note that the class equation must be  $1 + 1 + 2 + 2 + 2$ . Also every non-identity element must be either order 2 or 4.

But if every element were order 2 the group would be abelian. So there is at least one element  $x$  of order 4.

□

**Exercise (7.4.7).** *Let  $G$  be a group of order  $n$  that operates nontrivially on a set of order  $r$ . Prove that if  $n > r!$ , then  $G$  has a proper normal subgroup.*

*Proof.*

□

**Exercise (7.4.9).** *Let  $x$  be an element of a group  $G$ , not the identity, whose centralizer  $Z(x)$  has order  $pq$ , where  $p$  and  $q$  are primes. Prove that  $Z(x)$  is abelian.*

*Proof.* Suppose  $Z(x)$  is not abelian.

$Z(x)$  is itself a group. Which must have a center containing  $x$  and 1, so  $|Z| > 1$ . Since the center is a subgroup its order must divide the order of the group, so it is either  $p, q, pq$ .

If the order of  $Z$  is  $pq$ ,  $Z(x)$  is abelian and we are done.

With out loss of generality suppose the order is  $p$ , consider the the centralizer of some element  $y \in Z(x) - Z$ . Its centralizer must have order greater than  $p$  and it must divide  $pq$ , so it must be order  $pq$ . So it commutes with the whole group.

So all elements not in the center commute, and so does the center. But this is a contradiction, so  $Z(x)$  must be abelian.

□

**Exercise (7.5.3).** *Determine the orders of the elements of the symmetric group  $S_7$ .*

*Proof.* We have the obvious cases for 1 through 7 cycles which gives order 1, 2, 3, 4, 5, 6, 7 elements.

Then we have the orders that arise from products of disjoint cycles. We start counting these down from 7 cycles noting that 7 and 6 cycles cannot form disjoint products with any of the cycles in  $S_7$ .

With 5 cycles we can form products with disjoint 2 cycles yielding an element of order 10.

With 4 cycles we can form products with disjoint 2 and 3 cycles yielding elements of order 8, and 12.

Counting lower results in double counting so our full list of the order of all elements of  $S_7$  is 1, 2, 3, 4, 5, 6, 7, 8, 10, 12.

□

**Exercise (7.5.6).** *Find all subgroups of  $S_4$  of order 4, and decide which ones are normal.*

*Proof.* We have the obvious cases, the subgroups generated by any of the 12 ( $4!$ ) four cycles. The canonical example of these is  $\{1, (\mathbf{1234}), (\mathbf{13})(\mathbf{24}), (\mathbf{1432})\}$ .

Then by inspection we note that 2 disjoint transpositions generate subgroups of order 4. There 6 ( $4$  choose  $2$ ) of these, the canonical example being  $\{1, (\mathbf{12}), (\mathbf{34}), (\mathbf{12})(\mathbf{34})\}$ .

So in total there are 18 subgroups of order 4 in  $S_4$ .

□

**Exercise (7.5.7).** *Prove that  $A_n$  is the only subgroup of  $S_n$  of index 2.*

*Proof.*

□

**Exercise (7.5.12).** *Determine the class equations of  $S_6$  and  $A_6$ .*

*Proof.*

□

**Exercise (7.6.4).** Let  $H$  be a normal subgroup of prime order  $p$  in a finite group  $G$ . Suppose that  $p$  is the smallest prime that divides the order of  $G$ . Prove that  $H$  is in the center  $Z(G)$ .

*Proof.* □

**Exercise (7.6.5).** Let  $p$  be a prime integer and let  $G$  be a  $p$ -group. Let  $H$  be a proper subgroup of  $G$ . Prove that the normalizer  $N(H)$  of  $H$  is strictly larger than  $H$ , and that  $H$  is contained in a normal subgroup of index  $p$ .

*Proof.* □

## Pre-Lecture Problems

**Exercise (7.5.1).** 1. Prove that the transpositions  $(\mathbf{12}), (\mathbf{23}), \dots, (\mathbf{n-1, n})$  generate the symmetric group  $S_n$ .

2. How many transpositions are needed to write the cycle  $(\mathbf{123 \dots n})$ ?

3. Prove that the cycles  $(\mathbf{12 \dots n})$  and  $(\mathbf{12})$  generate the symmetric group  $S_n$ .

*Proof.* 1. Let some arbitrary cycle be  $(\mathbf{i_1, i_2, \dots, i_j})$ . Then we can generate this with the given transpositions. First we note that we can generate adjacent indicies as follows

$$(\mathbf{i_1, i_1 \pm 1})(\mathbf{i_1 \pm 1, i_1 \pm 2}) \dots (\mathbf{i_1 \pm k_1, i_2}) = (\mathbf{i_1, i_2})$$

Simillarly  $(\mathbf{i_2, i_3}) = (\mathbf{i_2, i_2 \pm 1}) \dots (\mathbf{i_2 \pm k_2, i_3})$ . So we can write  $(\mathbf{i_1, i_2, i_3}) = (\mathbf{i_1, i_2})(\mathbf{i_2, i_3})$ .

Generating the rest of the given cycle follows by induction.

2.  $n - 1$  by counting:

$$(\mathbf{123 \dots n}) = (\mathbf{12})(\mathbf{23}) \dots (\mathbf{n-1, n})$$

$\begin{matrix} 1 & 2 & & n-1 \end{matrix}$

3. Using the given cycles we can generate any adjacent transpostions since  $(\mathbf{i, i+1}) = (\mathbf{123 \dots n})^{i+1}(\mathbf{12})(\mathbf{123 \dots n})^{1-i}$ . Then applying the first part of this problem we can generate  $S_n$ .

□