General Physics I Classnotes

Jonathan Henrique Maia de Moraes (ID: 1620855)

01/27/2016

January 22

1 Metric Prefixes

10^{12}	=1,000,000,000,000	tera	T
10^{9}	=1,000,000,000	giga	G
10^{6}	=1,000,000	mega	M
10^{3}	=1,000	kilo	k
10^{0}	= 1	_	_
10^{-2}	= 0.01	centi	c
10^{-3}	= 0.001	milli	m
10^{-6}	= 0.000,001	micro	μ
10^{-9}	=0.000,000,001	nano	n
10^{-12}	= 0.000, 000, 000, 001	pico	p

2 Basic Quantities

	Metric	English
Length	m = meter	ft = foot
Mass	kg = kilogram	sl = slug
Time	s = second	s = second

$$1 \text{day} = 24 \times 60 \times 60 = 86,400 \ s$$

$$1 \text{day} = 10 \times 100 \times 100 = 100,000 \ s$$

JANUARY 22 ii

3 Derived Quantities

velocity/speed	mi/s	km/h	m/min	 [L]/[T]
area	cm^2	m^2		 $[L]^2$
density	g/cm^3	kg/m^3		 $[M]/[L]^3$

4 Conversions

$$\begin{array}{ccccc}
1 & min & \equiv & 60 & s \\
1 & h & \equiv & 60 & min \\
1 & ft & \equiv & 12 & in \\
1 & mi & \equiv & 5280 & ft \\
1 & L & \equiv & 1,000 & cm^3 \\
1 & mi^2 & \equiv & 640 & acres \\
1 & in & \equiv & 2.54 & cm
\end{array}$$

Example:

$$70 \ mi/h = ? \ m/s$$

$$= 70 \ mi/h \times \left(\frac{5280 \ ft}{1 \ mi}\right) \times \left(\frac{12 \ in}{1 \ ft}\right) \times \left(\frac{2.54 \ cm}{1 \ in}\right) \times \left(\frac{1 \ m}{100 \ cm}\right)$$

$$\times \left(\frac{1 \ h}{60 \ min}\right) \times \left(\frac{1 \ min}{60 \ s}\right)$$

$$= 31.2928 \ m/s$$

$$350in^{3} = ? L$$

$$= 350 in^{3} \times \left(\frac{2.54 cm}{1 in}\right)^{3} \times \left(\frac{1 L}{1000 cm^{3}}\right)$$

$$= 5.7355 L$$

JANUARY 22

Homework:

$$\begin{aligned} 1acre &= ? \ in^2 \\ &= 1 \ acre \times \left(\frac{1 \ mi^2}{640 \ acres}\right) \times \left(\frac{5280 \ ft}{1 \ mi}\right)^2 \times \left(\frac{12 \ in}{1 \ ft}\right)^2 \\ &= 6,272,640 \ in^2 \end{aligned}$$

January 25

5 Position

Let

$$x = \text{position}$$

 $x_i = \text{initial position}$
 $x_f = \text{final position}$
 $\Delta x = \text{Displacement}$
 $= x_f - x_i$

Example:

$$x_{i} = +3 ft$$

$$x_{f} = +5 ft$$

$$\Delta x = x_{f} - x_{i}$$

$$= 5 ft - 3 ft$$

$$= +2 ft$$

$$x_i = +5 ft$$

$$x_f = -1 ft$$

$$\Delta x = x_f - x_i$$

$$= -1 ft - 5 ft$$

$$= -6 ft$$

JANUARY 25 v

Example:

$$x_i = +3 ft$$

$$x_2 = +5 ft$$

$$x_f = -1 ft$$

$$\Delta x = x_f - x_i$$

$$= -1 ft - 3 ft$$

$$= -4 ft$$
Distance Traveled = 2 ft + 6 ft
$$= 8 ft$$

6 Velocity

$$\begin{split} \bar{v} &= \text{average velocity} \\ \bar{v} &\equiv \frac{\Delta x}{\Delta t} = \frac{\text{displacement}}{\text{time elapsed}} \\ \text{average speed} &= \frac{\text{distance travelled}}{\text{time elapsed}} \end{split}$$

Start at
$$x = +3$$
 ft
Move to $x = +5$ ft
End at $x = -1$ ft
Trip takes 4 s
Find a) average velocity
b) average speed

$$\bar{v} \equiv \frac{\Delta x}{\Delta t}$$

$$= \frac{-1 ft - 3 ft}{4 s} = \frac{-4 ft}{4 s} = -1 ft/s$$
average speed =
$$\frac{\text{distance}}{\text{time}} = \frac{8 ft}{4 s} = 2 ft/s$$

JANUARY 25 vi

$$v = \text{instantaneous velocity}$$

$$= \lim_{\Delta t \to 0} \frac{\Delta x}{\Delta t}$$

$$v \equiv \frac{dx}{dt}$$

$$x = 3 \ m + (17 \ m/s)t + (7 \ m/s^3)t^3$$
 Find
$$a) \text{position at } t = 2 \ s$$

$$b) \text{position at } t = 4 \ s$$

$$c) \text{average velocity from } 2 \ s \to 4 \ s$$

a)

$$x = 3 m + (17 m/s)(2 s) + (7 m/s^{3})(2 s)^{3}$$

$$= 3 m + 34 m + 56 m$$

$$= 93 m$$
(1)

b)
$$x = 3 m + (17 m/s)(4 s) + (7 m/s^3)(4 s)^3$$
$$= 3 m + 68 m + 448 m$$
$$= 519 m$$

c)
$$\bar{v} = \frac{\Delta x}{\Delta t} = \frac{519 \ m - 93 \ m}{4 \ s - 2 \ s}$$
$$= \frac{426 \ m}{2 \ s}$$
$$= 213 \ m/s$$

JANUARY 25 vii

d)

$$v = \frac{dx}{dt}$$

$$= \frac{d}{dt} \left[3 m + (17 m/s)t + (7 m/s^3)t^3 \right]$$

$$= 0 + 17 m/s + (21 m/s^3)t^2$$

$$v(3 s) = 17 m/s + (21 m/s^3)(3 s)^2$$

$$= 17 m/s + 189 m/s = 208 m/s$$

January 27

7 Summary

$$x = \text{position}$$

$$\Delta x = \text{displacement}$$

$$= x_f - x_i$$

$$\bar{v} = \text{average velocity}$$

$$= \frac{\Delta x}{\Delta t}$$

$$v = \text{instantaneous velocity}$$

$$= \frac{dx}{dt} = \text{slope of x vs. t}$$
Avg Speed = $\frac{\text{distance traveled}}{\text{time elapsed}}$

8 Acceleration

Let
$$\bar{a}=$$
 average acceleration
$$\bar{a}\equiv \frac{\Delta v}{\Delta t} = \frac{v_f - v_i}{\Delta t} = \frac{\text{change in velocity}}{\text{time elapsed}}$$

Example: A car goes from $20 \ mph$ to $60 \ mph$ in $8 \ s$. What is its average acceleration?

JANUARY 27 ix

$$v_{i} = 20 \ mi/h$$

$$v_{f} = 60 \ mi/h$$

$$\Delta t = 8 \ s$$

$$\bar{a} = \frac{\Delta v}{\Delta t} = \frac{v_{f} - v_{i}}{\Delta t}$$

$$= \frac{60 \ mi/h - 20 \ mi/h}{8 \ s}$$

$$= \frac{40 \ mi/h}{8 \ s}$$

$$= 5 \ \frac{mi}{h \times s}$$

Example: Justin Bieber's Limo goes from $30 \ m/s$ to a stop in $0.10 \ s$. What is its average acceleration?

$$v_{i} = 30 \ m/s$$

$$v_{f} = 0 \ m/s$$

$$\Delta t = 0.10 \ s$$

$$\bar{a} = \frac{\Delta v}{\Delta t} = \frac{v_{f} - v_{i}}{\Delta t}$$

$$= \frac{0 \ m/s - 30 \ m/s}{0.10 \ s}$$

$$= \frac{-30 \ m/s}{0.10 \ s}$$

$$= -300 \ \frac{m/s}{s} = -300m/s^{2}$$

(- means slowing)

JANUARY 27 x

Let

$$a = \text{instantaneous acceleration}$$

$$= \lim_{\Delta t \to 0} \frac{\Delta v}{\Delta t}$$
 $a \equiv \frac{dv}{dt} = \text{rate of change of velocity}$

$$= \text{slope of tangent line to v vs. t}$$

Example:

$$x = 3 m + (17 m/s) t + (7 m/s^3) t^3$$

Find: a) velocity at 3 s

- b) velocity at 5 s
- c) average acceleration from $3 s \rightarrow 5 s$
- c) instantaneous acceleration at 4 s

a)

$$v = \frac{dx}{dt} = 17 \ m/s + (21 \ m/s^3) t^2$$
$$v(3 \ s) = 17 \ m/s + (21 \ m/s^3) (3 \ s)^2$$
$$= 17 \ m/s + 189 \ m/s$$
$$= 206 \ m/s$$

b)

$$v(5 s) = 17 m/s + (21 m/s^3) (5 s)^2$$

= 17 m/s + 525 m/s
= 542 m/s

c)

$$\bar{a} = \frac{\Delta v}{\Delta t} = \frac{v_f - v_i}{\Delta t} = \frac{542 \ m/s - 206 \ m/s}{5 \ s - 3 \ s}$$
$$= \frac{336 \ m/s}{2 \ s} = 168 \ m/s^2$$

JANUARY 27 xi

d)

$$a = \frac{dv}{dt}$$

$$= \frac{d}{dt} \left[17 \ m/s + (21 \ m/s^3)t^2 \right]$$

$$= 0 + (42 \ m/s^3)t$$

$$a(4 \ s) = (42 \ m/s^3)(4 \ s)$$

$$= 168 \ m/s^2$$

Suppose a = constant

$$\bar{a} = a
\frac{v - v_0}{t} = a
v - v_0 = at
v = v_0 + at : v(t)
x = x_0 + v_0 t + \frac{1}{2} a t^2 : x(t)
x = x_0 + \frac{1}{2} (v_0 + v) t : \text{no } a$$
(2)

$$2a(x - x_0) = v^2 - v_0^2 : \text{ no } t$$
 (5)