

Dräger

Anaesthetics are inhaled by patients.

However, volatile anaesthetics only produce 0.1% of US greenhouse emissions, but their long-term environmental impact must be considered.

Only **<5%** are metabolised


Isoflurane and nitrous oxides also damage the ozone layer.

A diagram of Earth showing the atmosphere. The boundary layer is the lowest part of the atmosphere, closest to the surface. It is shown as a thin layer at the bottom of the atmosphere, with a dashed line indicating its upper boundary. The boundary layer is labeled 'Boundary layer' and 'Boundary layer height'.

Desflurane has a global warming potential of 2,540, meaning the emissions generated by 1 kg of desflurane has the same effect as emitting **2.54 tonnes of CO₂**.

Putting it into perspective

■ Sevoflurane ■ Isoflurane ■ Nitrous Oxide (60%) ■ Desflurane



Sources: Sherman J.D. & Berkow L. (2019). Scaling up inhaled anesthetic practice improvement: The role of environmental sustainability metrics. *Anesth Analg.* 128(6), pp. 1060-1062.

Dräger anaesthesia machines support a safe and convenient application of minimal- and low-flow anaesthesia allowing you to reduce your ecological footprint.

Gas Consumption Analytics allows you to discover clinical, financial, and environmental insights into your anaesthetic gas consumption, e.g. the carbon dioxide pollution equivalent.



Cost savings