

WE BELIEVE THAT SMARTER ANESTHESIA DOSING INCREASES PATIENT SAFETY

Dräger

Keeping the right balance is hard when mixing anesthesia drugs.


With Dräger's SmartPilot View the resulting anesthetic depth from a combination of IV and volatile drugs is calculated based on proven population models [1] and visualized intuitively.

Information from therapy and IV-pumps is automatically transferred to Smart Pilot View. It is plug and play.


Additionally, manual entry is possible, e.g. for manual bolus.



One dimensional view: Resulting level of anaesthesia vs. time



Two-dimensional view: Resulting level of anaesthesia by the analgetic and anaesthetic agent concentrations (isoboles)



Abbreviations

TOSS 50 Tolerance of Shake and Shout where 50% of patients do not react

TOSS 90 Tolerance of Shake and Shout where 90% of patients do not react

MAC scale: Used on mixed anaesthesia.

Example:

MAC90: 90% of patients do not react on surgical stimuli (comparable to MAC 1.3); MAC50 would be comparable to MAC 1.0

TOL Scale: Tolerance On Laryngoscopy; used on Total Intravenous Anaesthesia.

Example:

TOL 90: 90% of patients tolerate a laryngoscopy

[1] Volatile agents: Bailey, Yasuda, Eger Bailey JM. The pharmacokinetics of volatile anesthetic agent elimination Minto et al. determined the PK-characteristics of Remifentanyl with a population of Minto CF, Schnider TW, Minto CF, Gambus PL, Andresen C, Goodale DB, Shafer SL, Youngs EJ. The influence of method of administration and covariates on the pharmacokinetics of Propofol in adult volunteers. Anesthesiology 1998; 88(5): 1170-82.