Lecture 2 (June 24)

Wednesday, June 24, 2026 12:01 PM

chapter1b_naive_pro...

Chapter 1B: Naive Probability

MATH/STAT 394: Probability I

Arman Jahangiri

University of Washington Department of Mathematics

Summer 2026

Learning Goals

By the end of this lecture, students should be able to:
- state the naive definition of probability;
- identify when the naive definition is appropriate;
- compute probabilities by counting favorable outcomes;
- use complements to simplify probability calculations;

Naive Probability

From Events to Probabilities

$f(x) = x^2$ $\mathbb{R} \to \mathbb{R}^{\geq 0}$

In the previous lecture:

S = {all possible outcomes}, event $A \subseteq S$.

S: sample space, $A \subseteq S$, $A \in P(S)$

Now we want to assign a number to an event:

$\mathbb{P}: P(A) \to [0,1]$

$\mathbb{P}(A)$ = probability that A occurs.

$A \mapsto [0,1]$

Question

If every outcome in a finite sample space is equally likely, how should we define $\mathbb{P}(A)$?

$S = \{1,2,3\} \Rightarrow P(S) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1,2\}, \{2,3\}, \{1,3\}, \{1,2,3\}\}$

$\mathbb{P}(\{1\})$ $\mathbb{P}(\{1,2\})$

Definition: Naive Probability

example) roll a die $S = \{1,2,\ldots,6\}$, $|S| = 6$, $P(\{1,2,3\}) = \frac{|\{1,2,3\}|}{S} = \frac{3}{6} = \frac{1}{2}$

Naive Definition of Probability

Let A be an event for an experiment with a finite sample space S.
If all outcomes in S are equally likely, define

$$\mathbb{P}_{\text{naive}}(A) = \frac{|A|}{|S|} = \frac{\text{number of outcomes favorable to } A}{\text{total number of outcomes in } S}.$$

Here $|A|$ denotes the number of elements in the set A (aka *card*(A)).
Question: Why does this definition is suitable for equally-likely outcomes?

Naive Probability - Properties

Naive probability satisfies

$$0 \leq \mathbb{P}(A) \leq 1, \ \mathbb{P}(S) = 1, \ \mathbb{P}(\emptyset) = 0, \quad \text{for every } A \subseteq S.$$

Why? First, if $A \subseteq S$, then

$$0 \leq |A| \leq |S|,$$

so $|A| \leq |S| \Rightarrow \frac{|A|}{|S|} \leq 1$

$$0 \leq \frac{|A|}{|S|} \leq 1.$$

$E = A \cup B$ $F = A \cap B$

Also,

$$\mathbb{P}(S) = \frac{|S|}{|S|} = 1, \quad \mathbb{P}(\emptyset) = \frac{0}{|S|} = 0.$$

Additivity for Disjoint Events

Suppose $A, B \subseteq S$,
Then, since

$$|A \cup B| = |A| + |B| - |A \cap B|$$

Therefore, (naive def. of prob.)

$$\mathbb{P}(A \cup B) = \frac{|A \cup B|}{|S|} = \frac{|A| + |B| - |A \cap B|}{|S|} = \frac{|A|}{|S|} + \frac{|B|}{|S|} - \frac{|A \cap B|}{|S|}$$

$|A \cup B| = |A| + |B| - |A \cap B|$

Thus

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$

A speical case is when A and B are exclusive ($A \cap B = \emptyset$), when we have

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$$

$|A| + |B|$

Looking ahead

Later, we will define probability more generally using properties like these as axioms. Naive probability will then become a special case of the general theory.

What the Definition Means

The naive definition says:

$$\mathbb{P}(A) = \frac{\text{how many outcomes make } A \text{ happen}}{\text{how many outcomes are possible}}.$$

This requires two assumptions:
1. The sample space S is finite (we can't divide by infinity).
2. All outcomes in S are equally likely.

Important

The naive definition is not wrong. It is just limited. It should only be used when its assumptions are justified.

Pebble World Interpretation

Suppose each pebble has the same mass.

$$\mathbb{P}(A) = \frac{\text{number of pebbles in } A}{\text{number of pebbles in } S}$$



$$\mathbb{P}(A) = \frac{|A|}{|S|}.$$

Example: One Fair Die

Roll one fair six-sided die.

$$S = \{1, 2, 3, 4, 5, 6\}.$$

Let

$$A = \{\text{roll is even}\} = \{2, 4, 6\}.$$

Then

$$|S| = 6, \qquad |A| = 3.$$

Therefore

$$\mathbb{P}(A) = \frac{|A|}{|S|} = \frac{3}{6} = \frac{1}{2}.$$

Example: One Card

Draw one card from a standard well-shuffled 52-card deck.
Let

$$A = \{\text{card is an ace}\}.$$

There are 4 aces and 52 cards total, so

$$\mathbb{P}(A) = \frac{|A|}{|S|} = \frac{4}{52} = \frac{1}{13}.$$

Let

$$H = \{\text{card is a heart}\}.$$

Then

$$\mathbb{P}(H) = \frac{13}{52} = \frac{1}{4}.$$

Example: Red or Ace

Draw one card from a standard 52-card deck.
Let

$$R = \{\text{card is red}\}, \qquad A = \{\text{card is an ace}\}.$$

and

We want

$$\mathbb{P}(R \cup A).$$

intersection of R & A

There are:

26 red cards, 4 aces, 2 red aces.

So

$$|R \cup A| = 26 + 4 - 2 = 28.$$

Therefore

$$\mathbb{P}(R \cup A) = \frac{28}{52} = \frac{7}{13}.$$

$P(R \cup A) = P(R) + P(A) - P(R \cap A) = \frac{26}{52} + \frac{4}{52} - \frac{2}{52} = \frac{7}{13}$

Complements

Complement Rule Under the Naive Definition

Let $A \subseteq S$.
The complement of A is

$$A^c = \{s \in S : s \notin A\}.$$

$P(A^c) = 1 - P(A)$

Since every outcome is either in A or not in A,

$$|A^c| = |S| - |A|.$$

$A \subseteq S$, $A^c = S - A$

Therefore,

$$\mathbb{P}(A^c) = \frac{|A^c|}{|S|} = \frac{|S| - |A|}{|S|} = 1 - \frac{|A|}{|S|}.$$

So

$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$$

$A \cup A^c = S$

$P(A^c) = 1 - P(A) \Leftarrow 1 = P(S) = P(A \cup A^c) = P(A) + P(A^c) - P(A \cap A^c)$

Why Complements Are Useful

Sometimes it is easier to count when an event does **not** happen.
Instead of computing $\mathbb{P}(A)$ directly, compute:

$$\mathbb{P}(A) = 1 - \mathbb{P}(A^c).$$

Strategy

When a probability asks for phrases like "at least one", "at least two", "not all", first ask whether the complement is easier.

if you see these keywords then think about computing both $P(A)$ & $P(A^c)$. And then compute the one that is easier.

Example: At Least One Head

$S = \{(s_1, s_2, s_3, s_4, s_5) : s_i = H \text{ or } s_i = T\}$ $|S| = 2^5$

Flip a fair coin 5 times.
Let

$$A = \{\text{at least one head occurs}\}.$$

let X = number of heads, then A says that $\{X \geq 1\}$

The complement is

$$A^c = \{\text{no heads occur}\}.$$

$\Rightarrow A^c : \{X < 1\}$ $\Rightarrow A^c : \{X = 0\}$ (no heads) all tails

There is only one outcome with no heads: $TTTTT$.
The total number of outcomes is $2^5 = 32$.
Therefore

$$\mathbb{P}(A^c) = \frac{1}{32}, \qquad \mathbb{P}(A) = 1 - \frac{1}{32} = \frac{31}{32}.$$

Example: At Least One Six

(at home)

Roll a fair die 4 times.
Let

$$A = \{\text{at least one six occurs}\}.$$

It is easier to count

$$A^c = \{\text{no six occurs}\}.$$

Total outcomes: 6^4. Outcomes with no six: 5^4.
Therefore,

$$\mathbb{P}(A) = 1 - \mathbb{P}(A^c) = 1 - \frac{5^4}{6^4}.$$

Common Mistakes

Example: Two Fair Dice

Roll two fair dice.
A correct sample space is

$$S = \{(i, j) : i, j \in \{1, 2, 3, 4, 5, 6\}\}.$$

$(1,1), (1,2), (1,3), \ldots (1,6)$
$(2,1), (2,2), \ldots (2,6)$
$(6,1), (6,2), \ldots (6,6)$

Here (i, j) means:

$$i = \text{result of die 1}, \qquad j = \text{result of die 2}.$$

let X = first die, Y = second die

Thus

$$|S| = 36 = 6^2$$

$P(X + Y = 11) = \frac{2}{36}$

$P(X + Y = 12) = \frac{1}{36}$

Important

The outcomes are ordered pairs. For example, $(5, 6)$ and $(6, 5)$ are different outcomes.

Example: Sum 11 versus Sum 12

For two fair dice:

$$S = \{(i, j) : i, j \in \{1, \ldots, 6\}\}.$$

Let

$$A = \{\text{sum is } 11\}, \qquad B = \{\text{sum is } 12\}.$$

Then

$$A = \{(5, 6), (6, 5)\}, \qquad B = \{(6, 6)\}.$$

So

$$\mathbb{P}(A) = \frac{2}{36} = \frac{1}{18},$$

while $\mathbb{P}(B) = \frac{1}{36}$.
Thus a sum of 11 is twice as likely as a sum of 12.

Mini-Exercises

(at home)

1. A fair die is rolled. What is the probability of rolling a number at least 5?
2. A fair coin is flipped 4 times. What is the probability of getting all tails?
3. A card is drawn from a standard deck. What is the probability that it is not a heart?
4. Two fair dice are rolled. What is the probability that the sum is 7?
5. Why is the statement "the probability of rain tomorrow is 1/2 because either it rains or it does not" invalid?

Answers

$\frac{2}{6} = \frac{1}{3}$, $\frac{1}{16}$, $1 - \frac{13}{52} = \frac{3}{4}$, $\frac{6}{36} = \frac{1}{6}$

Summary

- The naive definition is
$$\mathbb{P}(A) = \frac{|A|}{|S|}.$$
- It applies when S is finite and all outcomes are equally likely.
- Naive probability satisfies:
$$0 \leq \mathbb{P}(A) \leq 1, \quad \mathbb{P}(S) = 1, \quad \mathbb{P}(\emptyset) = 0$$
If $A \cap B = \emptyset$, then
$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B).$$
- Complements are often useful:
$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$$

Next lecture

Counting methods: multiplication rule, sampling with replacement, and sampling without replacement.

---

chapter1cd_counting

Chapter 1C–1D: Counting, Multiplication Rule, and Binomial Coefficients

MATH/STAT 394: Probability I

Arman Jahangiri

University of Washington Department of Mathematics

Summer 2026

Learning Goals

By the end of this lecture, students should be able to:
- apply the multiplication rule for counting;
- distinguish between sampling with and without replacement;
- compute permutations and ordered selections;
- compute combinations and unordered selections;
- understand the binomial coefficient combinatorially;
- recognize and avoid overcounting;
- prove combinatorial identities using counting arguments.

The Multiplication Rule (Product Rule)

The Basic Counting Principle

Multiplication Rule

Suppose a procedure consists of two stages:
- Stage 1 can be done in a ways;
- for each choice in Stage 1, Stage 2 can be done in b ways.

Then the entire procedure can be done in

$$ab$$

ways.

More generally, if a process has stages with

$$n_1, n_2, \ldots, n_k$$

possible choices respectively, then the total number of outcomes is $n_1 n_2 \cdots n_k$.

Tree Interpretation

The multiplication rule can be visualized using a tree.

Suppose:
- there are 3 choices for the first step;
- each leads to 2 choices for the second step.

Then, according to the Multiplication Rule, there are $3 \times 2 = 6$ possible outcomes.



Example: License Plates

Suppose a license plate consists of 3 letters, followed by 4 digits. Assume repetition is allowed.



Figure: WA license plate. Source: Wikipedia.

26 . 26 . 26 . 10 . 10 . 10 . 10
3 letters 4 digits

Choices:

26 possibilities for each letter,
10 possibilities for each digit.

Thus the total number of possible license plates is

$$(26 \times 26 \times 26) \times (10 \times 10 \times 10 \times 10) = 26^3 \, 10^4 = 175,760,000$$

Example: Binary Strings

10011010 a binary string of size 8

A binary string of length n consists only of 0's and 1's.
Each position has:

2 choices.

Using the multiplication rule:

$$\underset{\text{digit 1}}{2} \times \underset{\text{digit 2}}{2} \times \cdots \times \underset{\text{digit } n}{2} = 2^n.$$

Therefore, Number of binary strings of length n is 2^n.
Remark: This is also equal to the number of all possible subsets of a set with n elements. (Why?)

Sampling With and Without Replacement

sample 3 balls (without replacement)

Ordered Sampling With Replacement

Suppose we have a set of size n, and we perform k draws with replacement, where $1 \leq k \leq n$.
With replacement means: after each draw, the selected object is returned before the next draw.
Therefore, each draw has exactly n possible choices.
Using the multiplication rule:

$$\underset{\text{draw 1}}{n} \times \underset{\text{draw 2}}{n} \times \cdots \times \underset{\text{draw } k}{n} = n^k$$

Hence, the number of possible ways to draw k ordered items, with replacement, from a set of size n, is

$$n^k$$

Example: PIN Codes

A 4-digit PIN code uses digits 0–9.
Digits may repeat.
Each position has:

10 choices.

Therefore:

$$10^4 = 10000$$

possible PINs.

Ordered Sampling Without Replacement

Now suppose selected objects are not returned after each draw.
Then the number of available choices decreases after every selection.
Using the multiplication rule:

for any $n \in \mathbb{N}$, $n! = n(n-1)\cdots$

$n - (k-1)$ (Gamma)

$$\underset{\text{draw 1}}{n} \times \underset{\text{draw 2}}{n-1} \times \underset{\text{draw 3}}{n-2} \times \cdots \times \underset{\text{draw } k}{n-k+1}$$
$$= n(n-1)(n-2)\cdots(n-k+1) = \frac{n!}{(n-k)!}$$

Important

Without replacement means that the choices depend on previous selections.

Example: Arranging Books

Suppose 5 distinct books are arranged on a shelf.
Choices:

5 choices for the first position,
4 for the second,
3 for the third,

etc.
Therefore, the number of arrangements is

$$5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5! = 120.$$

$n = 5$

Factorials and Permutations

Factorials

Definition

For a positive integer n,

$$n! = n(n-1)(n-2)\cdots 2 \cdot 1.$$

Also define

$$0! = 1.$$

Examples:

$$1! = 1, \quad 2! = 2, \quad 3! = 6, \quad 4! = 24, \quad 5! = 120.$$

Permutations

A **permutation** is an ordered arrangement.
The number of ordered ways to choose k objects from n distinct objects is

$$n(n-1)\cdots(n-k+1).$$

Using factorial notation:

$$\frac{n!}{(n-k)!}$$

This is denoted by

$$P(n, k).$$

So, in summary

$$P(n, k) = \frac{n!}{(n-k)!} = n(n-1)\cdots(n-k+1).$$

Example: Choosing Officers

Suppose a club has 12 members.
We choose:
- a president,
- a vice president,
- a secretary.

Since positions are different, order matters.
Therefore:

$$P(12, 3) = \frac{12!}{9!} = 12 \cdot 11 \cdot 10 = 1320.$$

$n = 12$, $k = 3$: $\frac{12!}{(12-3)!} = P(n,k) = P(12,3)$

Combinations

When Order Does Not Matter

Suppose we choose a committee of size k from n people.
Now:

$$(A, B, C)$$

and

$$(B, A, C)$$

represent the same committee.
Thus order does not matter.
This requires a different counting formula.

$\{A,B,C\} = \{A,C,B\} = \{B,A,C\} = \cdots$

Binomial Coefficients

(A,B,C): ABC, ACB, BAC, BCA, CAB, CBA; $3! = 6$

Definition

The number of subsets of size k from a set of size n is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

aka the binomial coefficient

choose n, k; aka $C(n,k)$

This is read as:

"n choose k".

Note: $\binom{n}{0} = \binom{n}{n} = \binom{0}{0} = 1$ can be justified by the formula or combinatorially.

$P(n,k) = \frac{n!}{(n-k)!}$, $C(n,k) = \frac{P(n,k)}{k!}$

Why Divide by $k!$?

Suppose we first count ordered selections:

$$P(n, k) = \frac{n!}{(n-k)!}$$

But each unordered subset is counted many times.
For example $\{A, B, C\}$ can appear as:

$$ABC, \ ACB, \ BAC, \ BCA, \ CAB, \ CBA.$$

There are $3! = 6$ orderings.
Therefore we divide by $k!$:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{P(n, k)}{k!} \Rightarrow \binom{n}{k} \leq P(n, k)$$

Example: Poker Hands

A poker hand consists of 5 cards from a 52-card deck.
Order does not matter.
Thus the number of poker hands is

$$\binom{52}{5} = \frac{52!}{5!47!}.$$

at home

Example: Committees

(at home)

How many committees of size 4 can be formed from 10 people?
Since order does not matter:

$$\binom{10}{4} = \frac{10!}{4!6!}$$

Simplifying:

$$\binom{10}{4} = 210.$$

Properties of Binomial Coefficients

Symmetry Identity

$\binom{n}{k} = \frac{n!}{k!(n-k)!}$, $\binom{n}{n-k} = \frac{n!}{(n-k)!(n-(n-k))!} = \frac{n!}{(n-k)!\,k!}$

Identity

$$\binom{n}{k} = \binom{n}{n-k}$$

Reason:
Choosing k elements to include is equivalent to choosing $n-k$ elements to exclude. :)
Algebraic Proof:

Khayyam–Pascal's Identity

Khayyam-Pascal's Identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

$\frac{n!}{k!(n-k)!}$

Algebraic Proof: (Boring)

$$\binom{n-1}{k} = \frac{(n-1)!}{k!(n-1-k)!}, \quad \binom{n-1}{k-1} = \frac{(n-1)!}{(k-1)!(n-k)!}$$

$$= \frac{(n-1)!\,(n-k)}{k!\,(n-k)!} + \frac{(n-1)!\,k}{k!\,(n-k)!} = \frac{(n-1)!\,[n-k+k]}{k!(n-k)!} = \frac{n!}{k!(n-k)!} = \binom{n}{k}$$

Combinatorial Proof of Khayyam-Pascal's Identity

Suppose a committee of size k is selected from n people.
Focus on one distinguished person.
Two possibilities:
1. The distinguished person is not selected.
   Then choose all k members from the remaining $n-1$ people:
$$\binom{n-1}{k}$$
2. The distinguished person is selected.
   Then choose the remaining $k-1$ members from the remaining $n-1$ people:
$$\binom{n-1}{k-1}$$

Adding the two cases gives (why addition?)

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

Khayyam–Pascal Triangle: The Main Idea

The Khayyam–Pascal triangle is a triangular arrangement of numbers.
Each row begins and ends with 1.
Every interior entry is obtained by adding the two entries directly above it:

new entry = upper-left entry + upper-right entry



Source: Wikipedia: Pascal's Triangle

$\binom{3}{1} = 3$

Building the Triangle

future: $(a+b)^n = \sum_{k=0}^{n} \binom{n}{k} a^k b^{n-k}$ important identity (for Binomial distribution)

Start with 1 at the top.
Each new number is the sum of the two numbers above it.

1
1 1
1 1+1 1
1 1+2 2+1 1
1 1+3 3+3 3+1 1

The Formula Inside the Triangle

$\binom{n}{k} = \binom{n}{n-k}$

$\binom{n}{n} = \frac{n!}{n!\,(n-n)!} = 1$

row 0: $\binom{0}{0}$
row 1: $\binom{1}{0}$ $\binom{1}{1}$
row 2: $\binom{2}{0}$ $\binom{2}{1}$ $\binom{2}{2}$
row 3: $\binom{3}{0}$ $\binom{3}{1}$ $\binom{3}{2}$ $\binom{3}{3}$
row $n-1$: $\binom{n-1}{0}$ … $\binom{n-1}{k-1}$ $\binom{n-1}{k}$ … $\binom{n-1}{n-1}$
row n: $\binom{n}{0}$ … $\binom{n}{k}$ … $\binom{n}{n}$

Each interior entry is obtained by adding the two entries directly above it:

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

This identity is known as **Khayyam-Pascal's Rule**.

$(a+b)^2 = a^2 + 2ab + b^2$

Khayyam-Pascal's Rule/Identity

(at home)

The rule for building the triangle is

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

for $1 \leq k \leq n-1$.

For example,

$$\binom{4}{2} = \binom{3}{1} + \binom{3}{2} = 3 + 3 = 6.$$

So every interior entry is obtained by adding the two entries above it.

Connection to Binomial Expansions

The rows of the triangle give the coefficients in binomial expansions.

$$(a+b)^0 = 1$$
$$(a+b)^1 = a + b$$
$$(a+b)^2 = a^2 + 2ab + b^2$$
$$(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$
$$(a+b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$$

- The coefficients are exactly the rows of the Khayyam-Pascal triangle.
- Hence, we can show that $(a+b)^n = \sum_{k=0}^{n} \binom{n}{k} a^k b^{n-k}$ (a very useful identity in Probability theory)
- Also proves why $2^n = \sum_{k=0}^{n} \binom{n}{k}$
  Hint: $2 = 1 + 1$

$a = 1$, $b = 1$ $\Rightarrow$ $a + b = 2$

Story Proofs and Overcounting

Story Proofs

A **story proof** proves an identity by showing that both sides count the same object in different ways.
Advantages:
- gives intuition;
- explains why identities are true;
- often simpler than algebraic proofs.

Example Story Proof

Identity

$$\sum_{k=0}^{n} \binom{n}{k} = 2^n.$$

Why?
The left-hand side counts subsets of an n-element set by grouping according to subset size.
The right-hand side counts subsets directly (see slide 60):
each element is either:

included or not included

Thus there are

$$2^n$$

subsets total.

$(\underline{\ }\ \underline{\ }\ \cdots\ \underline{\ })$

bijection between all subsets of a set of size n & all binary sequences of length n

Mini-Exercises

at home

1. How many binary strings of length 8 are there?
2. How many 3-letter strings can be formed from the English alphabet if repetition is allowed?
3. How many ordered arrangements of 4 people can be selected from 10 people?
4. How many committees of size 3 can be formed from 8 people?

Answers

$2^8 = 256$, 26^3, $\frac{10!}{6!}$, $\binom{8}{3} = 56$.

Summary

- Number of ordered length-k sequences from n options (repetition allowed):
$$n^k.$$
- Number of ordered length-k sequences from n distinct objects (no repetition):
$$\frac{n!}{(n-k)!}.$$
- Number of permutations of k objects chosen from n:
$$P(n, k) = \frac{n!}{(n-k)!}.$$
- Number of combinations of k objects chosen from n:
$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$
- Fundamental identities for binomial coefficients:
$$\binom{n}{k} = \binom{n}{n-k},$$
and
$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

---

chapter1f_axiomatic_p...

Chapter 1F: Axiomatic Probability and the Addition Rule

MATH/STAT 394: Probability I

Arman Jahangiri

University of Washington Department of Mathematics

Summer 2026

Learning Goals

By the end of this lecture, students should be able to:
- state the axioms of probability;
- understand probability as a mathematical function on events;
- derive basic consequences of the axioms;
- apply the addition rule;
- use complements and inclusion–exclusion;
- identify mutually exclusive events;
- understand why axiomatic probability generalizes naive probability.

Why Axiomatize Probability (or anything)?

Limitations of Naive Probability

Previously, we defined (naive) probability by

$$\mathbb{P}(A) = \frac{|A|}{|S|}.$$

S $A \subseteq S$
S = set of outcomes
s_1 & s_2

This works well when:
- the sample space is finite; ✓
- all outcomes are equally likely; ✓

$P(\{s_1\}) = \frac{1}{n}$
$P(\{s_2\}) = \frac{1}{n}$

But many important situations do not satisfy these assumptions.
Examples:
- infinitely many outcomes;
- continuous measurements;
- unequal likelihoods;
- etc.

The Axiomatic Point of View

Instead of defining probability by counting, we define probability abstractly using properties that probabilities should satisfy.

Idea

A probability measure assigns a number to each event:

$$A \longmapsto \mathbb{P}(A),$$

subject to certain axioms (i.e., principles, rules) .

The naive definition will become a special case of the general theory.

The Axioms of Probability

Probability Space

A probability model consists of:
1. S sample space (set of all possible outcomes in an experiment);
2. $\mathcal{F}$ a collection of subsets of S (aka "events") ;
3. $\mathbb{P}(\cdot)$ a probability function $\mathbb{P} : \mathcal{F} \to [0, 1]$.

$\mathbb{P}: \mathcal{F} \to [0,1]$

So for any event $A \subseteq S$ (or $A \in \mathcal{F}$), $\mathbb{P}(A)$ is the probability that the function $\mathbb{P}(\cdot)$ assigns to A.
These 3 elements together form a Probability Model:

$$(S, \mathcal{F}, \mathbb{P})$$

Note that the probability function assigns numbers to sets (events), not to numbers (this might be new for us). So we will see things like $\mathbb{P}([1, 2])$, $\mathbb{P}((-\infty, 4])$, or $\mathbb{P}(\{2, 4, 6\})$

Axiom 1: Nonnegativity

Axiom 1

For every event $A \subseteq S$,

$$\mathbb{P}(A) \geq 0.$$

Interpretations:
- probabilities cannot be negative;
- impossible events have probability 0;
- larger probabilities correspond to more likely events.

Axiom 2: Normalization

Axiom 2

The probability that "something happens" is 1:

$$\mathbb{P}(S) = 1.$$

In other words, $\mathbb{P}(\cdot)$ assigns the number 1 to the event $S \in \mathcal{F}$.

Axiom 3: Additivity

$\emptyset \cap A_i = \emptyset$ $A_1, \ldots, A_n, \emptyset, \emptyset, \ldots$

Axiom 3

If

$$A_1, A_2, A_3, \ldots$$

are pairwise disjoint events, then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i).$$

$\Rightarrow P\left(\bigcup_{i=1}^{n} A_i\right) = \sum_{i=1}^{n} P(A_i)$

Pairwise disjoint means:

$$A_i \cap A_j = \emptyset \quad \text{whenever } i \neq j.$$

$P(A \cup B) = P(A) + P(B)$

Finite Additivity

Axiom 3 immediately implies the finite version:
If

$$A_1, \ldots, A_n$$

are disjoint, then

$$\mathbb{P}(A_1 \cup \cdots \cup A_n) = \mathbb{P}(A_1) + \cdots + \mathbb{P}(A_n).$$

In particular, if

$$A \cap B = \emptyset,$$

then

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B).$$

Basic Consequences of the Axioms

Probability of the Empty Set

Proposition

$$\mathbb{P}(\emptyset) = 0.$$

$S \cap \emptyset = \emptyset$, $\emptyset$ is disjoint from any set

Reason:

$\emptyset \subseteq S$
$\emptyset \notin S$

$$S = S \cup \emptyset,$$

and

$$S \cap \emptyset = \emptyset.$$

Using additivity,

$$\mathbb{P}(S) = \mathbb{P}(S) + \mathbb{P}(\emptyset).$$

$\Rightarrow P(S) = P(S \cup \emptyset) = P(S) + P(\emptyset)$ (Axiom 3) $\Rightarrow P(\emptyset) = 0$

Since $\mathbb{P}(S) = 1$, we obtain $\mathbb{P}(\emptyset) = 0$.

Complements

Proposition (Complement Rule)

For every event A,

$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A).$$

Reason:

$$A \cup A^c = S,$$

and

$$A \cap A^c = \emptyset.$$

Thus,

$$1 = \mathbb{P}(S) = \mathbb{P}(A) + \mathbb{P}(A^c).$$

Axiom 3, Axiom 2

$\Rightarrow P(A^c) = 1 - P(A)$

A, A^c disjoint

Probabilities Are Between 0 and 1

Proposition

For every event A,

$$0 \leq \mathbb{P}(A) \leq 1.$$

$A \subseteq S$

Why?
Axiom 1 gives:

$$\mathbb{P}(A) \geq 0.$$

Also,

$$\mathbb{P}(A^c) \geq 0.$$

Using

$$0 \leq \mathbb{P}(A^c) = 1 - \mathbb{P}(A),$$

we obtain

$$1 - \mathbb{P}(A) \geq 0 \Rightarrow 1 \geq P(A)$$

Monotonicity

Proposition (Monotonicity)

For any event $A, B \subseteq S$ if $A \subseteq B$, then

$$\mathbb{P}(A) \leq \mathbb{P}(B).$$

Proof.

If $A \subseteq B$, then $B = A \cup (B \setminus A)$, where the two events ($A$ and $B - A$) are disjoint (why?). Thus,

$$\mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \geq \mathbb{P}(A),$$

since $\mathbb{P}(B - A) \geq 0$ according to Axiom 1. □

Monotonicity: Picture



$$B = A \cup (B \setminus A)$$
$$A \cap (B \setminus A) = \emptyset$$

Since B contains all of A plus the extra part $B \setminus A$,

$$\mathbb{P}(B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \geq \mathbb{P}(A).$$

$B = (B - A) \cup A$

$P(B) = P(B - A) + P(A) \geq P(A)$ (Ax 1)

$A \subseteq B \Rightarrow P(A) \leq P(B)$

The Addition Rule

Why Finite Additivity Is Not Enough

Suppose A and B are not disjoint.
Then:

$$\mathbb{P}(A \cup B) \neq \mathbb{P}(A) + \mathbb{P}(B).$$

Why?
Because outcomes in

$$A \cap B$$

are counted twice.

(skip)

The Addition Rule

Proposition: Addition Rule

For any events A and B,

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B).$$

Proof.

Partition $A \cup B$ into three disjoint subsets and use a similar trick that was applied to prove Monotonicity previously. □

$A \cup B = (A - B) \cup (B - A) \cup (A \cap B)$

Why the Addition Rule Works

When adding

$$\mathbb{P}(A) + \mathbb{P}(B),$$

outcomes in

$$A \cap B$$

are counted twice.
Therefore we subtract one copy:

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B).$$

Mutually Exclusive Events

Definition

Events A and B are called **mutually exclusive** (or disjoint) if

$$A \cap B = \emptyset.$$

In that case,

$$\mathbb{P}(A \cap B) = 0,$$

so the addition rule becomes:

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B).$$

Example: Die Roll

Roll one fair die.
Let

$$A = \{\text{roll is even}\} = \{2, 4, 6\},$$
$$B = \{\text{roll is odd}\} = \{1, 3, 5\}.$$

Then

$$A \cap B = \emptyset.$$

(at home)

Thus

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B).$$

Indeed,

$$\frac{1}{2} + \frac{1}{2} = 1.$$

Inclusion–Exclusion

Three-Set Inclusion–Exclusion

For three events:

$$\mathbb{P}(A \cup B \cup C)$$

we first add:

$$\mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C).$$

Then subtract pairwise intersections:

$$-\mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C).$$

But now the triple intersection has been subtracted too many times, so we add it back.

Inclusion–Exclusion Formula

Three-Set Inclusion–Exclusion

$$\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C)$$
$$- \mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C)$$
$$+ \mathbb{P}(A \cap B \cap C).$$

Partitions

Partitions of the Sample Space

A_1 A_2 A_n S B

Definition

Events

$$A_1, \ldots, A_n$$

form a partition of S if:
1. $$A_i \cap A_j = \emptyset \quad (i \neq j),$$
2. $$A_1 \cup \cdots \cup A_n = S.$$

A partition divides the sample space into non-overlapping pieces.

Probability of a Partition

If

$$A_1, \ldots, A_n$$

form a partition of S, then

$$\mathbb{P}(A_1) + \cdots + \mathbb{P}(A_n) = 1.$$

Reason:
The events are disjoint and their union is S.

$$1 = P(S) = P\left(\bigcup_{i=1}^{n} A_i\right) = \sum_{i=1}^{n} P(A_i)$$

by (∗)

Naive Probability as a Special Case

Recovering Naive Probability

$P(A) = \frac{|A|}{|S|} \geq 0$

Suppose:
- S is finite;
- all outcomes are equally likely.

Let $S = \{s_1, s_2, \ldots, s_n\}$
Assign equal probability weights to each outcome:

$P(S) = \frac{|S|}{|S|} = 1$

$$\mathbb{P}(s_1) = \mathbb{P}(s_2) = \cdots = \mathbb{P}(s_n) = \frac{1}{n}.$$

Then for an event A, we have

$$\mathbb{P}(A) = \mathbb{P}(\cup_{s_i \in A} s_i) = \sum_{i \in A} \frac{1}{n} = |A| \times \frac{1}{n} = \frac{|A|}{|S|}.$$

Thus the naive definition satisfies the probability axioms.

Mini-Exercises

1. If
$$\mathbb{P}(A) = 0.3,$$
what is
$$\mathbb{P}(A^c)?$$
2. If
$$\mathbb{P}(A) = 0.6, \quad \mathbb{P}(B) = 0.5, \quad \mathbb{P}(A \cap B) = 0.1,$$
compute
$$\mathbb{P}(A \cup B).$$
3. Can two disjoint events both have probability 1?

Answers

0.7, 1.0.

Summary

- Probability is defined axiomatically.
- The axioms are:
$$\mathbb{P}(A) \geq 0, \qquad \mathbb{P}(S) = 1,$$
and countable additivity.
- Important consequences include:
$$\mathbb{P}(\emptyset) = 0, \qquad \mathbb{P}(A^c) = 1 - \mathbb{P}(A), \qquad 0 \leq \mathbb{P}(A) \leq 1.$$
- The addition rule is:
$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B).$$
- Naive probability is a special case of axiomatic probability.

Next lecture

Conditional probability and independence.