

Forward School

Program Code: J620-002-4:2020

Program Name: FRONT-END SOFTWARE DEVELOPMENT

Title : Hypothesis Testing Exercise In Python

Name: Ooi Caarob

IC Number: 990701-07-5837

Date : 16/7/2023

Introduction : Exercise caculating the value from the given data

Conclusion :Still need to practice more and do revision

Creating confidence intervals in python

In this assessment, you will look at data from a study on toddler sleep habits.

The confidence intervals you create and the questions you answer in this Jupyter notebook will be used to answer questions in the following graded assignment.

In [1]:

```
import numpy as np
import pandas as pd
from scipy.stats import t
pd.set_option('display.max_columns', 30) # set so can see all columns of the DataFrame
```

Your goal is to analyse data which is the result of a study that examined differences in a number of sleep variables between napping and non-napping toddlers. Some of these sleep variables included: Bedtime (lights-off time in decimalized time), Night Sleep Onset Time (in decimalized time), Wake Time (sleep end time in decimalized time), Night Sleep Duration (interval between sleep onset and sleep end in minutes), and Total 24-Hour Sleep Duration (in minutes). Note: [Decimalized time](https://en.wikipedia.org/wiki/Decimal_time) (https://en.wikipedia.org/wiki/Decimal_time) is the representation of the time of day using units which are decimally related.

The 20 study participants were healthy, normally developing toddlers with no sleep or behavioral problems. These children were categorized as napping or non-napping based upon parental report of children's habitual sleep patterns. Researchers then verified napping status with data from actigraphy (a non-invasive method of monitoring human rest/activity cycles by wearing of a sensor on the wrist) and sleep diaries during the 5 days before the study assessments were made.

You are specifically interested in the results for the Bedtime, Night Sleep Duration, and Total 24- Hour Sleep Duration.

ref: Akacem LD, Simpkin CT, Carskadon MA, Wright KP Jr, Jenni OG, Achermann P, et al. (2015) The Timing of the Circadian Clock and Sleep Differ between Napping and Non-Napping Toddlers. PLoS ONE 10(4): e0125181. <https://doi.org/10.1371/journal.pone.0125181>
(<https://doi.org/10.1371/journal.pone.0125181>)

In [2]:

```
# Import the data  
df = pd.read_csv("nap_no_nap.csv")
```

In [3]:

```
# First, Look at the DataFrame to get a sense of the data
df
```

Out[3]:

	id	sex	age (months)	dlmo time	days napped	napping	nap lights outl time	nap sleep onset	nap midsleep	nap sleep offset	nap wake time	du
0	1	female	33.7	19.24	0	0	NaN	NaN	NaN	NaN	NaN	
1	2	female	31.5	18.27	0	0	NaN	NaN	NaN	NaN	NaN	
2	3	male	31.9	19.14	0	0	NaN	NaN	NaN	NaN	NaN	
3	4	female	31.6	19.69	0	0	NaN	NaN	NaN	NaN	NaN	
4	5	female	33.0	19.52	0	0	NaN	NaN	NaN	NaN	NaN	
5	6	female	36.2	18.22	4	1	14.00	14.22	15.00	15.78	16.28	
6	7	male	36.3	19.28	1	1	14.75	15.03	15.92	16.80	16.08	1
7	8	male	30.0	21.06	5	1	13.09	13.43	14.44	15.46	15.82	1
8	9	male	33.2	19.38	2	1	14.41	14.42	15.71	17.01	16.60	1
9	10	female	37.1	19.93	3	1	13.12	13.42	14.31	15.19	15.30	1
10	11	male	32.9	18.79	4	1	13.99	14.03	14.85	15.68	16.10	
11	12	female	35.0	19.65	5	1	13.18	13.45	14.33	15.21	15.35	1
12	13	male	35.1	19.83	3	1	13.94	14.48	15.26	16.03	15.78	
13	14	female	35.6	19.88	4	1	12.68	13.08	13.92	14.76	15.00	1
14	15	female	36.6	19.94	4	1	12.71	12.88	13.80	14.72	14.88	1
15	16	male	36.5	20.25	3	1	13.74	14.68	15.66	16.64	16.45	1
16	17	female	33.7	20.33	5	1	13.15	13.87	14.49	15.11	15.40	
17	18	male	36.4	20.16	5	1	12.47	12.56	13.30	14.05	14.25	
18	19	female	33.6	19.68	3	1	14.71	14.85	15.46	16.07	16.20	
19	20	male	33.8	20.51	3	1	12.68	13.54	14.30	15.07	15.23	

Question: What variable is used in the column 'napping' to indicate a toddler takes a nap?

In [4]:

```
df['napping'].describe()
```

Out[4]:

```
count    20.000000
mean      0.750000
std       0.444262
min       0.000000
25%       0.750000
50%       1.000000
75%       1.000000
max       1.000000
Name: napping, dtype: float64
```

Question: What is the sample size n ?

In [5]:

```
n = len(df)
n
```

Out[5]:

20

Hypothesis testing

We will look at two hypothesis test, each with $\alpha = .025$:

1. Is the average bedtime for toddlers who nap later than the average bedtime for toddlers who don't nap?

$$H_0 : \mu_{nap} = \mu_{no\ nap}, \quad H_a : \mu_{nap} > \mu_{no\ nap}$$

Or equivalently:

$$H_0 : \mu_{nap} - \mu_{no\ nap} = 0, \quad H_a : \mu_{nap} - \mu_{no\ nap} > 0$$

2. The average 24 h sleep duration (in minutes) for napping toddlers is different from toddlers who don't nap.

$$H_0 : \mu_{nap} = \mu_{no\ nap}, \quad H_a : \mu_{nap} \neq \mu_{no\ nap}$$

Or equivalently:

$$H_0 : \mu_{nap} - \mu_{no\ nap} = 0, \quad H_a : \mu_{nap} - \mu_{no\ nap} \neq 0$$

Aside: This α level is equivalent to $\alpha = .05$ and then applying the [Bonferonni correction](https://en.wikipedia.org/wiki/Bonferroni_correction) (https://en.wikipedia.org/wiki/Bonferroni_correction).

Before any analysis, we will convert 'night bedtime' into decimalized time.

In [6]:

```
# Convert 'night bedtime' into decimalized time
df.loc[:, 'night bedtime'] = np.floor(df['night bedtime'])*60 + np.round(df['night bedtim
```

Now, isolate the column 'night bedtime' for those who nap into a new variable, and those who didn't nap into another new variable.

In [7]:

```
nap_bedtime = df[df['napping'] == 1]['night bedtime']  
nap_bedtime
```

Out[7]:

```
5    1235.0  
6    1260.0  
7    1321.0  
8    1224.0  
9    1278.0  
10   1185.0  
11   1218.0  
12   1222.0  
13   1226.0  
14   1228.0  
15   1246.0  
16   1243.0  
17   1202.0  
18   1190.0  
19   1218.0
```

Name: night bedtime, dtype: float64

In [8]:

```
nap_bedtime
```

Out[8]:

```
5    1235.0  
6    1260.0  
7    1321.0  
8    1224.0  
9    1278.0  
10   1185.0  
11   1218.0  
12   1222.0  
13   1226.0  
14   1228.0  
15   1246.0  
16   1243.0  
17   1202.0  
18   1190.0  
19   1218.0
```

Name: night bedtime, dtype: float64

In [9]:

```
no_nap_bedtime = df[df['napping'] == 0]['night bedtime']  
no_nap_bedtime
```

Out[9]:

```
0    1245.0  
1    1163.0  
2    1200.0  
3    1186.0  
4    1161.0
```

Name: night bedtime, dtype: float64

In [10]:

no_nap_bedtime

Out[10]:

```
0    1245.0
1    1163.0
2    1200.0
3    1186.0
4    1161.0
```

Name: night bedtime, dtype: float64

Now find the sample mean bedtime for nap and no_nap.

In [11]:

```
nap_mean_bedtime = nap_bedtime.mean()
nap_mean_bedtime
```

Out[11]:

1233.0666666666666

In [13]:

```
no_nap_mean_bedtime = no_nap_bedtime.mean()
no_nap_mean_bedtime
```

Out[13]:

1191.0

Question: What is the sample difference of mean bedtime for nappers minus no nappers?

Now find the sample standard deviation for X_{nap} and $X_{no\ nap}$.

In [15]:

no_nap_bedtime.std()

Out[15]:

34.30014577228499

In [16]:

nap_bedtime.std()

Out[16]:

34.445540177143954

	Nap Bedtime	No Nap Bedtime
Mean	1233.07	1191
Std Dev, s	34.35	34.30

Nap Bedtime No Nap Bedtime

We expect the variance in sleep time for toddlers who nap and toddlers who don't nap to be the same. Calculate the pooled standard error of $\bar{X}_{nap} - \bar{X}_{no\ nap}$.

$$s.e.(\bar{X}_{nap} - \bar{X}_{no\ nap}) = S_P \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

$$S_P^2 = \frac{S_1^2(n_1-1) + S_2^2(n_2-1)}{n_1-1 + n_2-1}$$

Question: What is the pooled s.e. $(\bar{X}_{nap} - \bar{X}_{no\ nap})$?

In [17]:

```
import math
n1 = len(df[df['napping'] == 1])
se_nap = nap_bedtime.std() / math.sqrt(n1)
se_nap
n2 = len(df[df['napping'] == 0])
se_no_nap = no_nap_bedtime.std() / math.sqrt(n2)
se_no_nap
sp = math.sqrt((((nap_bedtime.std())**2) * (n1 - 1)) + ((no_nap_bedtime.std())**2) * (n2 - 1))
pooled_se = sp * math.sqrt((1 / n1) + (1 / n2))
pooled_se
```

Out[17]:

17.77094313065816

Question: Given our sample size of n , how many degrees of freedom (df) are there for the associated t distribution?

In [18]:

```
n1 + n2 - 2
```

Out[18]:

18

Now find the t -test statistic for our hypothesis test using

- pooled s.e. $(\bar{X}_{nap} - \bar{X}_{no\ nap})$
- $\bar{X}_{nap} - \bar{X}_{no\ nap}$
- $\mu_{0, nap} - \mu_{0, no\ nap}$, the population difference in means under the null hypothesis

Question: What is the t -test statistic for the hypothesis test?

To find the p-value, we can use the function:

```
t.cdf(y, df)
```

Which for $X \sim t(df)$ returns $P(X \leq y)$.

Because of the symmetry of the t distribution, we have that

```
1-t.cdf(y, df)
```

returns $P(X > y)$

This function, `t.cdf(y, df)`, will give you the same value as finding the one-tailed probability of y on a t -table with the specified degrees of freedom.

Using the function `t.cdf` and your t -test statistics, find the p -value.

In [19]:

```
from scipy.stats import t
t_statistics = ((nap_mean_bedtime - no_nap_mean_bedtime) / pooled_se)
p_value = 1 - t.cdf(t_statistics, n-2)
print("t-statistics:", t_statistics)
print("p-value:", p_value)
```

```
t-statistics: 2.367160052079275
p-value: 0.014667451430902756
```

In [61]:

```
t-statistics : 2.367160052079275
p-value : 0.014667451430902756
```

Question: What is the p -value to the nearest hundredth?

In [20]:

```
p_value*100
```

Out[20]:

```
1.4667451430902756
```

In [62]:

```
p-value to the nearest hundredth : 1.4667451430902756
```

Calculate the t test statistics and corresponding p -value using the `scipy` function `scipy.stats.ttest_ind(a, b, equal_var=True)` and check with your answer.

In [21]:

```
from scipy.stats import ttest_ind
t_statistics2, p_value2 = ttest_ind(nap_bedtime, no_nap_bedtime, equal_var=True)
print("t-statistics:", t_statistics2)
print("p-value:", p_value2)
```

```
t-statistics: 2.367160052079275
p-value: 0.029334902861805394
```


In [60]:

Out[60]:

Ttest_indResult(statistic=2.367160052079275, pvalue=0.029334902861805394)

Does `scipy.stats.ttest_ind` return values for a one-sided or two-sided test? Can you think of a way to recover the results you got using `1-t.cdf` from the p-value given by `scipy.stats.ttest_ind`?

In [22]:

```
##two-sided
p_value*2
```

Out[22]:

0.029334902861805512

Question: Do you reject or fail to reject the null hypothesis that the difference in average bedtimes for napping versus non napping toddlers is 0?

Using the `scipy.stats.ttest_ind` function, find the t -test statistic and p-value for the second hypothesis test:
2. The average total 24 h sleep duration (in minutes) for napping toddlers is different from toddlers who don't nap.

$$H_0 : \mu_{nap} = \mu_{no\ nap}, H_a : \mu_{nap} \neq \mu_{no\ nap}$$

Or equivalently:

$$H_0 : \mu_{nap} - \mu_{no\ nap} = 0, H_a : \mu_{nap} - \mu_{no\ nap} \neq 0$$

Question: What is the t -test statistic and p-value?

In [25]:

```
# t_statistics2, p_value2 = ttest_ind(nap_bedtime, no_nap_bedtime, equal_var=True)
# print("t-statistics:", t_statistics2)
# print("p-value:", p_value2)

nonap24 = df[df['napping']==0]['24 h sleep duration']
nap24 = df[df['napping']==1]['24 h sleep duration']
ttest_ind(nonap24, nap24)
```

Out[25]:

Ttest_indResult(statistic=-1.4811248223284985, pvalue=0.1558664953018476)

Question: For $\alpha = .025$, do you reject or fail to reject the null hypothesis?

In []:

```
print('Failed to reject the null hypothesis, p-value is smaller than 0.025')
```

