

CS6210 - Homework/Assignment-5

Arnab Das(u1014840)

November 27, 2016

1: Chapter-9: Question-5

An $n \times n$ linear system of equations $Ax = b$ is modified such that every b_i is replaced with $b_i - x_i^3$.

(a) For the system $Ax = b$, thus we have:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{bmatrix}, x = \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} \text{ and } b = \begin{bmatrix} b_1 \\ b_2 \\ \dots \\ b_n \end{bmatrix}$$

The b vector gets modified as $\begin{bmatrix} b_1 - x_1^3 \\ b_2 - x_2^3 \\ \dots \\ b_n - x_n^3 \end{bmatrix}$, so that now we have a non-linear system that can be expressed as

$\mathbf{f}(\mathbf{x}) = 0$, where \mathbf{f} is a vector of functions represented as :

$$f_1(x) = (a_{11}x_1 + x_1^3) + a_{12}x_2 + \dots + a_{1n}x_n - b_1 = 0$$

$$f_2(x) = a_{21}x_1 + (a_{22}x_2 + x_2^3) + \dots + a_{2n}x_n - b_2 = 0$$

...

$$f_i(x) = a_{i1}x_1 + a_{i2}x_2 + \dots + (a_{ii}x_i + x_i^3) + \dots + a_{in}x_n - b_i = 0$$

...

$$f_n(x) = a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n + x_n^3 - b_n = 0$$

Then, the Jacobian of this vector of functions, \mathbf{f} , will be:

$$J = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} & \dots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} & \dots & \frac{\partial f_2}{\partial x_n} \\ \dots & \dots & \dots & \dots & \dots \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial f_n}{\partial x_2} & \frac{\partial f_n}{\partial x_3} & \dots & \frac{\partial f_n}{\partial x_n} \end{bmatrix}$$

$$= \begin{bmatrix} (a_{11} + 3x_1^2) & a_{12} & a_{13} & \dots & a_{1i} & \dots & a_{1n} \\ a_{21} & (a_{22} + 3x_2^2) & a_{23} & \dots & a_{2i} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & a_{i3} & \dots & (a_{ii} + 3x_i^2) & \dots & a_{in} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{ni} & \dots & (a_{nn} + 3x_n^2) \end{bmatrix} = A + D, \text{ where D is:}$$

$$D = \begin{bmatrix} 3x_1^2 & 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 3x_2^2 & 0 & \dots & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 3x_i^2 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & \dots & 3x_n^2 \end{bmatrix}$$

(b) Given that A is strictly diagonally dominant with positive elements in its diagonal. We know that a strictly diagonally dominant matrix is non-singular, hence **A is non-singular**. Now, diagonal dominance means:

$$\forall i, |a_{ii}| \geq \sum_{j=1}^n |a_{ij}|, j \neq i$$

Since, D is a diagonal matrix with non-negative elements along its diagonal, hence adding D to A , conserves the diagonal dominance in the resulting matrix J . Thus, at any iterate, the Jacobian, J , also remains diagonally dominant and hence non-singular.

(c) In Newton's method, we are theoretically basically solving the following:

$$x_{k+1} - x_k = -J(x_k)f(x_k) \quad (1)$$

where, x_k is the value of x in the k 'th iteration. For convergence of the method, we want the following:

$$\frac{\|x_{k+1} - x^*\|}{\|x_k - x^*\|} = \lambda \quad (2)$$

where, λ is some positive constant, and x^* is the optimal solution.

The error at the $(k+1)$ 'th iteration can be written as:

$$\|e_{k+1}\| = \|x_{k+1} - x^*\| = \|x_k - J(x_k)^{-1}f(x_k) - x^*\| = \|e_k - J(x_k)^{-1}f(x_k)\| \quad (3)$$

Now, from Taylor's series expansion, we have:

$$f(x_k) = f(x^*) + J(x_k)(x_k - x^*) + \frac{1}{2}(x_k - x^*)^T H(x_k)(x_k - x^*) + \dots$$

Approximating higher order terms:

$$f(x_k) = f(x^*) + J(x_k)e_k + \frac{1}{2}(e_k)^T H(x_k)(e_k)$$

At the optima, $f(x^*) = 0$. Multiplying both sides by $J(x_k)^{-1}$ thus we get:

$$J(x_k)^{-1}f(x_k) = 0 + e_k + \frac{1}{2}J(x_k)^{-1}(e_k)^T H(x_k)(e_k)$$

Replacing this in (3)

$$\|e_{k+1}\| = \|e_k - e_k - \frac{1}{2}J(x_k)^{-1}(e_k)^T H(x_k)(e_k)\| = \|\frac{1}{2}J(x_k)^{-1}(e_k)^T H(x_k)(e_k)\| \leq \frac{\|J^{-1}\|\|H\|}{2}\|e_k\|^2 \quad (4)$$

Thus (4) indicates that if the Jacobian is non-singular, then it quadratically converges. So, the only remaining piece is to show that the Jacobian is non-singular. We know that a symmetric positive definite matrix is non-singular. If we can prove that J is symmetric positive definite, then it suits our purpose: Consider any non-zero vector, u , then:

$$u^T J u = u^T (A + D)u = u^T A u + u^T D u$$

we know that A is symmetric positive definite, hence $u^T A u > 0$. So, we need to show now $u^T D u > 0$.

$$u^T D u = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} 3x_1^2 & 0 & 0 & \dots & 0 \\ 0 & 3x_2^2 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 3x_n^2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{bmatrix} = x_1^2 u_1^2 + x_2^2 u_2^2 + \dots + x_i^2 u_i^2 + \dots + x_n^2 u_n^2 > 0$$

Hence,

$$u^T J u = u^T (A + D)u = u^T A u + u^T D u > 0$$

Thus, the Jacobian is also symmetric positive definite and hence invertible and non-singular. Hence, it is guaranteed to converge as per (4).

2: Chapter-9: Question-6

(a) Given a linear system of equations, $Ax = b$, on which we apply Newton's method. For a linear system,

$$\text{the Jacobian, } J, \text{ will be : } J = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \frac{\partial f_1}{\partial x_3} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \frac{\partial f_2}{\partial x_3} & \cdots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial f_n}{\partial x_2} & \frac{\partial f_n}{\partial x_3} & \cdots & \frac{\partial f_n}{\partial x_n} \end{bmatrix}$$

$$= \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1i} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2i} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{i1} & a_{i2} & a_{i3} & \cdots & a_{ii} & \cdots & a_{in} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{ni} & \cdots & a_{nn} \end{bmatrix} = A$$

Thus, Newton's method applied to this problem becomes:

$$x^* = x_1 = x_0 + A^{-1}(b - Ax_0) = A^{-1}b$$

Hence, it will converge in a single iteration.

(b) The Newton's method is of the form:

$$J(x_k)(x_{k+1} - x_k) = -f(x_k)$$

If the Jacobian is singular at the solution, then theoretically it means $\Delta x_k = x_{k+1} - x_k$, will be potentially large in the region close to the solution which means the values of x at the k 'th and $(k+1)$ 'th iterate will have a large difference implying that convergence will be extremely slow and even the convergence might be lost, since two consecutive iterations will not result in close 'x' values that can meet the convergence limits. Also, since J is singular, hence the condition number given by $\|J\|\|J^{-1}\|$ will be infinitely large indicating an ill-conditioned system, which will not potentially converge.

7: Chapter-10: Question-25

(a) Let the quadratic polynomial be

$$p(x) = c_0 + c_1x + c_2x^2 \tag{5}$$

Then , at $x=0$:

$$p(0) = c_0 = f(0) = 1$$

and

$$p'(0) = c_1 = f'(0) = -1$$

At, $x=1$:

$$c_0 + c_1 + c_2 = f(1) = 2$$

Solving the above 3 equations, gives

$$c_0 = 1$$

$$c_1 = -1$$

$$c_2 = 2$$

Plugging these coefficients back to (5), we get the following quadratic interpolant:

$$p(x) = 1 - x + 2x^2 \tag{6}$$

By taking the first derivative and the second derivative of (6), we get:

$$p'(x) = -1 + 4x$$

$$p''(x) = 4 > 0$$

Equating the first derivative to 0, we get a stationary point at $x = \frac{1}{4}$, and since the second derivative, $p''(x) > 0$, hence this point is a minima. Thus we have found $x^* = \frac{1}{4}$. (Answer)

(b) We already found the minima $x^* = \frac{1}{4}$ which is $0 < \frac{1}{4} < 1$. Since, it is quadratic, it has only a single maxima or minima because its first derivative is linear and hence cannot have multiple roots. Since, we have found a minima, hence it has to be the unique minima satisfying $0 < x^* < 1$. For the original function, given it has continuous second order derivative, however nothing is mentioned about its higher derivatives. Now, since $f(1) > f(0)$, there is a point in $[0, 1)$, from where the function f starts to increase. Since, $f'(0) < 0$, hence $x = 0$ is not the point where f achieves a minimum. So, f achieves a minimum at $x^* = 0 + \delta$, where δ is a positive quantity between 0 and 1. Hence, f has a unique minimum at a point $0 < x < 1$. Since, f has continuous second order derivative in this region, we do not know about its higher order derivatives. If it is quadratic, then this minima is unique for same reasons as the quadratic polynomial. But, if the underlying function is not quadratic, then the uniqueness cannot be asserted since there can be many more minimas in this region.

8: Chapter-11: Question-3

Let $f \in C^3[a, b]$ be given at equidistant points, $x_i = a + ih$, where $i = 0, 1, 2, \dots, n$ and $nh = b - a$. $f'(a)$ is given as well.

(a) Construct an algorithm for C^1 continuous quadratic interpolation:

$$v(x) = S_i(x) = a_i + b_i(x - x_i) + c_i(x - x_i)^2; x_i \leq x \leq x_{i+1}$$

That is an algorithm to determine the 3n coefficients.

Algorithm: The polynomial $S_i(x)$ in the interval $x_i \leq x \leq x_{i+1}$ will match the function value exactly. So, we have our first condition as:

$$S_i(x_i) = a_i = f(x_i)$$

Thus, we have n equations in n variables due to this. This gives us all the n a_i values.

Also, since $S_i(x)$ is valid in the interval $x_i \leq x \leq x_{i+1}$, hence it should satisfy the function value at x_{i+1} as well. Thus we have our next condition as:

$$S_i(x_{i+1}) = f(x_{i+1})$$

$$a_i + b_i(x_{i+1} - x_i) + c_i(x_{i+1} - x_i)^2 = f(x_{i+1})$$

since, $x_{i+1} - x_i = h$, we get:

$$b_i h + c_i h^2 = a_{i+1} - a_i$$

This gives us n equations in 2n variables. In combination with the above we get 2n equations in 3n variables.

Next, since we want the interpolant to be C^1 continuous and the given function is C^3 , hence, the first derivative of $S_i(x_i)$ should match the first derivative of the function at x_i . Thus, we get:

$$\begin{aligned} S'_i(x_i) &= f'(x_i) \\ b_i &= f'(x_i) \end{aligned}$$

Also, the derivatives should match at x_{i+1} . Hence,

$$\begin{aligned} S'_i(x_{i+1}) &= f'(x_{i+1}) \\ b_i + 2c_i(x_{i+1} - x_i) &= f'(x_{i+1}) \\ b_i + 2c_i h &= f'(x_{i+1}) \end{aligned}$$

There is a certain pattern that emerges. If we know the derivative value at x_i , we can use it further to compute the derivative value at x_{i+1} . Since, we need to evaluate all the derivative values except the first one, let us denote the derivative with the variable d_i .

Thus, from the above relations we have the following set of equations:

n equations for

$$a_i = f(x_i)$$

n equations for

$$b_i h + c_i h^2 - (a_{i+1} - a_i) = 0$$

n equations for

$$b_i - d_i = 0$$

(n-1) equations for

$$b_i + 2c_i h - d_{i+1} = 0$$

1 equations for the derivative given at $x=a$, that is d_0

$$d_0 = f'(a)$$

Thus, in total we have $4n$ variables with $4n$ equations which can be solved to get the $3n$ coefficients $\{a_i, b_i, c_i\}$ and the derivatives d_i .

One more method: Since, we want the piecewise interpolant to be C^1 continuous, hence at the interior point x_i , the first derivative of $S_{i-1}(x)$ should equal the first derivative of $S_i(x)$. Hence, we have the following relation:

$$\begin{aligned} S'_{i-1}(x_i) &= S'_i(x_i) \\ b_{i-1} + 2c_{i-1}(x_i - x_{i-1}) &= b_i \\ b_{i-1} - b_i + 2c_{i-1}h &= 0 \end{aligned}$$

This will have (n-1) equations since we equate between two consecutive interior points.

Thus, now we can consider the following set of equations:

n equations for

$$a_i = f(x_i)$$

n equations for

$$b_i h + c_i h^2 - (a_{i+1} - a_i) = 0$$

(n-1) equations for

$$b_{i-1} - b_i + 2c_{i-1}h = 0$$

and 1 equation for the condition that the derivative at $x = a$, $f'(a)$, is given.

$$\begin{aligned} S'_0(a) &= b_0 = f'(a) \\ b_0 - f'(a) &= 0 \end{aligned}$$

Thus, now we have $3n$ equations and $3n$ variables, which can be solved to get the $3n$ coefficients $\{a_i, b_i, c_i\}$.

(b) The error estimate for an n -degree interpolate, $p_n(x)$, with respect to the original function, $f(x)$, is expressed as:

$$f(x) - p_n(x) = \frac{f^{(n+1)}(\zeta)}{(n+1)!} \prod_{i=0}^n (x - x_i)$$

For piecewise quadratic interpolant, $v(x) = S_i(x)$, we have $n=2$. Hence, we get,

$$|f(x) - v(x)| \leq (x - x_{i-1})(x - x_i) \max_{a \leq \zeta \leq b} \frac{f^3(\zeta)}{3!} \quad (7)$$

That is the 3rd derivative is bounded by its max value at a point over the entire interval $[a, b]$. For the term, $(x - x_{i-1})(x - x_i)$, the maximum occurs at the midpoint of the interval, that is, at $\frac{x_{i-1} + x_i}{2}$. Thus, we get:

$$|(x - x_{i-1})(x - x_i)| \leq \left(\frac{x_i - x_{i-1}}{2} \right)^2$$

and since, $x_i - x_{i-1} = h$, we get

$$|(x - x_{i-1})(x - x_i)| \leq \left(\frac{h}{2} \right)^2$$

Plugging this back to (7), we get the following error bound:

$$ErrorBound = |f(x) - v(x)| \leq \frac{h^2}{24} \max_{a \leq \zeta \leq b} |f^3(\zeta)|$$

9: Chapter-11: Question-12

Derive a B-spline basis representation for piecewise linear interpolation and for piecewise cubic interpolation:

For $(n+1)$ interpolating points and degree $(k-1)$, the bspline representation is written as:

$$P(u) = \sum_{i=0}^n p_i N_{i,k}(u) \quad (8)$$

where $N_{i,k}$ are the bspline basis functions and $(k-1)$ is the degree of the curve. Hence, for a piece-wise linear interpolant, the degree of the curve is 1, hence $k = 2$. Thus, we have $(n+1)$ interpolating points and $k=2$. The idea of bspline formulation is to generate curve that are affected locally by the control points. So, if a point is changed, the curve formulation will change locally, and not cause changes in the entire curve unlike bezier curves. The entire interval of points $\{p_0, p_1, \dots, p_n\}$ is divided into a sequence of knots (t_0, t_1, \dots, t_r) , such that the effect of a point will have its control in a knot region $[t_{i-1}, t_i]$ and not anywhere else: The basis function, $N_{i,k}$, is defined as :

$$N_{i,k}(u) = \frac{(u - t_i)N_{i,k-1}(u)}{t_{i+k} - t_i} + \frac{(t_{i+k} - u)N_{i+1,k-1}(u)}{t_{i+k} - t_{i+1}} \quad (9)$$

The base case of the basis functions is given as:

$$\begin{aligned} N_{i,1}(u) &= 1; t_i \leq u < t_{i+1} \\ &= 0; \text{otherwise} \end{aligned}$$

This allows for the termination condition of the recursion defined in (9):

In general, the knot values, t_i are defined as:

$$\begin{aligned} t_i &= 0; i < k \\ t_i &= i - k + 1; k \leq i \leq n \\ t_i &= n - k + 2; i > n \end{aligned}$$

For the piecewise linear case, the value of $k = 2$, hence the above relations translate to:

$$\begin{aligned} t_i &= 0; i < 2 \\ t_i &= i - 1; 2 \leq i \leq n \\ t_i &= n; i > n \end{aligned}$$

Thus, the set of $(n + k + 1) = (n + 3)$ knot values will be:

$$t = (0, 0, 1, 2, \dots, n - 1, n, n)$$

(8) in this case translates to :

$$P(u) = \sum_{i=0}^n p_i N_{i,2}(u) = p_0 N_{0,2}(u) + p_1 N_{1,2}(u) + \dots + p_i N_{i,2}(u) + \dots + p_n N_{n,2}(u)$$

where, the basis function $N_{i,2}$ in this case will be :

$$N_{i,2}(u) = \frac{(u - t_i)N_{i,1}(u)}{t_{i+2} - t_i} + \frac{(t_{i+2} - u)N_{i+1,1}(u)}{t_{i+2} - t_{i+1}} \quad (10)$$

Here, the base cases of the recursion will become:

$$N_{0,1} = 1 \text{ if } t_0 \leq u < t_1 \text{ or } 0 \leq u < 0$$

$$N_{0,1} = 0, \text{otherwise,}$$

$$N_{0,1} = 0$$

$$N_{1,1} = 1 \text{ if } t_1 \leq u < t_2 \text{ or } 0 \leq u < 1$$

$$N_{1,1} = 0, \text{otherwise}$$

$$N_{2,1} = 1 \text{ if } t_2 \leq u < t_3 \text{ or } 1 \leq u < 2$$

$$N_{2,1} = 0, \text{otherwise}$$

...

$$N_{n,1} = 1 \text{ if } t_n \leq u < t_{n+1} \text{ or } n - 1 \leq u < n$$

$$N_{n,1} = 0, \text{otherwise}$$

The above base cases plugged into the recursion of the *eqrefeq : linbasis* gives the representation for the bspline basis function for piecewise linear. (Answer).

For the case of the piecewise cubic hermite interpolant, we have a degree 3 interpolant. Hence, (k-1) is 3, thus k=4. Hence, the relation for deriving the knot values will become:

$$\begin{aligned} t_i &= 0; i < 4 \\ t_i &= i - 3; 4 \leq i \leq n \\ t_i &= n - 2; i > n \end{aligned}$$

Thus, the set of $(n + k + 1) = (n + 5)$ knot values will be:

$$t = (0, 0, 0, 0, 1, 2, \dots, n - 3, n - 2, n - 2, n - 2, n - 2)$$

(8) in this case translates to :

$$P(u) = \sum_{i=0}^n p_i N_{i,4}(u) + p_1 N_{1,4}(u) + \dots + p_i N_{i,4}(u) + \dots + p_n N_{n,4}(u) \quad (11)$$

Where the recursions over the basis functions can be further expanded following the relation(1), to reach the terminal base cases which in these case will be defined as follows:

$$\begin{aligned} N_{0,1} &= 1 \text{ if } t_0 \leq u < t_1 \text{ or } 0 \leq u < 0 \\ N_{0,1} &= 0, \text{ otherwise,} \\ \text{So, } N_{0,1} &= 0 \end{aligned}$$

$$\begin{aligned} N_{1,1} &= 1 \text{ if } t_1 \leq u < t_2 \text{ or } 0 \leq u < 0 \\ N_{1,1} &= 0, \text{ otherwise,} \\ \text{So, } N_{1,1} &= 0 \end{aligned}$$

$$\begin{aligned} N_{2,1} &= 1 \text{ if } t_2 \leq u < t_3 \text{ or } 0 \leq u < 0 \\ N_{2,1} &= 0, \text{ otherwise,} \\ \text{So, } N_{2,1} &= 0 \end{aligned}$$

$$\begin{aligned} N_{3,1} &= 1 \text{ if } t_3 \leq u < t_4 \text{ or } 0 \leq u < 1 \\ N_{3,1} &= 0, \text{ otherwise,} \end{aligned}$$

$$\begin{aligned} N_{4,1} &= 1 \text{ if } t_4 \leq u < t_5 \text{ or } 1 \leq u < 2 \\ N_{4,1} &= 0, \text{ otherwise,} \end{aligned}$$

$$\begin{aligned} N_{5,1} &= 1 \text{ if } t_5 \leq u < t_6 \text{ or } 2 \leq u < 3 \\ N_{5,1} &= 0, \text{ otherwise,} \end{aligned}$$

...

$$\begin{aligned} N_{n,1} &= 1 \text{ if } t_n \leq u < t_{n+1} \text{ or } (n - 3) \leq u < (n - 2) \\ N_{n,1} &= 0, \text{ otherwise,} \end{aligned}$$

Now consider (11), the basis functions recursively depends exactly on 4 base cases if we break them down as follows:

$$N_{0,4} : (N_{0,1}, N_{1,1}, N_{2,1}, N_{3,1})$$

$$N_{1,4} : (N_{1,1}, N_{2,1}, N_{3,1}, N_{4,1})$$

$$N_{2,4} : (N_{2,1}, N_{3,1}, N_{4,1}, N_{5,1})$$

$$N_{3,4} : (N_{3,1}, N_{4,1}, N_{5,1}, N_{6,1})$$

and so on,

In the region , $0 \leq u < 1$, only $N_{3,1} = 1$, other $N'_{i,1}$ s are zero. Now , $N_{3,1}$ is present only in the recursive list of the basis functions associated with (p_0, p_1, p_2, p_3) . Hence, in the region $0 \leq u < 1$, only these 4 points serve as the control points with local control over the curve. Other points do not have any effect here. Similarly, in the region $1 \leq u < 2$, only $N_{4,1}$ is 1, which is present in the recursive list of the basis functions associated with (p_1, p_2, p_3, p_4) . Hence, in the region $1 \leq u < 2$, only these 4 points serve as the local control points.

Thus plugging in the base cases , $N'_{i,1}$ s, present in the recursive list of respective basis functions, we obtain the representation of the bspline basis functions. (Answer).